



Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 228-406	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Riverdale DTD 8000 Lawn Weed Killer	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas Inc. 1333 Burr Ridge Parkway Suite 125A Burr Ridge, IL 60527 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAY 18 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change relative to PR Notice 2001-06. This notification is consistent with the guidance in PR Notice 2001-06 and the requirements of EPA's regulations at 40CFR156.10 and 40CFR152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 2001-06 and the requirements of 40CFR156.10 and 40CFR152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name George Meindl	Title Registration Manager	Telephone No. (Include Area Code) 708.755.7277
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		8. Date Application Received (Stamped)
2. Signature 	3. Title Registration Manager	
4. Typed Name George Meindl	5. Date 4/27/2005	

2/10

(RESIDENTIAL LABEL)

NOTIFICATION

MAY 18 2005

Riverdale®

DTDA™ 8000 LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN RESIDENTIAL TURF

CONTAINS 2,4-D, TRICLOPYR, AND DICAMBA

KILLS (OR CONTROLS): DANDELION, CHICKWEED, BLACK MEDIC, KNOTWEED, PLANTAIN, OXALIS, CLOVER, COCKLEBUR, THISTLE AND MANY OTHER SPECIES OF BROADLEAF WEEDS.

ONE QUART TREATS (COVERS) 8000 SQUARE FEET

ACTIVE INGREDIENTS:

Trisopropanolamine Salt of 2,4-D Acid*.....	12.9%
Triethylamine Salt of Triclopyr Acid**.....	1.2%
Dicamba Acid***.....	0.9%

OTHER INGREDIENTS:	<u>85.0%</u>
TOTAL:	100.0%

Isomer Specific Method. Equivalent to:

*2,4-D Acid.....	6.944%, 0.604 lbs./gal.
**Triclopyr Acid.....	0.862%, 0.075 lbs./gal.
***Dicamba Acid.....	0.862%, 0.075 lbs./gal.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-406
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

000228-00406.20050426.Res

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated contact may cause allergic reactions in some individuals. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

The maximum number of broadcast applications per treatment site is 2 per year.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on Residential Turf.

USE PRECAUTIONS

Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants.

Do not use on Centipedegrass, St. Augustinegrass, Dichondragrass, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Do not exceed recommended rates because damage to turf may occur. Shake well before using.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 24 hours, nor should lawns be irrigated (watered) for 24 hours following application. If lawn needs watering, water thoroughly before application; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder	Common chickweed	Horsetail	Prostrate spurge	Thistle
Annual yellow sweet clover	Common mullein	Indiana mallow	Prostrate vervain	Tick trefoil
Artichoke	Common sowthistle	Ironweed	Puncture vine	Toadflax
Aster	Corn Chamomile	Jewelweed	Purslane	Trailing Crownvetch
Austrian fieldcress	Creeping jenny	Jimsonweed	Ragweed	Tumble mustard
Bedstraw	Crimson clover	Kochia	Red clover	Tumble pigweed
Beggartick	Croton	Knawel	Redroot pigweed	Tumbleweed
Biden	Cudweed	Knotweed	Red sorrel	Velvet leaf
Bindweed	Curly dock	Lambsquarter	Redstem filaree	Venice mallow
Bird vetch	Curly indigo	Lespedeza	Rough cinquefoil	Veronica
Bitterweed	Dandelion	Locoweed	Rough fleabane	Vervain
Bitter wintercress	Dead nettle	Lupine	Roundleafed marigold	Vetch
Black-eyed Susan	Dock	Mallow	Rush	Virginia buttonweed
Black medic	Dogbane	Marshelder	Russian pigweed	Virginia creeper
Black mustard	Dogfennel	Matchweed	Russian thistle	Virginia pepperweed
Black-seed plantain	Elderberry	Mexicanweed	St. Johnswort	Wavyleaf bullthistle
Blessed thistle	English daisy	Milk Vetch	Scarlet pimpernel	Western clematis
Blue lettuce	Fall dandelion	Milkweed bloodflower	Scotch thistle	Western salsify
Blue vervain	False dandelion	Mugwort	Sheep sorrel	White clover
Box elder	False flax	Morningglory	Shepherdspurse	White mustard
Bracted plantain	False sunflower	Mouseear chickweed	Slender plantain	Wild mustard
Brassbuttons	Fiddleneck	Musk thistle	Smallflower galinsoga	Wild aster
Bristly oxtongue	Field bindweed	Mustard	Smartweed	Wild buckwheat
Broadleaf dock	Field pansy	Narrowleaf plantain	Smooth dock	Wild carrot
Broadleaf plantain	Flea bane (daisy)	Narrowleaf vetch	Smooth pigweed	Wild four-o'clock
Broomweed	Flixweed	Nettle	Sneezeweed	Wild garlic
Buckhorn	Florida betony	Orange hawkweed	Southern wild rose	Wild geranium
Buckhorn plantain	Florida pusley	Oxalis	Sowthistle	Wild lettuce
Bulbous buttercup	Frenchweed	Oxeye daisy	Spanishneedle	Wild marigold
Bull nettle	Galinsoga	Parsley-piert	Spatterdock	Wild onion
Bull thistle	Garlic mustard	Parsnip	Speedwell	Wild parsnip
Burdock	Goathead	Pearlwort	Spiny Amaranth	Wild radish
Burning nettle	Goatsbeard	Pennywort	Spiny cocklebur	Wild rape
Bur ragweed	Goldenrod	Pennywort	Spotted catsear	Wild strawberry
Burweed	Ground ivy	Peppergrass	Spotted knapweed	Wild sweet potato
Buttercup	Gumweed	Pepperweed	Spotted spurge	Wild vetch
Canada thistle	Hairy bittercress	Pigweed	Spurge	Willow
Carolina geranium	Hairy fleabane	Pineywoods bedstraw	Spurweed	Witchweed
Carpetweed	Hawkweed	Plains coreopsis	Stinging nettle	Woolly morningglory
Catchweed bedstraw	Healall	Plantain	Stinkweed	Woodsorrel
Catsear	Heartleaf drymary	Poison hemlock	Stitchwort	Woolly croton
Catnip	Hedge bindweed	Poison ivy	Strawberry clover	Woolly plantain
Chickweed	Hedge mustard	Poison oak	Sumac	Wormseed
Chicory	Hemp	Pokeweed	Sunflower	Yarrow
Cinquefoil	Henbit	Poorjoe	Sweet clover	Yellow rocket
Clover	Hoary cress	Povertyweed	Tall nettle	Yellowflower
Cockle	Hoary plantain	Prickly lettuce	Tail vervain	pepperweed, and
Cocklebur	Hoary vervain	Prickly sida	Tansy mustard	other broadleaf weeds.
Coffebean	Honeysuckle	Primrose	Tansy ragwort	
Coffeeweed	Hop clover	Prostrate knotweed	Tanweed	
	Horsenettle	Prostrate pigweed	Tarweed	

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-kill (or control) broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks. In the event further applications are needed, use as a spot treatment.

Apply this product at the rate of 1.6 to 2.4 pints in 4 to 50 gallons of water per 8,000 square feet (2 to 3 fluid ounces in 1/2 to 5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Ryegrass, Bent (excluding golf course greens and tees), Bahiagrass, Bermudagrass and Zoysiagrass.

Note: Some hybrid Bermudagrasses may be sensitive to this product. Contact your local Extension Service weed control specialist.

Disposable Sprayer and Bottle Instructions: Determine area to be sprayed. Apply at a rate of 32 fluid ounces (1 quart) per 8,000 square feet. Remove bottle cap and screw bottle into nozzle. Connect spray nozzle to hose while holding sprayer at waist level, point

in direction away from face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and the Lawn Weed Killer will mix automatically at the proper ratio. Walk back and forth at steady pace.

2,000 Square Feet

4,000 Square Feet

(Optional marks on left for 1 quart disposable bottle. Not to scale.)

6,000 Square Feet

8,000 Square Feet

For Refill of DTDA Spot Weed Killer: Put 2.9 fluid ounces of this product into the labeled empty quart container. Then add 29.1 fluid ounces of tap water. Trigger Sprayer - adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. Repeat as necessary.

STORAGE AND DISPOSAL

STORAGE: To be stored in original container and placed in an area inaccessible to children.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith. (RV012505)

DTDA is a Trademark of Nufarm Americas Inc.
Riverdale is a Registered Trademark of Nufarm Americas Inc.

6/10

NOTIFICATION

MAY 18 2005

(COMMERCIAL LABEL)

Riverdale®

DTDA™ 8000

LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS,
TURF GRASSES AND OTHER LISTED SITES.

CONTAINS 2,4-D, TRICLOPYR, AND DICAMBA

KILLS (OR CONTROLS): DANDELION, CHICKWEED, BLACK MEDIC, KNOTWEED, PLANTAIN,
OXALIS, CLOVER, COCKLEBUR, THISTLE AND MANY OTHER SPECIES OF BROADLEAF WEEDS.

ONE QUART TREATS (COVERS) 8000 SQUARE FEET

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-D Acid*.....	12.9%
Triethylamine Salt of Triclopyr Acid**.....	1.2%
Dicamba Acid***.....	0.9%

OTHER INGREDIENTS:	<u>85.0%</u>
TOTAL:	100.0%

Isomer Specific Method, Equivalent to:

*2,4-D Acid.....	6.944%, 0.604 lbs./gal.
**Triclopyr Acid.....	0.862%, 0.075 lbs./gal.
***Dicamba Acid.....	0.862%, 0.075 lbs./gal.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-406
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

000228-00406.20050426.Com

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated contact may cause allergic reactions in some individuals. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

The maximum number of broadcast applications per treatment site is 2 per year.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
<p align="center">HOT LINE NUMBER</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs) and similar turf areas. *Excluding Bentgrass Tees.

USE PRECAUTIONS

Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants.

Do not use on Centipedegrass, St. Augustinegrass, Dichondragrass, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Do not exceed recommended rates because damage to turf may occur. Shake well before using.



For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) avoid applying if rainfall is expected within 24 hours, nor should lawns be irrigated (watered) for 24 hours following application. If lawn needs watering, water thoroughly before application; (3) turf should not be mowed 1 to 2 days before and following application; (4) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder	Cocklebur	Hoary cress	Parsnip	St. Johnswort
Artichoke	Cocklebur, Spiny	Honeysuckle	Parsnip, Wild	Stinkweed
Aster	Coffebean	Horsenettle	Pearwort	Stitchwort
Aster, Wild	Coffeeweed	Horsetail	Pennycress	Strawberry, Wild
Bedstraw	Coreopsis, Plains	Ironweed	Pennywort	Sumac
Bedstraw, Pineywoods	Creeping jenny	Ivy, Poison	Peppergrass	Sunflower
Beggartick	Croton	Jewelweed	Pepperweed	Sunflower, False
Betony, Florida	Croton, Woolly	Jimsonweed	Pepperweed, Virginia	Sweet Potato, Wild
Biden	Cudweed	Knapweed, Spotted	Pigweed	Tansy Ragwort
Bindweed	Curly indigo	Knawel	Pigweed, Prostrate	Tanweed
Bindweed, Field	Daisy, English	Knotweed	Pigweed, Redroot	Tarweed
Bindweed, Hedge	Daisy, Oxeye	Knotweed, Prostrate	Pigweed, Russian	Thistle
Bitter wintercress	Dandelion	Kochia	Pigweed, Smooth	Thistle, Blessed
Bittercress, Hairy	Dandelion, Fall	Lambsquarter	Pigweed, Tumble	Thistle, Bull
Bitterweed	Dandelion, False	Lespedeza	Scarlet Pimpernel	Thistle, Canada
Black-eyed Susan	Dock	Lettuce, Blue	Plantain	Thistle, Musk
Bloodflower, Milkweed	Dock, Broadleaf	Lettuce, Prickly	Plantain, Black-seed	Thistle, Russian
Boxelder	Dock, Curly	Lettuce, Wild	Plantain, Bracted	Thistle, Scotch
Brassbuttons	Dock, Smooth	Locoweed	Plantain, Broadleaf	Toadflax
Bristly Oxtongue	Dogbane	Lupine	Plantain, Buckhorn	Trailing Crownvetch
Broomweed	Dogfennel	Mallow	Plantain, Hoary	Trefoil, Tick
Buckhorn	Elderberry	Mallow, Indiana	Plantain, Narrowleaf	Tumbleweed
Buckwheat, Wild	Fiddleneck	Mallow, Venice	Plantain, Slender	Velvet leaf
Bur ragweed	Fieldcress, Austrian	Marigold, Roundleafed	Plantain, Woolly	Veronica
Burdock	Filaree, Redstem	Marigold, Wild	Pokeweed	Vervain
Burweed	Flax, False	Marshelder	Poorjoe	Vervain, Blue
Buttercup	Flea bane (daisy)	Matchweed	Povertyweed	Vervain, Hoary
Buttercup, Bulbous	Fleabane, Hairy	Medic, Black	Prickly Sida	Vervain, Prostrate
Carpetweed	Fleabane, Rough	Mexicanweed	Primrose	Vervain, Tall
Carrot, Wild	Flixweed	Morningglory	Puncture vine	Vetch
Catchweed Bedstraw	Florida pusley	Morningglory, Woolly	Purslane	Vetch, Bird
Catnip	Four-O'-Clock, Wild	Mugwort	Radish, Wild	Vetch, Milk
Catsear	Frenchweed	Mullein, Common	Ragweed	Vetch, Narrowleaf
Catsear, Spotted	Galinsoga	Mustard	Rape, Wild	Vetch, Wild
Chamomile, Corn	Galinsoga, Smallflower	Mustard, Black	Rose, Southern Wild	Virginia Buttonweed
Chickweed	Garlic, Wild	Mustard, Garlic	Rush	Virginia Creeper
Chickweed, Common	Geranium, Carolina	Mustard, Hedge	Salsify, Western	Wavyleaf Bullthistle
Chickweed, Mousear	Geranium, Wild	Mustard, Tansy	Shepherdspurse	Willow
Chicory	Goathead	Mustard, Tumble	Smartweed	Witchweed
Cinquefoil	Goatsbeard	Mustard, White	Sneezeweed	Woodsorrel
Cinquefoil, Rough	Goldenrod	Mustard, Wild	Sorrel, Red	Wormseed
Clematis, Western:	Ground-Ivy	Nettle	Sorrel, Sheep	Yarrow
Clover	Groundsel	Nettle, Bull	Sowthistle	Yellow rocket
Clover, Annual Yellow	Gumweed	Nettle, Burning	Sowthistle, Common	Yellowflower
Sweet	Hawkweed	Nettle, Dead	Spanishneedle	Pepperweed
Clover, Crimson	Hawkweed, Orange	Nettle, Stinging	Spatdock	and other broadleaf
Clover, Hop	Healall	Nettle, Tall	Speedwell	weeds
Clover, Red	Heartleaf drymary	Oak, Poison	Spiny Amaranth	
Clover, Strawberry	Heathaster	Onion, Wild	Spurge	
Clover, Sweet	Hemlock, Poison	Oxalis	Spurge, Prostrate	
Clover, White	Hemp	Pansy, Field	Spurge, Spotted	
Cockle	Henbit	Parsley-piert	Spurweed	

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-kill (or control) broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks. In the event further applications are needed, use

as a spot treatment.

Apply this product at the rate of 1.6 to 2.4 pints in 4 to 50 gallons of water per 8,000 square feet (2 to 3 fluid ounces in 1/2 to 5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Ryegrass, Bentgrass (excluding golf course greens and tees), Bahiagrass, Bermudagrass and Zoysiagrass.

Note: Some hybrid Bermudagrasses may be sensitive to this product. Contact your local Extension Service weed control specialist.

Disposable Sprayer and Bottle Instructions: Determine area to be sprayed. Apply at a rate of 32 fluid ounces (1 quart) per 8,000 square feet. Remove bottle cap and screw bottle into nozzle. Connect spray nozzle to hose while holding sprayer at waist level, point in direction away from face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and the Lawn Weed Killer will mix automatically at the proper ratio. Walk back and forth at steady pace.

2,000 Square Feet

4,000 Square Feet

(Optional marks on left for 1 quart disposable bottle. Not to scale.)

6,000 Square Feet

8,000 Square Feet

Roadsides (including aprons and guard rails), rights-of-way, drainage ditchbanks and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 0.8 to 2.6 pints of this product per 5 to 30 gallons of water. This mixture will cover 4,350 square feet. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (Spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.

For Refill of DTDA Spot Weed Killer: Put 2.9 fluid ounces of this product into the labeled empty quart container. Then add 29.1 fluid ounces of tap water. Trigger Sprayer - adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. Repeat as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well-ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities by burning. If burned, stay out of smoke.

NOTICE: Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith. (RV012505)

DTDA is a Trademark of Nufarm Americas Inc.
Riverdale is a Registered Trademark of Nufarm Americas Inc.





10/10
Nufarm Americas, Inc.
George Meindl
Registration Manager
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527
Phone: 708.755-7277 Fax: 708.754.0314
george.meindl@us.nufarm.com

April 27, 2005

Via Overnight Courier

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U. S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1801 Bell Street
Arlington, VA 22202-4501

Subject: Riverdale® DTDA 8000 Lawn Weed Killer
EPA Reg. No. 228-406
Label notification per PRN 2001-06

Dear Ms. Hobgood:

Nufarm Americas Inc. has removed the statement "Partially filled container may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash" from the Storage and Disposal section of the subject registration's labeling. This can be done as a notification per PRN2001-06.

To process this request please find enclosed the following:

- Application for Pesticide Registration (EPA form 8570-1)
(which includes the PRN2001-06 certification statement)
- Final print labeling (1 copy "clean")
- Label highlighting the changes (1 copy)

If you should have any questions regarding this matter, please feel free to contact me at 708.755.7277 or email at george.meindl@us.nufarm.com.

Sincerely,

George Meindl
Registration Manager
Nufarm Americas, Inc.

