Please read instructions on reverse before completing form.	Form Approved. OMB No. 2070-0060
SEPA Environmental Protection Washington, DC 204	Registration Amendment OPP Identifier Number Amendment
Application	on for Pesticide - Section I
1. Company/Product Number 228-422	2. EPA Product Manager Joanne Miller 3. Proposed Classification X None Restricted
4. Company/Product (Name) Riverdale Battieship 2 Selective Herbicide	PM# 23
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas Inc. 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527	6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.
Check if this is a new address	Product Name
	Section - il
Amendment - Explain below. Resubmission in response to Agency letter dated	Final printed labels in repsonse to Agency letter dated "Me Too" Application. MAY 1 8 2005 Other - Explain below.
confidential statement of formula of this product. I understand that it is a	iffect cold storage instructions. EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the violation of 18 U.S.C. Sec. 1001 to wilfully make any false statement to EPA. I further votice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject
	Section - III
Meterial This Product Will Be Peckaged In:	
Child-Resistent Peckaging Yes No **Certification must be submitted Unit Peckaging Yes Ves Void Peckaging Void Peckaging wgt. No. per Unit Peckaging wgt.	Water Soluble Packaging Yes No If "Yes" No. per Package wgt container 2. Type of Container Metal Plastic Glass Paper Other (Specify)
3. Location of Net Contents Information 4. Size(s) Reta	5. Location of Label Directions
6. Manner in Which Lebel is Affixed to Product Lithogr.	dued
	Section - IV
1. Contact Point (Complete items directly below for identification	of individual to be contacted, if necessary, to process this application.)
	Title Telephone No. (Include Arer Code) VP, Registrations & Regulatory Affairs 708 755 7232
Certificat I certify that the statements I have made on this form and I acknowledge that any knowlingly false or misleading stat both under applicable law.	all attachments thereto are true, accurate and complete. Received
2. Signature Smith	VP, Registrations & Regulatory Affairs
	5. Date
Jeannie Smith	May 5, 2005

NOTIFICATION
MAY 1 & 2005

RIVERDALE®

BATTLESHIP®2 SELECTIVE HERBICIDE

FOR THE CONTROL OF ANNUAL AND PERENNIAL BROADLEAF WEEDS IN ORNAMENTAL TURF INCLUDING SOD FARMS

CONTAINS MCPA, FLUROXYPYR, AND TRICLOPYR

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Sait of 2-Methyl-4-Chlorophenoxyacetic Acid*	37.84%
1-Methylheptyl Ester of Fluroxypyr: ((4-amino-3-5-dichloro-6-fluoro- 2-pyridinyl)oxy)acetic Acid, 1-methylheptyl ester**	8.90%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid***	3.45%
OTHER INGREDIENTS:	49.81%
TOTAL	00.00%

By Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	30.90%, 2.85 lbs./gal.
**((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid	. 6.18%, 0.22 lbs./gal.
***3.5.6-Trichloro-2-Pyridinyloxyacetic Acid	. 2.47%, 0.57 lbs./gal.

Intended For Use By Professional Turf Maintenance Personnel, Landscaping or Commercial Applicators Only.

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO.228-422 EPA EST. NO. 228-IL-1 MANUFACTURED BY NUFARM AMERICAS INC. BURR RIDGE, IL 60527-0866

NET CONTENTS

GALS.

000228-00422.20050505.notif



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

NON-WPS TURF USE: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170)-in general only agricultural plant uses are covered by the WPS-must wear eye protection (goggles, face shield, or safety glasses) when using this product. It is recommended that safety glasses include front, brow, and temple protection.

WPS SOD FARM USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170)-in general, agricultural plant uses are covered-must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: If the mechanical system is used in a manner that meets the requirements fisted in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- · Wash hands, face and arms with soap and water before eating, smoking, drinking, or using the toilet.
- · Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing,
- Remove PPE immediately after handling this product.
- · Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

Probable mucosal damage may contraindicate use of gastric lavage.

FIRST AID		
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
IF SWALLOWED	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 	
IF ON SKIN OR CLOTHING	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. 	
	HOT LINE NUMBER Intainer or label with you when calling a poison control center or doctor, or going for treatment. In 1-877-325-1840 for emergency medical treatment information.	
	NOTE TO PHYSICIAN	

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spille. Percent of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT, USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTORS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for

•::::

pesticide regulation.

This product should be used strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposure.

Under some conditions, this product may have a potential to runoff to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no-till, limited till and contour plowing; these methods also reduce pesticide runoff. Where feasible, use application techniques such as T-banding, and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetaion filter strips is recommended along rivers, creeks, streams, wetlands, etc. or on the downhilf side of fields where runoff could occur to minimize water runoff.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and for high temperatures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application Do not enter treatment areas until spray has dried.

This product is a broad-spectrum herbicide for control of actively growing annual and perennial broadleaf weeds. It is for use on established Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields, Roadsides and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas; also for use on sod farms. Do not use on Golf Greens or Tees. This product may be applied to the following established turfgrass species:

ESTABLISHED COOL SEASON TURFGRASSES		ESTABLISHED WARM SEASON TURFGRASSES†	
Common Name	Scientific Name	Common Name	Scientific Name
Bentgrass	Agrostis species	Bahiagrass	Paspalum notatum
Bluegrass, Kentucky	Poa pratensis	Bermudagrass	Cynodon dactylon
Fescue, chewing	Festuca rubravar, commutata	Buffalograss	Buchloe dactyloides
Fescue, creeping red	Festuca rubra	Centipedegrass	Eremochloa ophiuroides
Fescue, sheeps	Festuca ovina	Zoysiagrass	Zoysia japonica
Fescue, tall	Festuca arundinaceae	Zoysiagrass	Zoysia tenuifolia
Ryegrass, perennial	Lolium perenne	Fescue, tall (growing in warm season areas)	Festuca arundinacea

†Do not treat warm season turfgrass with this product when mowing height is less than 1/2 inch.

Note: This product may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply this product only to healthy, well-established turf that has a well-anchored root system. Do not apply this product to grass species not listed on this label unless potential turf injury can be tolerated.

USE PRECAUTIONS

Do not apply this product directly to, or otherwise permit drift or spray mist to come into contact with cotton, citrus, grapes, tobacconvegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not pour spray solutions flear desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Spray drift can be reduced by spraying when wind velocity is low, keeping spray boom as low as possible, using nozzles that deliver larger spray droplets and minimize the production of fine spray particles, and by using higher spray volumes and lower pressures. Applications should be made only when hazards from spray drift are at a minimum. Very small quantities of spray, which may not be visible, may seriously figure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injusted. Do not apply to exposed roots of shallow rooted trees and shrubs. Maximum kill of weeds will be obtained from spring or early fall applications.



when weeds are actively growing. Do not apply more than 4 pints per acre per application. Mow newly seeded turf at least twice before treatment with this product.

For optimum results: (1) Add a spray adjuvant, such as DYNE-AMIC for best results; (2) Use a water conditioning agent, such as QUEST for high PH and water hardness; (3) Avoid applying during excessively dry or hot periods unless irrigation is used; (4) Turf should not be moved 1 to 2 days before and following application; (5) Reseed no sooner than 3 to 4 weeks after application of this

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Creeping ienny Crimson clover Annual yellow sweet clover Artichoke Croton Cudweed Aster Austrian fieldcress Curly dock Redstraw Curly indigo Beggartick Dandelion Dead nettle Biden Bindweed Dock Bird vetch Dogbane Bitterweed Dogfennel Ritter wintercress Elderberry Black-eyed Susan English daisy Black medic Fall dandelion Black mustard False dandelion Black-seed plantain False flax Blessed thistle False sunflower Blue lettuce Fiddleneck Field bindweed Blue vervain Box elder Field pansy Bracted plantain Flea bane (daisy) Brassbuttons Ffixweed Bristly oxtongue Florida betony Broadleaf dock Florida pustey Broadleaf plantain Frenchweed Broomweed Galinsoga Buckhorn Garlic mustard Buckhorn plantain Goathead Bulbous buttercup Goatsbeard Bull nettle Goldenrod **Bull thistle** Ground ivy Burdock Gumweed Burning nettle Hairy bittercress Hairy fleabane Bur ragweed Burweed Hawkweed Healall Buttercup Canada thistle Heartleaf drymary Hedge bindweed Carolina geranium Hedge mustard Carpetweed Catchweed bedstraw Hemp Henbit Catsear Catnip Hoary cress Hoary plantain Chickweed Chicory Hoary vervain Honeysuckle Cinquefoil Hop clover Clover Cockle Horsenettle Cocklebur Horsetail Coffeebean Indiana mallow Ironweed Coffeeweed Common chickweed Jewelweed Common mullein Jimsonweed Common sowthistle Kochia Knawel Corn Chamomile

Knotweed Lambsquarter Lespedeza Locoweed Lupine Mallow Marshelder Matchweed Mexicanweed Milk vetch Milkweed bloodflower Mugwort Morningglory Mousear chickweed Musk thistle Mustard Narrowleat plantain Narrowleaf vetch Nettle Orange hawkweed Oxalis Oxeye daisy Parsley-piert Parsnip Pearlwort Pennycress Pennywort Peppergrass Pepperweed Pigweed Pineywoods bedstraw Plains coreopsis Plantain Poison hemlock Poison ivv Poison oak Pokeweed Poorioe Povertyweed Prickly lettuce Prickly sida Primrose Prostrate knotweed Prostrate pigweed Prostrate source Prostrate vervain

Rough fleabane Roundleafed marigold Rush Russian pigweed Russian thistle St. Johnswort Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smartweed Smooth dock Smooth pigweed Sneezeweed Southern wild rose Sowthistle Spanishneedle Spatterdock Speedwell Spiny Amaranth Spiny cocklebur Spotted catsear Spotted knapweed Spotted spurge Spurge Spurweed Stinging nettle Stinkweed Stitchwort Strawberry clover Sumac Sunflower Sweet clover Tall nettle Tall vervain Tansy mustard Tansy ragwort Tanweed Tarweed Thistle Tick trefoil Toadflax Trailing Crownvetch Tumble mustard Tumble pigweed Tumbleweed Velvet leaf. Venice mallow

Veronica

Vervain

Vetch

Puncture vine

Purstane

Ragweed

Red clover

Red sorrel

Redroot pigweed

Redstem filaree

Rough cinquetoil

Virginia buttonweed Wild aster Wild parsnip Woodsorrel Virginia creeper Wild buckwheat Wild radish Wooly croton Virginia pepperweed Wild carrot Wild rape Wooly plantain Wavyleaf bullthistle Wild four-o'-clock Wild strawberry Wormseed Western clematis Wild gartic Wild sweet potato Yarrow Western salsify Wild geranium Wild vetch Yellow rocket White clover Wild lettuce Willow Yellowflower pepperweed White mustard Wild marigold Witchweed and other broadleaf weeds. Wild mustard Wild onion Wooly morningglory

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks.

TURF APPLICATION DIRECTIONS

Apply 3 to 4 pints of this product per acre (1.1 to 1.5 ozs./1,000 sq.ft.) to control above listed broadleaf weeds.

The maximum use rate for most warm season grasses is 3 pints per acre.

Make application using equipment that will ensure uniform coverage (see specific application directions below). Foliar sprays should be applied during warm weather when weeds are actively growing. Application under drought conditions may provide less than desirable results. When hard-to-control species are prevalent, when applications are made in late summer, when weeds are mature, or during drought conditions, use the higher rates.

On closely mowed golf course fairway bentgrass, apply this product at a maximum rate of 2 pints in 20 to 240 gallons of water per acre (0.73 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet). During conditions which promote turf stress use lower rates, 1 to 1.5 pints in 20 to 240 gallons of water per acre (0.36 to 0.55 fluid ounces in 0.3 to 5.5 gallons per 1,000 square feet). Slight turf yellowing will disappear after one week. To minimize grass injury, a second application should not be made for at least 4 weeks. Avoid swath overlaps. Do not use on golf course greens or tees.

Note: Broadleaf weeds germinate at different times of the year. Only emerged weeds present at the time of application will be controlled by this product.

Standard Volume Broadcast Applications: Apply 3 to 4 pints of this product in enough water to deliver 20 to 200 gallons of total spray mix per acre (0.5 to 5 gallons spray per 1,000 square feet). Higher application volumes may be used when this product is tank mixed with liquid fertilizers.

Low Volume Broadcast Applications: Apply 3 to 4 pints of this product in enough water to deliver from 5 to 40 gallons of total spray mixture per acre. Use low pressures and application equipment capable of delivering a uniform droplet size that can wet the weed leaf surface.

Targeted Zone Applications or Spot Treatment Using Portable Sprayers: Mix 0.5 to 1.0 ounce of this product in enough water to make one gallon of spray and apply at any time broadleaf weeds are actively growing. Target plants should be sprayed to wet but not beyond point of runoff.

Controlled Droplet Applicators (CDA), Atomizers, and Spinning Disk Applicators: Use this product at the rate of 3 to 4 pints per acre (1.1 to 1.5 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

Note: Follow CDA equipment spray instructions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always store pesticides in a secured warehouse or storage building. Protect from freezing. Store at temperatures above 25°F. If allowed to freeze, remix before using; this does not alter the product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater if these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL FOR NON-REFILLABLE CONTAINERS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings which have been opened during use and replaced all caps. Contact Nufarm Customer Service at 1-800-345-3330, to arrange for return of the empty refillable container.

WARRANTY


Seller warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions



under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use, including the use of this product on turf species not recommended on this label.

(RV050505)

Riverdale is a Registered Trademark of Nufarm Americas Inc. Battleship is a Registered Trademark of Helena Chemical Company





Nufarm Americas Inc.
Jeannie K. Smith
Vice President
Registrations & Regulatory Affairs
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527
Phone: 708.755-7332 Fax: 708.754.0314
jeannie.smith@us.nufarm.com

May 5, 2005

Via Overnight Courier

Document Processing Desk (NOTIF)
Office of Pesticide Programs 7504C
US Environmental Protection Agency
1801 South Bell Street
Arlington, VA 22202-4501

Subject:

Riverdale® Battleship® 2 Selective Herbicide

EPA Reg. No.: 228-422

Enclosed are the following items to support a notification amendment.

1. EPA Application Form 8570-1

2. Two (2) Copies of Amended Labeling

The pesticide storage statement in the storage and disposal section was updated to reflect cold storage.

Let me know if you have any questions or concerns. Thank you.

Regards,

Jeannie K. Smith Vice President

Registrations & Regulatory Affairs