



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

228-442

Date of Issuance:

MAY 26 2005

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Riverdale® Escalade™ II

Name and Address of Registrant (include ZIP Code):

Nufarm Americas, Inc.
Riverdale - A Nufarm Company
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527-0866

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit, within one (1) year of the date of this registration notice, storage stabilities (830.6317) and corrosion characteristic (830.6320) studies to the Agency.
3. Make the following labeling changes:

Signature of Approving Official:

Joanne J. Miller

Date:

MAY 26 2005

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 228-442".
 - b. Under the heading entitled "**PRECAUTIONARY STATEMENTS**" on the page two (2) of your product labeling, please incorporate the following additional precautionary statement: "Avoid contact on skin, eyes or clothing."
4. Submit one (1) copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosures

RIVERDALE® ESCALADE™ II

FOR SELECTIVE BROADLEAF WEED CONTROL
IN ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS 2,4-D, FLUROXYPYR AND DICAMBA

CONTROLS: Black medic, Chickweed, Clover, Cocklebur, Dandelion, Knotweed, Oxalis, Plantain, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	39.53%
1-Methylheptyl Ester of Fluroxypyr:[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid,1-methylheptyl Ester**	5.90%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	4.10%

OTHER INGREDIENTS:

.....	50.47%
TOTAL	100.00%

By Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	32.83%, 3.2 lbs./gal.
**[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid	4.10%, 0.4 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	4.10%, 0.4 lbs./gal.

**KEEP OUT OF REACH OF CHILDREN
WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-UUE
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

000228-00UUE.20050511.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

MAY 26 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-442

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING - AVISO**

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

NON-WPS TURF USES: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) (in general only agricultural plant uses are covered by the WPS) must wear face shield, goggles, or safety glasses and long pants, long-sleeved shirt, socks, shoes and chemical-resistant gloves when mixing, loading or applying this product. It is recommended that safety glasses include front, brow, and temple protection. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical-resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

NON-WPS INDUSTRIAL USES: When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield, goggles, or safety glasses and chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow, and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

WPS USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) (in general, agricultural plant uses are covered) must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear, when cleaning equipment, mixing or loading. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned. If this container is over one gallon and less than five gallons, mixers and loaders who do not use a mechanical system (such as a probe and pump or spigot) to transfer contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Engineering Controls Statements: If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This pesticide should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposure.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Where feasible, use application techniques such as T-banding and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur will minimize water run-off is recommended.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear and chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs) and similar turf areas; also for use on Sod Farms.

*Excluding Bentgrass Tees

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Centipede, St. Augustine, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

Alder	Bulbous buttercup	Curly dock	Hawkweed	Milkweed bloodflower
Annual yellow sweet clover	Bull nettle	Curly indigo	Healall	Mugwort
Artichoke	Bull thistle	Dandelion	Heartleaf drymary	Morningglory
Aster	Burdock	Dead nettle	Hedge bindweed	Mouseear chickweed
Austrian fieldcress	Burning nettle	Dock	Hedge mustard	Musk thistle
Bedstraw	Bur ragweed	Dogbane	Hemp	Mustard
Beggartick	Burweed	Dogfennel	Henbit	Narrowleaf plantain
Biden	Buttercup	Elderberry	Hoary cress	Narrowleaf vetch
Bindweed	Canada thistle	English daisy	Hoary plantain	Nettle
Bird vetch	Carolina geranium	Fall dandelion	Hoary vervain	Orange hawkweed
Bitterweed	Carpetweed	False dandelion	Honeysuckle	Oxalis
Bitter wintercress	Catchweed bedstraw	False flax	Hop clover	Oxeye daisy
Black-eyed Susan	Catsear	False sunflower	Horsenettle	Parsley-piert
Black medic	Catnip	Fiddleneck	Horsetail	Parsnip
Black mustard	Chickweed	Field bindweed	Indiana mallow	Pearlwort
Blackseed pantain	Chicory	Field pansy	Ironweed	Pennycress
Blackseed plantain	Cinquefoil	Flea bane (daisy)	Jewelweed	Pennywort
Blessed thistle	Clover	Flixweed	Jimsonweed	Peppergrass
Blue lettuce	Cockle	Florida betony	Kochia	Pepperweed
Blue vervain	Cocklebur	Florida pusley	Knawel	Pigweed
Box elder	Coffebean	Frenchweed	Knotweed	Pineywoods bedstraw
Bracted plantain	Coffeeweed	Galinsoga	Lambsquarter	Plains coreopsis
Brassbuttons	Common chickweed	Garlic mustard	Lespedez	Plantain
Bristly oxtongue	Common mullein	Goathead	Locoweed	Poison hemlock
Broadleaf dock	Common sowthistle	Goatsbeard	Lupine	Poison ivy
Broadleaf plantain	Corn Chamomile	Goldenrod	Mallow	Poison oak
Broomweed	Creeping jenny	Ground ivy	Marshelder	Pokeweed
Buckhorn	Crimson clover	Gumweed	Matchweed	Poorjoe
Buckhorn plantain	Croton	Hairy bittercress	Mexicanweed	Povertyweed
	Cudweed	Hairy fleabane	Milk vetch	

(con't.)

WEEDS CONTROLLED (continued)

Prickly lettuce	Scotch thistle	Stinkweed	Vervain	Wild rape
Prickly sida	Sheep sorrel	Stitchwort	Vetch	Wild strawberry
Primrose	Shepherdspurse	Strawberry clover	Virginia buttonweed	Wild sweet potato
Prostrate knotweed	Slender plantain	Sumac	Virginia creeper	Wild vetch
Prostrate pigweed	Smallflower galinsoga	Sunflower	Virginia pepperweed	Willow
Prostrate spurge	Smartweed	Sweet clover	Wavyleaf bullthistle	Witchweed
Prostrate vervain	Smooth dock	Tall nettle	Western clematis	Woolly morningglory
Puncture vine	Smooth pigweed	Tall vervain	Western salsify	Woodsorrel
Purslane	Sneezeweed	Tansy mustard	White clover	Woolly croton
Ragweed	Southern wild rose	Tansy ragwort	White mustard	Woolly plantain
Red clover	Sowthistle	Tanweed	Wild mustard	Wormseed
Redroot pigweed	Spanishneedle	Tarweed	Wild aster	Yarrow
Red sorrel	Spatdock	Thistle	Wild buckwheat	Yellow rocket
Redstem filaree	Speedwell	Tick trefoil	Wild carrot	Yellowflower
Rough cinquefoil	Spiny Amaranth	Toadflax	Wild four-o'clock	pepperweed
Rough fleabane	Spiny cocklebur	Trailing Crownvetch	Wild garlic	and other broadleaf
Roundleaved marigold	Spotted catsear	Tumble mustard	Wild geranium	weeds.
Rush	Spotted knapweed	Tumble pigweed	Wild lettuce	
Russian pigweed	Spotted spurge	Tumbleweed	Wild marigold	
Russian thistle	Spurge	Velvet leaf	Wild onion	
St. Johnswort	Spurweed	Venice mallow	Wild parsnip	
Scarlet pimpernel	Stinging nettle	Veronica	Wild radish	

Some of these species may require repeat spot applications and/or use of higher rate recommended on this product label even under ideal conditions for application.

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.75 to 1.1 fl. oz. in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Ryegrass, Bentgrass (excluding golf course greens and tees), Bahia, Bermudagrass and Zoysia.

Note: On closely mowed golf course fairway bentgrass, apply this product at a maximum rate of 2 pints in 20 to 240 gallons of water per acre (0.75 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 sq. ft.). During conditions which promote turf stress use lower rates, 1 to 1-1/2 pints in 20 to 240 gallons of water, per acre (0.36 to 0.55 fluid ounces in 0.3 to 5.5 gallons per 1,000 sq. ft.). Slight turf yellowing will disappear after one week. To minimize grass injury, a second application should not be made for at least 4 weeks. Avoid swath overlaps. Do not use on golf course greens or tees.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use this product at the rate of 2 to 3 pints per acre (0.75 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE: For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

SOD FARMS

This product is intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

Timing

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to Floratam St. Augustinegrass.

The application of this herbicide to grass seedlings is not recommended until after the second mowing.

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant Bermudagrass, dormant Zoysiagrass and dormant Bahiagrass are suggested.

Recommended Rates for Sod Farms

For Bluegrass, Ryegrass and Fescue, apply 2 to 3 pints per acre. For Creeping Bentgrass use 1.25 pints per acre. For Common and Hybrid Bermudagrass, Bahia, Zoysia and St. Augustine grass apply 1.5 to 1.8 pints per acre. Spray volume for Sod Farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

8/9

NON-TURF AREAS

Roadsides (including aprons and guardrails), rights-of-way, and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 2 to 5 pints of this product per 50 to 300 gallons of water. This mixture will cover 1 acre. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 2-1/4 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 4 to 5 pints of this product to 100 gallons of water applying 200 to 600 gallons of spray mixture per 43,500 square feet depending upon the height and thickness of the brush. Mix thoroughly before spraying.

COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Coop.'s Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding the product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Protect from freezing. Store at temperatures above 25°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings and replace all caps. Contact Nufarm Customer Service at 1-800-345-3330, to arrange for return of the empty refillable container.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Nufarm or Seller. All such risks shall be assumed by Buyer

9/9

and User, and Buyer and User agree to hold Nufarm and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. NUFARM MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or Nufarm, and buyer assumes the risk of any such use.

To the extent allowed by law, Nufarm or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF NUFARM AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF NUFARM OR SELLER, THE REPLACEMENT OF PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

(RV022305)

Riverdale is a Registered Trademark of Nufarm Americas Inc.
Escalade is a Trademark of Nufarm Americas Inc.