

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 6, 2014

Matthew Granahan
Regulatory Manager
Nufarm Americas Inc.
11901 S. Austin Ave.
Alsip IL 60477

Subject: Label Amendment – Revising Restrictions etc.
Product Name: Turnpower 3 Selective Herbicide
EPA Registration Number: 228-445
Application Date: August 4, 2014
Decision Number: 494300

Dear Mr. Granahan:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Shanta Adeeb by phone at 703-347-0502, or via email at adeeb.shanta@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn V. Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure


TRUPOWER³

SELECTIVE HERBICIDE

[Alternate brand name: Change Up Selective Herbicide]

**FOR SELECTIVE BROADLEAF WEED CONTROL IN
ORNAMENTAL LAWNS AND TURF GRASSES.**

CONTAINS MCPA, FLUROXYPYR AND DICAMBA



CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	51.05%
1-Methylheptyl Ester of Fluroxypyr:[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid, 1-methylheptyl ester**	6.00%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	4.17%
OTHER INGREDIENTS	38.78%
TOTAL:	100.00%

Isomer Specific Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	41.68%, 4.0 lbs./gal.
**[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic acid	4.17%, 0.4 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	4.17%, 0.4 lbs./gal.

DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840**

EPA REG. NO. 228-445
EPA EST. NO. _____

Manufactured for
NUFARM AMERICAS INC.
11901 S. Austin Avenue
Alsip, IL 60803



NET CONTENTS: GAL. (Liters)

[Designation as "NONREFILLABLE" or "REFILLABLE" for containers >5gal]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING / AVISO: Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistant selection chart.

Mixers, loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- chemical-resistant gloves when mixing, loading, or using any hand-held equipment
- protective eyewear (goggles, face shield, or safety glasses)

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS	
Users Should:	<ul style="list-style-type: none">• Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish, aquatic invertebrates and aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS.

Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Aerial application is prohibited.

This pesticide must be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposure. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Where feasible, use application techniques such as T-banding and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur will minimize water run-off is recommended.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore, the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls worn over short-sleeved shirt and short pants
- chemical-resistant footwear plus socks
- chemical-resistant gloves made of any waterproof material
- chemical-resistant headgear for overhead exposure, and
- protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter or allow others to enter the treated area until sprays have dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs); also for use on Sod Farms.

*Excluding Bentgrass Tees.

SPRAY DRIFT MANAGEMENT

Preventing spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not use on Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur. When treating Carpetgrass and St. Augustinegrass, avoid broadcast applications when air temperature exceeds 80°F. When air temperature exceeds 80°F limit application to spot treatment only.

APPLICATION RESTRICTIONS

- Do not apply more than 3 pints of this product per application (1.5 lbs MCPA ae, 0.15 lbs Fluroxypyr ae, 0.15 lbs Dicamba ae).
- Do not apply more than 2 applications of this product to the same treatment site per year.
- The minimum retreatment interval for this product is 21 days.
- Do not apply this product to improved varieties of St. Augustinegrass including Bitterblue, Floratam and Seville.
- Do not apply this product to lawn or turf during spring transition (green-up).
- Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species.
- Do not apply to newly seeded grasses until well established.
- Do not apply by air.

ACTIVE INGREDIENT APPLICATION RESTRICTIONS

- Application of MCPA is limited to 3.0 lbs ae per acre per year, no more than 1.5 lbs ae per application, and no more than 2 applications per year.
- Application of Fluroxypyr is limited to 0.5 lbs ae per acre per year
- Application of Dicamba is limited to 2.0 lbs ae per acre per year, no more than 1.0 lbs ae per application, and no more than 2 applications per year.

The suitable use of this product on non-labeled turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder	Galinsoga	Povertyweed	Vervain
Annual yellow sweet clover	Garlic mustard	Prickly lettuce	Vetch
Artichoke	Goathead	Prickly sida	Virginia buttonweed
Aster	Goatsbeard	Primrose	Virginia creeper
Austrian fieldcress	Goldenrod	Prostrate knotweed	Virginia pepperweed
Bedstraw	Ground ivy	Prostrate pigweed	Wavyleaf bullthistle
Beggartick	Gumweed	Prostrate spurge	Western clematis
Biden Bindweed	Hairy bittercress	Prostrate vervain	Western salsify
Bird vetch	Hairy fleabane	Puncture vine	White clover
Bitterweed	Hawkweed	Purslane	White mustard
Bitter wintercress	Healall	Ragweed	Wild mustard
Black-eyed Susan	Heartleaf drymary	Red clover	Wild aster
Black medic	Hedge bindweed	Redroot pigweed	Wild buckwheat
Black mustard	Hedge mustard	Red sorrel	Wild carrot
Black-seed plantain	Hemp	Redstem filaree	Wild four-o'clock
Blessed thistle	Henbit	Rough cinquefoil	Wild garlic
Blue lettuce	Hoary cress	Rough fleabane	Wild geranium
Blue vervain	Hoary plantain	Roundleafed	Wild lettuce
Box elder	Hoary vervain	marigold	Wild marigold
Bracted plantain	Honeysuckle	Rush	Wild onion
Brassbuttons	Hop clover	Russian pigweed	Wild parsnip
Bristly oxtongue	Horsenettle	Russian thistle	Wild radish
Broadleaf dock	Horsetail	St. Johnswort	Wild rape
Broadleaf plantain	Indiana mallow	Scarlet pimpernel	Wild strawberry
Broomweed	Ironweed	Scotch thistle	Wild sweet potato
Buckhorn	Jewelweed	Sheep sorrel	Wild vetch
Buckhorn plantain	Jimsonweed	Shepherdspurse	Willow
Bulbous buttercup	Kochia	Slender plantain	Witchweed
Bull nettle	Knawel	Smallflower	Woolly morningglory
Bull thistle	Knotweed	galinsoga	Woodsorrel
Burdock	Lambsquarter	Smartweed	Woolly croton
Burning nettle	Lespedeza	Smooth dock	Woolly plantain
Bur ragweed	Locoweed	Smooth pigweed	Wormseed
Burweed	Lupine	Sneezeweed	Yarrow
Buttercup	Mallow	Southern wild rose	Yellow rocket
Canada thistle	Marshelder	Sowthistle	Yellowflower
Carolina geranium	Matchweed	Spanishneedle	pepperweed
Carpetweed	Mexicanweed	Spatdock	and other broadleaf
Catchweed bedstraw	Milk vetch	Speedwell	weeds.
Catsear	Milkweed	Spiny Amaranth	
Catnip	bloodflower	Spiny cocklebur	
Chickweed	Mugwort	Spotted catsear	
Chicory	Morningglory	Spotted knapweed	
Cinquefoil	Mousear chickweed	Spotted spurge	
Clover	Musk thistle	Spurge	
Cockle	Mustard	Spurweed	
Cocklebur Coffeebean	Narrowleaf plantain	Stinging nettle	
Coffeeweed	Narrowleaf vetch	Stinkweed	
Common chickweed	Nettle	Stitchwort	
Common mullein Common	Nutgrass	Strawberry clover	
sowthistle Corn Chamomile	Orange hawkweed	Sumac	
Creeping jenny Crimson	Oxalis	Sunflower	
clover Croton	Oxeye daisy	Sweet clover	
Cudweed	Parsley-plant	Tall nettle	
Curly dock	Parsnip	Tall vervain	
Curly indigo Dandelion	Pearlwort	Tansy mustard	
Dead nettle	Pennycress	Tansy ragwort	
Dock	Pennywort	Tanweed	
Dogbane	Peppergrass	Tarweed	
Dogfennel Elderberry	Pepperweed	Thistle	
English daisy	Pigweed	Tick trefoil	
Fall dandelion	Pineywoods	Toadflax	
False dandelion False flax	bedstraw Plains	Trailing crownvetch	
False sunflower	coreopsis Plantain	Tumble mustard	
Fiddleneck	Poison hemlock	Tumble pigweed	
Field bindweed Field pansy	Poison ivy	Tumbleweed	
Flea bane (daisy) Flixweed	Poison oak	Velvet leaf	
Florida betony Florida	Pokeweed	Venice mallow	
pusley Frenchweed	Poorjoe	Veronica	

MIXING INSTRUCTIONS

Fill spray tank with water. Start agitation and slowly add the specified amount of concentrate. Maintain continuous agitation after mixing and during application. If this product is allowed to stand in tank for extended periods of time, some separation may occur. Reagitate before use.

ORNAMENTAL LAWNS AND TURF

This product may be applied by broadcast or spot treatment application methods. Spot treatment is defined as a treatment area no greater than 1,000 ft² per acre.

Rates for Ornamental Lawns and Turf

For Bluegrass, Ryegrass and Fescue, apply 2 to 3 pints per acre. For Creeping Bentgrass use 1.25 pints per acre. For Common and Hybrid Bermudagrass, Bahiagrass, Carpetgrass, Centipedegrass, Zoysiagrass and St. Augustinegrass apply 1.5 to [1.75] [2.0] pints per acre.

Make broadcast applications of 1.25 to 3 pints per acre in 20 to 240 gallons of water per acre (or spot treatment application of 0.46 to 1.10 fl. oz. in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf grasses.

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE: For all grasses (1) avoid double coverage resulting from overlap unless temporary turf injury can be tolerated and (2) use reduced rates if grass is stressed from heat, drought, etc.

SOD FARMS

This product is also intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

Timing

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds.

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Apply this herbicide to grass seedlings after the second mowing. Also, applications to dormant Bermudagrass, dormant Zoysiagrass and dormant Bahiagrass are suggested.

Rates for Sod Farms

For Bluegrass, Ryegrass and Fescue, apply 2 to 3 pints per acre. For Creeping Bentgrass use 1.25 pints per acre. For Common and Hybrid Bermudagrass, Bahiagrass, Carpetgrass, Centipedegrass, Zoysiagrass and St. Augustinegrass apply 1.5 to [1.75] [2.0] pints per acre. Spray volume for Sod Farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

OTHER NON-CROP AREAS

(fencerows, hedgerows, roadsides (including aprons and guard rails), ditches, rights-of-way, utility, pipelines, powerlines, railroads, airports, commercial plants, storage and lumber yards, barrier strips and firebreaks, equipment areas, nurseries and ornamental plantings, fuel tank farms, pumping stations and other industrial sites)

For broadcast application to control broadleaf weeds, mix at a rate of 2 to 3 pints per acre. Spray volume for these applications is 50 to 300 gallons of water per acre. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.

For small (spot) applications with small tank sprayers, apply at the rate of 0.75 to 2.0 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Certain species are more resistant to this herbicide and may display regrowth. Mix thoroughly before spraying.

COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. **[Optional Language:** Some suggested compatibility agents to try are Kalo Laboratories Complex, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Coop.'s Chem-Link.] These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product.)
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 25°F. Protect product from freezing. If allowed to freeze, remix well before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL AND HANDLING:

[**Note to Reviewer:** The following statement will be included on all Final Printed Labels bearing multiple Container Handling statements] **NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size."

[**Note to Reviewer:** The bracketed section headers will be included when multiple container types / sizes are listed on the label.][**Nonrefillable Containers 5 Gallons or Less:**] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[**Nonrefillable Containers Larger than 5 Gallons:**] Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[**Refillable Container Larger than 5 Gallons:**] Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV110514)

TruPower is a Registered Trademark of Nufarm Americas Inc.
All other trademarks are the property of their respective owners.

LABEL HISTORY

File Name	Revision Mark	Comments
000228-00445.20090617.EPA.RED.FIN	RV061709N	
000228-00445.20130709.EPA.ABN.PRN98-10	RV070913	Adding Change Up Selective Herbicide as a ABN
000228-00445.20130717.MASTER (RV071713)	RV071713	EPA approval of ABN
000228-00445.20130925.EPA Notification	RV092513	Do not apply this product to Floratam St. Augustinegrass. CA and NY issues.
000228-00445.20140116.MASTER	RV071713	EPA approval of the above submission.
000228-00445.20140516.Draft	RV051614	Revision to maximum application rate/acre to conform to MCPA RED.
000228-00445.20140709.DRAFT	RV070914	Changes requested by US EPA
000228-00445.20140710.MASTER	RV071014	EPA Approval
000228-00445.20140804.EPA Amendment	RV080414	EPA Amendment per SJ, JF and RF
000228-00445.20141023.EPA Amendment	RV102314	EPA Review
000228-00445.20141105.EPA Amendment	RV110514	EPA Review