

228-449

8/25/2010

10/10



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

228-449

Date of Issuance:

AUG 25 2010

Term of Issuance:

Name of Pesticide Product:

Chaser Ultra<sup>3</sup> Selective
Herbicide

NOTICE OF PESTICIDE:

\_ Registration
\_x Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Nufarm Americas, Inc.
150 Harvester Drive
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:

- 1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following revisions to the final printed labeling:
a. Remove "flaggers" from the list of handlers in the PPE section as this product is prohibited for aerial applications. Additionally, the eyewear text in the handler PPE section must be revised to "goggles or face shield"
b. Per the 2,4-Dp RED, the text, "Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them." must be added to the User Safety Requirements text currently on the label.
c. As this product is a toxicity category I for primary eye irritation, a Note to Physician which addresses technical information on symptomatology, supportive treatments, etc. should be added to the Note to Physician section of the First Aid box.

Continued on Page 2

Signature of Approving Official:

Kathryn V. Montague
Project Manager 23
Herbicide Branch
Registration Division (7505P)

[Handwritten signature of Kathryn V. Montague]

Date:

AUG 25 2010

2 of 10

- d. The Environmental Hazards statement must have the following revisions:
  - 1. Per the REDs, "This pesticide may be toxic to fish, aquatic invertebrates, aquatic plants and may adversely affect other non-target plants." must be the first sentence of the Environmental Hazards statement. Remove, "This may be toxic to fish, aquatic invertebrates and aquatic plants."
  - 2. Per the 2,4-Dp RED, add "Application around a cistern or well may result in contamination of drinking water or groundwater." to the end of the second paragraph.
  - 3. The third paragraph, "Do not discharge...Regional Office of the EPA." is not required for end-use products and must be removed.
  
- e. Move the following text to the spray drift management section:  
"Low humidity and high temperatures increase evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and high temperatures."
  
- f. The typographical error, "definition *foe*..." must be corrected in the spray drift section.
  
- g. The restriction, "For residential turf, golf courses, sod farms and grass grown for seed, do not apply more than 1.5 lbs ae/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days" must be revised per the REDs to read as follows:  
  
"Do not apply more than 1.5 lbs ae **MCPA**/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of **30** days."
  
- 3. A stamped copy of your labeling is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment or the next printing of the label whichever occurs first. Products shipped after 12 months from the date of this Notice must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

# CHASER<sup>®</sup> ULTRA<sup>3</sup>

## SELECTIVE HERBICIDE

A PREMIUM THREE-WAY POST-EMERGENT SELECTIVE BROADLEAF HERBICIDE CONTAINING MCPA, FLUROXYPYR AND DICHLORPROP-p. FOR USE ON GOLF COURSES, PARKS, ORNAMENTAL TURF LAWNS, ALSO FOR USE ON SOD FARMS. CONTROLS DANDELIONS, CHICKWEEDS, PLANTAINS, OXALIS, SPURGE AND MANY OTHER BROADLEAF WEEDS, SOME OF WHICH ARE LISTED ON THIS LABEL.

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	41.60%
1-Methylheptyl Ester of Fluroxypyr:((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid, 1-methylheptyl Ester**	4.90%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***‡	8.10%

<b>OTHER INGREDIENTS:</b>	<b>45.40%</b>
<b>TOTAL:</b>	<b>100.00%</b>

Isomer Specific Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	33.97%	3.20 lbs./gal.
**((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid	3.40%	0.32 lbs./gal.
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	6.79%	0.64 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p.

**DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.**

**KEEP OUT OF REACH OF CHILDREN**

**DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-449  
EPA EST. NO. 228-IL-1

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**

**AUG 25 2010**


Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-449

MANUFACTURED BY  
NUFARM AMERICAS INC.  
BURR RIDGE, IL 60527



NET CONTENTS      GALS.



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, or on clothing. Avoid contact with skin.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**Mixers, loaders, applicators, flaggers, and other handlers must wear:**


- Long-sleeved shirt and long pants,
- Protective eyewear (goggles, face shield, or safety glasses)
- Chemical-resistant gloves (such as or made of any waterproof material), and
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls Statements:** If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 1 70.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	
<ul style="list-style-type: none"> <li>• Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product.</li> <li>• Wash the outside of gloves before removing.</li> <li>• As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>	

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate use of gastric lavage.	



50 of 10

### ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants and may be hazardous to aquatic organisms in water adjacent to treated areas. This may be toxic to fish, aquatic invertebrates and aquatic plants. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Dichlorprop-p and MCPA has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollution Discharge Elimination (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product should be used strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposure.

Under some conditions, this product may have a potential to runoff to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no-till, limited till and contour plowing; these methods also reduce pesticide runoff. Where feasible, use application techniques such as T-banding, and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips is recommended along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where runoff could occur to minimize water runoff.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and high temperatures.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees\* and Roughs). Also for use on Sod Farms.

\*Excluding Bentgrass Tees.

### USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Carpetgrass, Centipede, St. Augustine, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur.

Aerial application of this product is prohibited. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable tree and shrubs, since injury may result.

### SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. groundboom, sprayer) can influence pesticide drift. The applicator must evaluate all factors and make the appropriate adjustments when applying this product.

**Droplet Size:**

Use only medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles.

**Wind Speed:**

Do not apply at wind speeds greater than 10 mph.

**Temperature Inversions:**

If applying at wind speeds less than 3 mph, the applicator must determine if 1) conditions of temperature inversion exist or 2) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

**Additional Requirements for groundboom application:**

Do not apply with a nozzle height greater than four feet above the target site.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

### WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

- |                            |                   |                       |                       |
|----------------------------|-------------------|-----------------------|-----------------------|
| Alder                      | Creeping jenny    | Knotweed              | Rough fleabane        |
| Annual yellow sweet clover | Crimson clover    | Lambsquarter          | Roundleaved marigold  |
| Artichoke                  | Croton            | Lespedeza             | Rush                  |
| Aster                      | Cudweed           | Locoweed              | Russian pigweed       |
| Austrian fieldcress        | Curly dock        | Lupine                | Russian thistle       |
| Bedstraw                   | Curly indigo      | Mallow                | St. Johnswort         |
| Beggartick                 | Dandelion         | Marshelder            | Scarlet pimpernel     |
| Biden                      | Dead nettle       | Matchweed             | Scotch thistle        |
| Bindweed                   | Dock              | Mexicanweed           | Sheep sorrel          |
| Bird vetch                 | Dogbane           | Milk vetch            | Shepherdspurse        |
| Bitter wintercress         | Dogfennel         | Milkweed bloodf lower | Slender plantain      |
| Bitterweed                 | Elderberry        | Mugwort               | Smallflower galinsoga |
| Black-eyed Susan           | English daisy     | Morningglory          | Smartweed             |
| Black medic                | Fall dandelion    | Mouseear chickweed    | Smooth dock           |
| Black mustard              | False dandelion   | Musk thistle          | Smooth pigweed        |
| Black-seed plantain        | False flax        | Mustard               | Sneezeweed            |
| Blessed thistle            | False sunflower   | Narrowleaf plantain   | Southern wild rose    |
| Blue lettuce               | Fiddleneck        | Narrowleaf vetch      | Sowthistle            |
| Blue vervain               | Field bindweed    | Nettle                | Spanishneedle         |
| Box elder                  | Field pansy       | Orange hawkweed       | Spatterdock           |
| Bracted plantain           | Flea bane (daisy) | Oxalis                | Speedwell             |
| Brassbuttons               | Flixweed          | Oxeye daisy           | Spiny amaranth        |
| Bristly oxtongue           | Florida betony    | Parsley-piert         | Spiny cocklebur       |
| Broadleaf dock             | Florida pusley    | Parsnip               | Spotted catsear       |
| Broadleaf plantain         | Frenchweed        | Pearlwort             | Spotted knapweed      |
| Broomweed                  | Galinsoga         | Pennycress            | Spotted spurge        |
| Buckhorn                   | Garlic mustard    | Pennywort             | Spurge                |
| Buckhorn plantain          | Goathead          | Peppergrass           | Spunweed              |
| Bulbous buttercup          | Goatsbeard        | Pepperweed            | Stinging nettle       |
| Bull nettle                | Goldenrod         | Pigweed               | Stinkweed             |
| Bull thistle               | Ground ivy        | Pineywoods bedstraw   | Stitchwort            |
| Burdock                    | Gumweed           | Plains coreopsis      | Strawberry clover     |
| Burning nettle             | Hairy bittercress | Plantain              | Sumac                 |
| Bur ragweed                | Hairy fleabane    | Poison hemlock        | Sunflower             |
| Burweed                    | Hawkweed          | Poison ivy            | Sweet clover          |
| Buttercup                  | Healall           | Poison oak            | Tall nettle           |
| Canada thistle             | Heartleaf drymary | Pokeweed              | Tall vervain          |
| Carolina geranium          | Hedge bindweed    | Poorjoe               | Tansy mustard         |
| Carpeweed                  | Hedge mustard     | Povertyweed           | Tansy ragwort         |
| Catchweed bedstraw         | Hemp              | Prickly lettuce       | Tanweed               |
| Catsear                    | Henbit            | Prickly sida          | Tarweed               |
| Catnip                     | Hoary cress       | Primrose              | Thistle               |
| Chickweed                  | Hoary plantain    | Prostrate knotweed    | Tick trefoil          |
| Chicory                    | Hoary vervain     | Prostrate pigweed     | Toadflax              |
| Cinquefoil                 | Honeysuckle       | Prostrate spurge      | Trailing crownvetch   |
| Clover                     | Hop clover        | Prostrate vervain     | Tumble mustard        |
| Cockle                     | Horsenettle       | Puncture vine         | Tumble pigweed        |
| Cocklebur                  | Horsetail         | Purslane              | Tumbleweed            |
| Coffeebean                 | Indiana mallow    | Ragweed               | Velvet leaf           |
| Coffeeweed                 | Ironweed          | Red clover            | Venice mallow         |
| Common chickweed           | Jewelweed         | Redroot pigweed       | Veronica              |
| Common mullein             | Jimsonweed        | Red sorrel            | Vervain               |
| Common sowthistle          | Kochia            | Redstem filaree       | Vetch                 |
| Corn chamomile             | Knawel            | Rough cinquefoil      | (continued)           |

80A12

Virginia buttonweed  
Virginia creeper  
Virginia pepperweed  
Wavyleaf bullthistle  
Western clematis  
Western salsify  
White clover  
White mustard  
Wild mustard

Wild aster  
Wild buckwheat  
Wild carrot  
Wild four-o'clock  
Wild garlic  
Wild geranium  
Wild lettuce  
Wild marigold  
Wild onion

Wild parsnip  
Wild radish  
Wild rape  
Wild strawberry  
Wild sweet potato  
Wild vetch  
Willow  
Witchweed  
Woolly morningglory

Woodsorrel  
Woolly croton  
Woolly plantain  
Wormseed  
Yarrow  
Yellow rocket  
Yellowflower pepperweed  
and other broadleaf weeds.

**RECOMMENDED MIXING INSTRUCTIONS**

Fill spray tank with water. Start agitation and slowly add the recommended amount of concentrate. Maintain continuous agitation after mixing and during application. If this product is allowed to stand in tank for extended periods of time, some separation may occur. Reagitate before use.

**ORNAMENTAL LAWNS AND TURF**

For residential turf, golf course, sod farm and grass grown for seed uses, do not apply more than 1.5 lb ae/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Ryegrass, Bentgrass (excluding golf course greens and tees), Bahia, Bermudagrass and Zoysia grass.

**Lower Volume Equipment:** Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

**NOTE:** For all grasses (1) avoid double coverage resulting from overlap unless temporary turf injury can be tolerated. and (2) use reduced rates if grass is stressed from heat, drought, etc.

**COMPATIBILITY**

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, HACO Inc.'s Unite, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 1 68D, Amoco Oil's Amoco Spray Mate, Universal Coop.'s Chem-Link and Loveland Industries, Inc.'s E-Z Mix. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.



90912

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 25°F. Protect product from freezing. If allowed to freeze, remix well before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL:

**Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Nonrefillable containers larger than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable containers larger than 5 gallons:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Or

**Refillable Container:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

## WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

## LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF

INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV060909)

Chaser is a Registered Trademark of Loveland Products, Inc.