228.453 Flease read instructions on reverse before completing form.

01/09/2000

Form Approved. OMB No. 2070-0060

ね

•

.

\$epa	Environmenta _{Wesh}	ington, DC 2041	BO		Registra Amendr X Other		OPP Identifier Number	
		Application	n for Pestic	ide - Sect	ion I			
1. Company/Product Numbe	r			Product Mana	ger	3. P	roposed Classification	
228-453		-v	Joan	ne Miller			None Restricted	
4. Company/Product (Name) Battleship3 Selective Her			PM# . 23				· · · · · · · · · · · · · · · · · · ·	
S. Neme and Address of App Nufarm Americas, Inc. 1333 Burr Ridge Parkw Burr Ridge, IL 60527		ode)	(b)(i), to: EPA	my product is Reg. No			n FIFRA Section 3(c)(3) composition and labeling	
			Section -	uct Name _				
			Section -	<u> </u>	· ··· ···			
Amendment - Explain				Final printed labels in repsonse Agency letter dated			JAN - 9 2006	
Resubmission in resp	onse to Agency letter	dated	L_	"Me Too" A	pplication.		JAN - 9 2005	
X Notification - Explain	below.			Other - Expla	sin belaw.		4 2000	
Label notification consisten 98-10 and EPA regulations product. I understand that i notification is not consisten enforcement action and per	at 40 CFR 152.46, and it is a violation of 18 L t withe terms of PR N	nd no other cha J.S.C. Sec. 100 otice 98-10 and	nges have been 1 to willfully mak 140 CFR 152.46	made to the la te false statem	ibeling or the conf ents to EPA. I fur	idential s ther und	statement of formula of this erstand that if this	
			Section -					
1. Material This Product Will	Be Packaged In:							
Child-Resistant Packaging	Unit Packaging		Water Soluble	Packaging	2. Type of (Container		
Yes	Yes		Yes			Metal Plastic		
No	No		N₀			Glass		
 Certification must be submitted 	lf "Yes" Unit Packaging wgt.	No, per containar	If "Yes" Package wgt	No. per container		Paper Other (S	Specity)	
3. Location of Net Contents I	nformation	4. Size(s) Retai	l Container		5. Location of Leb	l Directio	ons	
Label	onteiner					i I		
6. Manner in Which Label is a	Affixed to Product	L Lithogra Paper pl Stancile	ph uedi d	Other				
			Section - I	V				
1. Contact Point <i>(Complete)</i>	items directly below fo	or identification	of individual to L	e contacted, if	necessary, to pro	cess this	application.)	
Name George Meindl george	e.meindl@us.nuf	1	Registration	Manager		1 '	e No. (Include Area Code) 55.7277	
I certify that the staten I acknowledge that any both under applicable I	knowlingly false or i	Certification the form and all	ON Il attachments th	ereto are true,		plete.	6. Date Application Received (Stamped)	
\mathcal{M} \mathcal{M} \mathcal{M}			Titte. Registrat	• gistration Manager				
4. Typed Name	ł	5.	Date					
George Mehndl			12/14/2	2005				

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

.

White - EPA File Copy {original}

Yallow - Applicant Copy

BATTLESHIP³ SELECTIVE HERBICIDE

FOR THE CONTROL OF ANNUAL AND PERENNIAL BROADLEAF WEEDS IN ORNAMENTAL TURF INCLUDING SOD FARMS

CONTAINS MCPA, FLUROXYPYR, AND TRICLOPYR

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	
1-Methylheptyl Ester of Fluroxypyr:((4-amino-3-5-dichloro-6-fluoro- 2-pyridinyl)oxy)acetic Acid, 1-meth	viheptyl ester** 4.45%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid***	4.07%
OTHER INGREDIENTS:	53.64%

By Isomer Specific AOAC Method, Equivalent to.	
*2-Methyl-4-Chlorophenoxyacetic Acid	30.90%, 2.85 lbs./gal.
**((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid	3.09%, 0.29 lbs./gal.
***3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	2.92%, 0.27 lbs./gal.

Intended For Use By Professional Turf Maintenance Personnel, Landscaping or Commercial Applicators Only.

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si ústed no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

EPA REG, NO.228-453

MANUFACTURED BY NUFARM AMERICAS INC. BURR RIDGE, IL 60527-0866



EPA EST. NO. 228-IL-1

NET CONTENTS GALS. ۰.

000228-00453.20051214 patent

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

NON-WPS TURF USE: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170)-in general only agricultural plant uses are covered by the WPS-must wear eye protection (goggles, face shield, or safety glasses) when using this product. It is recommended that safety glasses include front, brow, and temple protection.

WPS SOD FARM USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170)-in general, agricultural plant uses are covered-must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands, face and arms with soap and water before eating, smoking, drinking, or using the toilet,
- · Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- · Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing,

<u> </u>	FIRST AID
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
	HOT LINE NUMBER ntainer or label with you when calling a poison control center or doctor, or going for treatment. t 1-877-325-1840 for emergency medical treatment information
	NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate use of gastric lavage.

22 /IRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2.4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected hand'ers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

2

This product should be used strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposure.

Under some conditions, this product may have a potential to runoff to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no-till, timited till and contour plowing; these methods also reduce pesticide runoff. Where feasible, use application techniques such as T-banding, and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetaion filter strips is recommended along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where runoff could occur to minimize water runoff.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and for high temperatures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application Do not enter treatment areas until spray has dried.

This product is a broad-spectrum herbicide for control of actively growing annual and perennial broadleaf weeds. It is for use on established Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields, Roadsides and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas; also for use on sod farms. Do not use on Golf Greens or Tees. This product may be applied to the following established turfgrass species:

ESTABLISHED COOL SEASON TURFGRASSES		ESTABLISHED WARM SEASON TURFGRASSEST		
Common Name Scientific Name		Common Name	Scientific Name	
Bentgrass	Agrostis species	Bahiagrass	Paspalum notatum	
Bluegrass, Kentucky	Poa pratensis	Bermudagrass	Cynodon dactylon	
Fescue, chewing	Festuca rubravar, commutata	Buffalograss	Buchloe dactyloides	
Fescue, creeping red	Festuca rubra	Centipedegrass	Eremochloa ophiuroides	
Fescue, sheeps	Festuca ovina	Zoysiagrass	Zoysia japonica	
Fescue, tall	Festuca arundinaceae	Zoysiagrass	Zoysia tenuifolia	
Ryegrass, perennial	Lolium perenne	Fescue, tall (growing in warm season areas)	Festuca arundinacea	

†Do not treat warm season turfgrass with this product when moving height is less than 1/2 inch-

Note: This product may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply this product only to nealthy, well-established turf that has a well-anchored root system. Do not apply this product to grass species not fisted on this label unless potential turf injury can be tolerated.

USE PRECAUTIONS

Do not apply this product directly to, or otherwise permit drift or spray mist to come into contact with cotton, citrus, grapes, tobacco, vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not pour spray solutions near desirable plants. Avoid tine mists. Use lawn type spraver with coarse spray as wind drift is less likely. Spray drift can be reduced by spraying when wind velocity is low, keeping spray boom as low as possible using nozzles that deliver larger spray dioplets and minimize the production of fine spray particles, and by using higher spray volumes and lower pressures. Applications should be made only when hazards from spray drift are at a minimum. Very small quantities of spray which may not be visible, may seriously injure succeptible arits. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Do not apply the oxoosed roots of shallow rooted trees and shrubs. Maximum kill of weeds will be of think of the arity seeded turf at least twice

before treatment with this product.

For optimum results: (1) Add a spray adjuvant, such as DYNE-AMIC for best results; (2) Use a water conditioning agent, such as. QUEST for high PH and water hardness; (3) Avoid applying during excessively dry or hot periods unless irrigation is used: (4) Turt should not be mowed 1 to 2 days before and following application; (5) Reseed no sooner than 3 to 4 weeks after application of this product.

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder Annual yellow sweet clover Artichoke Aster Austrian fieldcress **Bedstraw** Beggartick Biden Bindweed Bird vetch Bitterweed Bitter wintercress Black-eyed Susan Black medic Black mustard Black-seed plantain Blessed thistle Blue lettuce Blue vervain Box elder Bracted plantain Brassbuttons Bristly oxtongue Broadleaf dock Broadleaf plantain Broomweed Buckhorn Buckhorn plantain Bulbous buttercup Bull nettle Bull thistle Burdock Burning nettle Bur ragweed Burweed Buttercup Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catsear Catnip Chickweed Chicory Cinquefoil Clover Cockle Cocklebur Coffeebean Coffeeweed Common chickweed Common mullein Common sowthistle Corn Chamomile

Creeping jenny Crimson clover Croton Cudweed Curly dock Curly indigo Dandelion Dead nettle Dock Dogbane Dogfennei Elderberry English daisy Fall dandelion False dandelion False flax False sunflower Fiddleneck Field bindweed Field pansy Flea bane (daisy) Flixweed Florida betony Florida pusley Frenchweed Galinsoga Garlic mustard Goathead Goatsbeard Goldenrod Ground avv Gumweed Hairy bittercress Hairy fleabane Hawkweed Heatall Heartleaf drymary medige bindweed Hedge mustard Hemp Henbit Heary cress Hoary plantain Houry vervain Honevsuckle Hop theyer Horschettle Horsetad Indiana mallow Ironweed Jewelweed Jimsonweed Kochia Knawel

Knotweed Lambsquarter Lespedeza Locoweed Lupine Mallow Marshelder Matchweed Mexicanweed Milk vetch Milkweed bloodflower Mugwort Morningglory Mousear chickweed Musk thistle Mustard Narrowleaf plantain Narrowleaf vetch Nettle Orange hawkweed Oxalis Oxeye daisy Parsley-piert Parsnip Pearlwort Pennycress Pennywort Peppergrass Pepperweed Plaweed Pineywoods bedstraw Plains coreopsis Plantain Poison hemlock Poison ivv Poison oak Pokeweed Poorjoe Povertyweed Prickly lettuce Prickly sida Primrose Prostrate knotweed Prostrate pigweed Prostrate spurge Prostrate vervain Puncture vine Purslane Ragweed Red clover Redroot pigweed Red sorrel Redstem filaree Rough cinquefoil

4

Rough fleabane Roundleafed marigold Rush Russian pigweed Russian thistle St. Johnswort Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smartweed Smooth dock Smooth pigweed Sneezeweed Southern wild rose Sowthistle Spanishneedle Spatterdock Speedwell Spiny Amaranth Spiny cocklebur Spotted catsear Spotted knapweed Spotted spurge Spurge Spurweed Stinging nettle Stinkweed Stitchwort Strawberry clover Sumab Sunflower Sweet clover Tall nettle Tall vervain Tansy mustard Tansy ragwort Tanweed Tarweed Thistle Tick trefoil Toadflax Trailing Crownvetch Tumble mustard Tumble pigweed Tumbleweed Velvet leaf Venice mallow Veronida Vervain Vetch (continuad) Virginia buttonweed Virginia creeper Virginia pepperweed Wavyleaf bullthistle Western ctematis Western salsify White clover White mustard Wild mustard Wild aster Wild buckwheat Wild carrot Wild four-o'-clock Wild garlic Wild geranium Wild lettuce Wild marigold Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild strawberry Wild vetch Willow Witchweed Wooly morningglory Woodsorrel Wooly croton Wooly plantain Wormseed Yafrow Yellow rocket Yellowriower pepperweed and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks,

RECOMMENDED MIXING INSTRUCTIONS

Fill spray tank with water. Start agitation and slowly add the recommended amount of concentrate. Maintain continuous agitation after mixing and during application. If this product is allowed to stand in tank for extended periods of time, some separation may occur. Reagitate before use.

TURF APPLICATION DIRECTIONS

Apply 3 to 4 pints of this product per acre (1.1 to 1.5 ozs./1,000 sq.tt.) to control above listed broadleat weeds.

The maximum use rate for most warm season grasses is 3 pints per acre.

Make application using equipment that will ensure uniform coverage (see specific application directions below). Foliar sprays should be applied during warm weather when weeds are actively growing. Application under drought conditions may provide less than desirable results. When hard-to-control species are prevalent, when applications are made in late summer, when weeds are mature, or during drought conditions, use the higher rates.

On closely mowed golf course fairway bentgrass, apply this product at a maximum rate of 2 pints in 20 to 240 gallons of water per acre (0.73 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet). During conditions which promote turt stress use lower rates, 1 to 1.5 pints in 20 to 240 gallons of water per acre (0.36 to 0.55 fluid ounces in 0.3 to 5.5 gallons per 1,000 square feet). Slight turt yellowing will disappear after one week. To minimize grass injury, a second application should not be made for at least 4 weeks. Avoid swath overlaps. Do not use on golf course greens or tees.

Note: Broadleaf weeds germinate at different times of the year. Only emerged weeds present at the time of application will be controlled by this product.

Standard Volume Broadcast Applications: Apply 3 to 4 pints of this product in enough water to deliver 20 to 200 gallons of total spray mix per acre (0.5 to 5 gallons spray per 1.000 square feet) Higher application volumes may be used when this product is tank mixed with liquid fertilizers.

Low Volume Broadcast Applications: Apply 3 to 4 pints of this product in enough water to deliver from 5 to 40 gallons of total spray mixture per acre. Use low pressures and application equipment capable of delivering a uniform droplet size that can wet the weed leaf surface.

Targeted Zone Applications or Spot Treatment Using Portable Sprayers: Mix 0.5 to 1.0 ounce of this product in enough water to make one gallon of spray and apply at any time broadleaf weeds are actively growing. Target plants should be sprayed to wet but not beyond point of runoff

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always store pesticides in a secured warehouse or storage building. Store at temperatures above 25°F. Protect from freezing. If allowed to freeze, remix well before using. Freezing does not after the product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two watlets high. Do not store near open containers of fertilizer seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillate. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. 'mproper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL FOR NON-REFILLABLE CONTAINERS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration or it allowed by State and local authorities, by burring. If burned, stay out of smoke,

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings which have been opened during use and replace all caps. Contact Nufarm Customer Service at 1 800-345-3330 to arrange for return of the empty refillable container.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Desirable plant injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Nufarm or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Nufarm and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. NUFARM MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to for beyond the control of) Seller or Nufarm, and Buyer assumes the risk of any such use.

To the extent allowed by law. Nufarm or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF NUFARM AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF NUFARM OR SELLER. THE REPLACEMENT OF PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

(RV083005)

Battleship is a Registered Trademark of Helena Chemical Company

U.S. Patent No. 6.831.038 (VBS NUBS chemical technology)



Nufarm Americas, Inc. George Meindl Registration Manager 1333 Burr Ridge Parkway, Suite 125. Burr Ridge, IL 60527 Phone: 708.755-7277 Fax: 708.754.0314 george.meindl@us.nufarm.com

December 14, 2005

Via Overnight Courier

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) U. S. Environmental Protection Agency Room 266A, Crystal Mall 2 1801 Bell Street Arlington, VA 22202-4501

Subject: Battleship®³ Selective Herbicide EPA Reg. No. 228-453 Label notification per PR98-10

Dear Ms. Hobgood:

Nufarm America, Inc would like to make a label change to the subject registration. We feel that this label change is permitted and consistent with PRN98-10 section II (Labeling notifications) and part N (Other revisions).

Specifically we have added a U.S patent number, which appears on the last page (page 6) of the label.

To process this request please find enclosed the following:

- Application for Pesticide Registration (EPA form 8570-1)
- Revised labeling with areas of change clearly identified († copy)

If you should have any questions regarding this matter, please feel free to contact me at 708.755.7277 or email at <u>george.meindl@us.nufarm.com</u>.

Sincerely,

George Meind Registration Manager Nufarm Americas, Inc.

H:\228-453\abel notification adding patent number per PRN98-10 submitted 12.14.2005.doc