

 <p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	<p>EPA Reg. Number: 228-455</p>	<p>Date of Issuance: SEP 28 2005</p>
	<p>Terms of Issuance: Conditional</p>	
	<p>Name of Pesticide Product: Strike 3 Ultra 3 Selective Herbicide</p>	
<p>Name and Address of Registrant (include ZIP Code): Nufarm Americas, Inc. 1333 Burr Ridge Parkway Suite 125A Burr Ridge, IL 60527</p>		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is registered in accordance with FIFRA sec.3(c)(7)(A) provided that you.</p> <ol style="list-style-type: none"> Add a "Note to Physician" statement to the label. It should read: Note To Physician: Probable mucosal damage may contraindicate the use of gastric lavage. <p>(Comments continued on the next page for this Notice of Registration)</p> <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions</p>		
<p>Signature of Approving Official: <i>Joanne J. Miller</i></p>	<p>Date: SEP 28 2005</p>	

2. A. Add your EPA establishment number and net contents to the label.
- B. Revise the EPA Registration Number to read, "EPA Reg. No. 228-455."
3. Submit storage stability (830.6317) and corrosion characteristics (830.6320) studies within one year of the date of this registration notice.
4. Submit and/or cite all data required for the registration/reregistration of this product when the Agency requires all registrants of similar products to submit data.
5. Submit one (1) copy of the final printed labeling before you release this product for shipment.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

3/10

STRIKE 3 ULTRA[®] 3 SELECTIVE HERBICIDE

A Three-Way Post-Emergent Selective Broadleaf Herbicide Containing 2,4-D, Fluroxypyr And Dichlorprop-p.
For Use On Golf Courses, Parks, Highways, Ornamental Turf Lawns And Similar Non-Crop Areas.
Also For Use On Sod Farms. Kills Dandelions, Chickweeds, Plantains, Oxalis, Spurge And Many Other
Broadleaf Weeds, Some Of Which Are Listed On This Label.

GET THE OPTICAL ADVANTAGE*

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	39.30%
1-Methylheptyl Ester of Fluroxypyr: [(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl) oxy] acetic Acid, 1-methylheptyl ester**	5.88%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy) propionic Acid***†	9.72%
OTHER INGREDIENTS:	<u>45.10%</u>
TOTAL:	100.00%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	32.64%, 3.2 lbs/gal.
**[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl) oxy] acetic acid	4.08%, 0.4 lbs/gal.
***(+)-R-2-(2,4-Dichlorophenoxy) propionic Acid	8.16%, 0.8 lbs/gal.

†CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p.

**INTENDED FOR USE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL,
LANDSCAPING OR COMMERCIAL APPLICATORS ONLY.
NOT INTENDED FOR SALE TO OR USE BY HOMEOWNERS**

**KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

NOTE: Spanish language is optional

EPA REG. NO. 228-
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

000228-00000.20050606

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
SEP 28 2005**

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-455

4/10

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO**

Corrosive, causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

NON-WPSTURF USES: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) in general only agricultural plant uses are covered by the WPS, must wear face shield, goggles, or safety glasses and long pants, long-sleeved shirt, socks, shoes and chemical-resistant gloves when mixing, loading or applying this product. It is recommended that safety glasses include front, brow, and temple protection. In addition to the clothing and eye protection listed above, commercial mixer/loaders must wear chemical-resistant in place of rubber gloves except when the product is to be applied to a golf course. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

NON-WPS INDUSTRIAL USES: When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield, goggles, or safety glasses and chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow, and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

WPS USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) in general, agricultural plant uses are covered, must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear, when cleaning equipment, mixing or loading. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned. If this container is over one gallon and less than five gallons, mixers and loaders who do not use a mechanical system (such as a probe and pump or spigot) to transfer contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Engineering Controls Statements: If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates and fish. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and 2,4-DP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling this product at mixing, loading, and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product is for use on ornamental turf lawns (residential, industrial and institutional), parks, cemeteries, athletic fields and golf courses (fairways, aprons, tees* and roughs) and similar turf areas.

*Excluding Bentgrass Tees.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Centipede, St. Augustine, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

The suitable use on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

This pesticide should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposure.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Where feasible, use application techniques such as T-banding and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur will minimize water run-off.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

WEEDS CONTROLLED

Alder	Canada thistle	Field bindweed	Knotweed
Annual yellow sweet clover	Carolina geranium	Field pansy	Lambsquarter
Artichoke	Carpetweed	Flea bane (daisy)	Lespedeza
Aster	Catchweed bedstraw	Flixweed	Locoweed
Austrian fieldcress	Catsear	Florida betony	Lupine
Bedstraw	Catnip	Florida pusley	Mallow
Beggartick	Chickweed	Frenchweed	Marshelder
Biden	Chicory	Galinsoga	Matchweed
Bindweed	Cinquefoil	Garlic mustard	Mexicanweed
Bird vetch	Clover	Goathead	Milk vetch
Bitterweed	Cockle	Goatsbeard	Milkweed bloodflower
Bitter wintercress	Cocklebur	Goldenrod	Mugwort
Black-eyed Susan	Coffeebean	Ground ivy	Morningglory
Black medic	Coffeeweed	Gumweed	Mousear chickweed
Black mustard	Common chickweed	Hairy bittercress	Musk thistle
Black-seed plantain	Common mullein	Hairy fleabane	Mustard
Blessed thistle	Common sowthistle	Hawkweed	Narrowleaf plantain
Blue lettuce	Corn Chamomile	Healall	Narrowleaf vetch
Blue vervain	Creeping jenny	Heartleaf drymary	Nettle
Box elder	Crimson clover	Hedge bindweed	Orange hawkweed
Bracted plantain	Croton	Hedge mustard	Oxalis
Brassbuttons	Cudweed	Hemp	Oxeye daisy
Bristly oxtongue	Curly dock	Henbit	Parsley-piert
Broadleaf dock	Curly indigo	Hoary cress	Parsnip
Broadleaf plantain	Dandelion	Hoary plantain	Pearlwort
Broomweed	Dead nettle	Hoary vervain	Pennycress
Buckhorn	Dock	Honeysuckle	Pennywort
Buckhorn plantain	Dogbane	Hop clover	Peppergrass
Bulbous buttercup	Dogfennel	Horsenettle	Pepperweed
Bull nettle	Elderberry	Horsetail	Pigweed
Bull thistle	English daisy	Indiana mallow	Pineywoods bedstraw
Burdock	Fall dandelion	Ironweed	Plains coreopsis
Burning nettle	False dandelion	Jewelweed	Plantain
Bur ragweed	False flax	Jimsonweed	Poison hemlock
Burweed	False sunflower	Kochia	Poison ivy
Buttercup	Fiddleneck	Knave	Poison oak

Pokeweed	Shepherdspurse	Tall vervain	Wild buckwheat
Poorjoe	Slender plantain	Tansy mustard	Wild carrot
Povertyweed	Smallflower galinsoga	Tansy ragwort	Wild four-o'clock
Prickly lettuce	Smartweed	Tanweed	Wild garlic
Prickly sida	Smooth dock	Tarweed	Wild geranium
Primrose	Smooth pigweed	Thistle	Wild lettuce
Prostrate knotweed	Sneezeweed	Tick trefoil	Wild marigold
Prostrate pigweed	Southern wild rose	Toadflax	Wild onion
Prostrate spurge	Sowthistle	Trailing Crownvetch	Wild parsnip
Prostrate vervain	Spanishneedle	Tumble mustard	Wild radish
Puncture vine	Spatterdock	Tumble pigweed	Wild rape
Purslane	Speedwell	Tumbleweed	Wild strawberry
Ragweed	Spiny Amaranth	Velvet leaf	Wild sweet potato
Red clover	Spiny cocklebur	Venice mallow	Wild vetch
Redroot pigweed	Spotted catsear	Veronica	Willow
Red sorrel	Spotted knapweed	Vervain	Witchweed
Redstem filaree	Spotted spurge	Vetch	Woolly morningglory
Rough cinquefoil	Spurge	Virginia buttonweed	Woodsorrel
Rough fleabane	Spurweed	Virginia creeper	Wooly croton
Roundleafed marigold	Stinging nettle	Virginia pepperweed	Wooly plantain
Rush	Stinkweed	Wavyleaf bullthistle	Wormseed
Russian pigweed	Stitchwort	Western clematis	Yarrow
Russian thistle	Strawberry clover	Western salsify	Yellow rocket
St. Johnswort	Sumac	White clover	Yellowflower pepperweed and other broadleaf weeds.
Scarlet pimpernel	Sunflower	White mustard	
Scotch thistle	Sweet clover	Wild mustard	
Sheep sorrel	Tail nettle	Wild aster	

Some of these species may require repeat spot applications and/or use of higher rate recommended on this product label even under ideal conditions for application.

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to bluegrass, fescue, ryegrass, bentgrass (excluding golf course greens and tees), bahiagrass, bermudagrass and zoysia grass.

Notes: On closely mowed golf course fairway bentgrass, apply this product at a maximum rate of 2 pints in 20 to 240 gallons of water per acre (0.73 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet). During conditions which promote turf stress use lower rates, 1 to 1-1/2 pints in 20 to 240 gallons of water per acre (0.36 to 0.55 fluid ounces in 0.3 to 5.5 gallons per 1,000 square feet). Slight turf yellowing will disappear after one week. To minimize grass injury, a second application should not be made for at least 4 weeks. Avoid swath overlaps. Do not use on golf course bentgrass greens or tees. On warm season grasses, exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur. If bermudagrass is dormant, up to 4 pints per acre may be used. However, some hybrid bermudagrasses may be sensitive to this product. Contact your local extension service weed control specialist.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use this product at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE: For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

SOD FARMS

This product is intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

Timing

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to Floratam St. Augustinegrass.

The application of this herbicide to grass seedlings is not recommended until after the second mowing.

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant bermudagrass, dormant zoysia grass and dormant bahiagrass are suggested.

Recommended Rates for Sod Farms

For bluegrass, ryegrass and fescue, apply 2 to 3 pints per acre. For creeping bentgrass use 1.25 pints per acre. For common and

8/10

hybrid bermudagrass, bahiagrass, zoysia grass and St. Augustine grass apply 1.5 to 1.8 pints per acre. Spray volume for sod farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages. The maximum application is 5 pints per acre per season.

NON-TURF AREAS

Roadsides (including aprons and guardrails), rights-of-way, and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 2 to 6 pints of this product per 50 to 300 gallons of water. This mixture will cover 1 acre. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 2-1/4 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 4 to 8 pints of this product to 100 gallons of water applying 200 to 600 gallons of spray mixture per 43,500 square feet depending upon the height and thickness of the brush. Mix thoroughly before spraying.

COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Protect from freezing. Store at temperatures above 25°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings and replace all caps. Contact Nufarm Customer Service at 1-800-345-3330 to arrange for return of the empty refillable container.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Nufarm or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Nufarm and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. NUFARM MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or Nufarm, and buyer assumes the risk of any such use.

To the extent allowed by law, Nufarm or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF NUFARM AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF NUFARM OR SELLER, THE REPLACEMENT OF PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

(NEW)

The Optical Advantage are Registered Trademarks of Nufarm Americas Inc.
Strike 3 Ultra is a Registered Trademark of Agrilience, LLC

10/10

**NOT TO BE PART OF THE PRINTED LABEL
FOR NUFARM INTERNAL REGULATORY USE ONLY**

06/06/05

New submission to EPA.

06-06-05

8

06-06-05