



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number: 228-489

Date of Issuance: AUG 8 2006

NOTICE OF PESTICIDE:

[x] Registration
[] Reregistration
(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product: Triplet Low Odor Premium Lawn Weed Killer Concentrate

Name and Address of Registrant (include ZIP Code):

Nufarm Americas, Inc.
1333 Burr Ridge Parkway Suite 125A
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label change:

a. Revise the EPA Registration Number to read: "EPA Reg. No. 228-489."

b. For the "commercial label" add the following Pesticide Disposal statement to the Storage and Disposal section:

"Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility."

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Handwritten signature: Joanne I. Miller

Date:

AUG 8 2006

2/14

(RESIDENTIAL LABEL)

TRIPLET[®] LOW ODOR PREMIUM LAWN WEED KILLER CONCENTRATE

FOR SELECTIVE BROADLEAF WEED CONTROL IN RESIDENTIAL TURF
WHEN USING A HOSE-END SPRAYER, JUST CONNECT HOSE, TURN ON WATER AND SPRAY.
ONE QUART COVERS UP TO 10,000 SQUARE FEET
(SEE DIRECTIONS FOR DIFFERENT GRASSES.)
CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

**ACCEPTED
with COMMENTS**
In EPA Letter Dated:
AUG 8 2006
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-489

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	13.24%
Dimethylamine Salt of Mecoprop-p,** ‡	2.29%
Dicamba Acid***	0.64%

OTHER INGREDIENTS:	83.83%
TOTAL:	100.00%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	7.10%	0.618 lbs./gal.
**Mecoprop-p Acid	1.89%	0.164 lbs./gal.
***Dicamba Acid	0.64%	0.056 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS GALS.

3/14

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pet on treatment area during application. Do not enter treatment areas until spray has dried.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. If spills occur, collect the material and dispose of by following disposal instructions on this label. Do not pour product or rinse water down any drain.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. Shake well before using.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. If necessary, contact your local Extension Agent.

4/14

WEEDS CONTROLLED

Alder	Dogbane	Narrowleaf vetch	Spotted spurge
Annual yellow sweet clover	Dogfennel	Nettle	Spurge
Artichoke	Elderberry	Nutgrass	Spurweed
Aster	English daisy	Orange hawkweed	Stinging nettle
Austrian fieldcress	Fall dandelion	Oxalis	Stinkweed
Bedstraw	False dandelion	Oxeye daisy	Stichwort
Beggartick	False flax	Parsley-piert	Strawberry clover
Biden	False sunflower	Parsnip	Sumac
Bindweed	Fiddleneck	Pearlwort	Sunflower
Bird vetch	Field bindweed	Pennycress	Sweet clover
Bitterweed	Field pansy	Pennywort	Tall nettle
Bitter wintercress	Flea bane (Daisy)	Peppergrass	Tall vervain
Black-eyed Susan	Flixweed	Pepperweed	Tansy mustard
Black medic	Florida betony	Pigweed	Tansy ragwort
Black mustard	Florida pusley	Pineywoods bedstraw	Tanweed
Blackseed plantain	Frenchweed	Plains coeopsis	Tarweed
Blessed thistle	Galinsoga	Plantain	Thistle
Blue lettuce	Garlic mustard	Poison hemlock	Tick trefoil
Blue vervain	Goathead	Poison ivy	Toadflax
Box elder	Goatsbeard	Poison oak	Trailing crown vetch
Bracted plantain	Goldenrod	Pokeweed	Tumble mustard
Brassbuttons	Ground ivy	Poorjoe	Tumble pigweed
Bristly oxtongue	Gumweed	Povertyweed	Tumbleweed
Broadleaf dock	Hairy bittercress	Prickly lettuce	Velvet leaf
Broadleaf plantain	Hairy fleabane	Prickly sida	Venice mallow
Broomweed	Hawkweed	Primrose	Veronica
Buckhorn	Healall	Prostrate knotweed	Vervain
Buckhorn plantain	Heartleaf drymary	Prostrate pigweed	Vetch
Bulbous buttercup	Hedge bindweed	Prostrate spurge	Virginia buttonweed
Bull nettle	Hedge mustard	Prostrate vervain	Virginia creeper
Bull thistle	Hemp	Puncture vine	Virginia pepperweed
Burdock	Henbit	Purslane	Wavyleaf bullthistle
Burning nettle	Hoary cress	Ragweed	Western clematis
Bur ragweed	Hoary plantain	Red clover	Western saisify
Burweed	Hoary vervain	Redroot pigweed	White clover
Buttercup	Honeysuckle	Red sorrel	White mustard
Canada thistle	Hop clover	Redstem filaree	Wild mustard
Carolina geranium	Horsenettle	Rough cinquefoil	Wild aster
Carpetweed	Horsetail	Rough fleabane	Wild buckwheat
Catchweed bedstraw	Indiana mallow	Roundleafed marigold	Wild carrot
Catsear	Ironweed	Rush	Wild four-o'clock,
Catnip	Jewelweed	Russian pigweed	Wild geranium
Chickweed	Jimsonweed	Russian thistle	Wild lettuce
Chicory	Kochia	St. Johnswort	Wild marigold
Cinquefoil	Knawel	Scarlet pimpernel	Wild parsnip
Clover	Knotweed	Scotch thistle	Wild radish
Cockle	Lambsquarter	Sheep sorrel	Wild rape
Cocklebur	Lespedeza	Shepherdspurse	Wild strawberry
Coffeebean	Locoweed	Slender plantain	Wild sweet potato
Coffeeweed	Lupine	Smallflower galinsoga	Wild vetch
Common chickweed	Mallow	Smartweed	Willow
Common mullein	Marshelder	Smooth dock	Witchweed
Common sowthistle	Matchweed	Smooth pigweed	Woolly morningglory
Corn chamomile	Mexicanweed	Sneezeweed southern wild rose	Woodsorrel
Creeping Jenny	Milk vetch	Sowthistle	Woolly croton
Crimson clover, croton	Milkweed bloodflower	Spanishneedle	Woolly plantain
Cudweed	Mugwort	Spatardock	Wormseed
Curly dock	Morningglory	Speedwell	Yarrow
Curly indigo	Mouseear chickweed	Spiny amaranth	Yellow Rocket
Dandelion	Musk thistle	Spiny cocklebur	Yellow flower pepperweed
Dead Nettle	Mustard	Spotted catsear	and other broadleaf weeds
Dock	Narrowleaf plantain	Spotted knapweed	

5/14

USE TIPS

1. Spray during the growing season when weeds are growing actively.
2. For best results, do not mow for 2 days before or after treating.
3. Do not treat when air temperatures exceed 90°F or damage to grass may occur.
4. For best results, do not water lawn within 24 hours after treatment. Poor weed control may result if rainfall occurs within 24 hours of application.
5. Shake well before using.
6. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
7. Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees. Clover will be damaged if sprayed.
8. Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.
9. If certain tough weeds are not dead within 3 weeks, repeat treatment. Broadcast application is not necessary; follow spot treatment directions.

HOW MUCH TO USE

SPOT TREATMENT

Pressure (Trigger) Sprayer: Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

ENTIRE LAWN

Hose-End Sprayer: Use a hose-end sprayer designed for lawns. Pour recommended amount of product (Column 2) in bottle and add water to indicated level (Column 3), and spray on area shown in Column 4. For example, to spray 2,500 square feet measure 20 fluid ounces (40 Tablespoons) in jar, add water to the 10 gallon level and mix. Turn on garden hose and spray as evenly as possible over 2,500 square feet.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

NOTE: Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

6/14

Pressure Sprayer: Measure indicated amount of this product, pour into tank, and dilute with 1 gallon of water according to the following chart. Use a spray pattern with coarse spray droplets without any fine mist.

AMOUNT OF CONCENTRATE IN 1 GALLON OF WATER REQUIRED TO TREAT 250 SQ. FT. OF LAWN	
GRASS TYPE	AMOUNT OF CONCENTRATE
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)
Warm Season Grasses Bahagrass Common Bermudagrass Zoysiagrass	1 Fluid Ounce (2 Tablespoon)
Warm Season Grasses Centipedegrass St. Augustinegrass*	1/2 Fluid Ounce (1 Tablespoon)

*Do not apply to Fioratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep from freezing. to be stored in original container and placed in areas inaccessible to children.

PESTICIDE DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

Seller warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(RV)

Triplet and The Optical Advantage are Registered Trademarks of Nufarm Americas, Inc.

[OPTIONAL BOX]

PRODUCT FACTS

What it Does:

Kills all Major Broadleaf Weeds including Chickweed, Clover and Dandelion (See inside for complete list)

Where to Use: (Icon weeds)

Lawns: Kills the Weeds not the Lawn

When to Use:

Do not apply when daytime temperatures are above 90°F as lawn injury can occur.

Questions and Comments: (Icon phone) (icon computer)

Call xxxxxxxx or visit our website at www.xxxxxxxxxx

7/14

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING HYPE - FRONT PANEL

Patented Low Odor Formulation
 Low Odor Formulation
 Now With Patented Low Odor Formulation
 Kills Over 250 Weed Types
 Kills 250+ Weed Types
 Kills Over 250 Weed Listed
 Kills Over 250 Weeds
 Kills All Major Broadleaf Weeds
 Kills Dandelion, Clover and Other Broadleaf Weeds
 Kills Dandelion, Clover and Other Listed Broadleaf Weeds
 Controls Over 250 Weeds
 Use When Weeds Are Actively Growing
 May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
 Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
 Overnight Results
 Works in 24 Hours
 Kills the Weeds, Not the Lawn
 Rainproof in Hours - Won't Wash Away
 Colorless Formulation
 Won't Harm the Home Lawn
 Illustration(s) of various weeds listed on label
 Contains Triplet Low Odor Formula
 Easy to Use
 For Thick, Weed-Free Turf
 Three Way Control
 Apply to a Moist Lawn
 Contains The Three Way Power of Triple Low Odor
 One Quart Covers Up To 10,000 Square Feet (One XXX covers up to XX,XXX square feet.)
 Easy to Mix
 Just Add Water
 New Formulation
 Specially formulated for residential use.

ALTERNATE HOSE-END SPRAYER INSTRUCTIONS

How To Use:

(Illustrations)

1. Attach garden hose to spray nozzle.
2. Turn on water supply.
3. Bend safety tab down and turn control to "water"
4. Point nozzle toward lawn and turn control to "on". Product will automatically mix with water. Walk back and forth at a steady pace.

8/14

(COMMERCIAL LABEL)

TRIPLET[®] LOW ODOR PREMIUM LAWN WEED KILLER CONCENTRATE

SELECTIVE BROADLEAF WEED CONTROL FOR ORNAMENTAL TURFGRASSES.
TO CONTROL DANDELION, CLOVER, HENBIT, PLANTAINS, WILD ONION, AND
MANY OTHER BROADLEAF WEEDS.
ALSO FOR HIGHWAYS, RIGHTS-OF-WAY AND OTHER SIMILAR NON-CROP AREAS

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA
1 GALLON TREATS (COVERS) UP TO 40,000 SQ. FT.

GET THE OPTICAL ADVANTAGE[®]

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

AUG 8 2006
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-489

ACTIVE INGREDIENTS:

Trisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	13.24%
Dimethylamine Salt of Mecoprop-p,**†	2.29%
Dicamba Acid***	0.77%

OTHER INGREDIENTS:	83.70%
TOTAL:	100.00%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	7.10%	0.618 lbs./gal.
**Mecoprop-p Acid	1.89%	0.164 lbs./gal.
***Dicamba Acid	0.64%	0.056 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-UIO
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527



NET CONTENTS GALS.

000228-00000.20060731.NUP 6J 04
Revisions per EPA 07/26/06.

9/14

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if inhaled. Avoid breathing spray mist. Wear long pants, long-sleeved shirt, socks, shoes, protective eyewear and chemical-resistant gloves when handling or applying this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pet on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters nor pour the washwater on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product can be used on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs), also along Highways, Rights-of-Way, Drainage Ditches and other similar turf areas. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Avoid broadcast applications when air temperature exceeds 90 degrees. When using small, spot treatment applications in temperatures over 90 degrees, turf injury may occur. When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 degrees. When air temperatures exceed 80 degrees, limit application to spot treatment only. Make new

10/14

dilution for each use. This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

Alder	Curly dock	Matchweed	Shepherdspurse
Annual yellow sweet clover	Curly indigo	Mexicanweed	Slender plantain
Artichoke	Dandelion	Milk vetch	Smallflower galinsoga
Aster	Dead Nettle	Milkweed bloodflower	Smartweed
Austrian fieldcress	Dock	Mugwort	Smooth dock
Bedstraw	Dogbane	Morningglory	Smooth pigweed
Beggartick	Dogfennel	Mouseear chickweed	Sneezweed southern wild rose
Biden	Elderberry	Musk thistle	Sowthistle
Bindweed	English daisy	Mustard	Spanishneedle
Bird vetch	Fall dandelion	Narrowleaf plantain	Spatterdock
Bitterweed	False dandelion	Narrowleaf vetch	Speedwell
Bitter wintercress	False flax	Nettle	Spiny amaranth
Black-eyed Susan	False sunflower	Nutgrass	Spiny cocklebur
Black medic	Fiddleneck	Orange hawkweed	Spotted catsear
Black mustard	Field bindweed	Oxalis	Spotted knapweed
Blackseed plantain	Field pansy	Oxeye daisy	Spotted spurge
Blessed thistle	Flea bane (Daisy)	Parsley-piert	Spurge
Blue lettuce	Flixweed	Parsnip	Spurweed
Blue vervain	Florida betony	Pearlwort	Stinging nettle
Box elder	Florida pusley	Pennycress	Stinkweed
Bracted plantain	Frenchweed	Pennywort	Stichwort
Brassbuttons	Galinsoga	Peppergrass	Strawberry clover
Bristly oxtongue	Garlic mustard	Pepperweed	Sumac
Broadleaf dock	Goathead	Pigweed	Sunflower
Broadleaf plantain	Goatsbeard	Pineywoods bedstraw	Sweet clover
Broomweed	Goldenrod	Plains coeopsis	Tall nettle
Buckhorn	Ground ivy	Plantain	Tall vervain
Buckhorn plantain	Gumweed	Poison hemlock	Tansy mustard
Bulbous buttercup	Hairy bittercress	Poison ivy	Tansy ragwort
Bull nettle	Hairy fleabane	Poison oak	Tanweed
Bull thistle	Hawkweed	Pokeweed	Tarweed
Burdock	Healall	Poorjoe	Thistle
Burning nettle	Heartleaf drymary	Povertyweed	Tick trefoil
Bur ragweed	Hedge bindweed	Prickly lettuce	Toadflax
Burweed	Hedge mustard	Prickly sida	Trailing crown vetch
Buttercup	Hemp	Primrose	Tumble mustard
Canada thistle	Henbit	Prostrate knotweed	Tumble pigweed
Carolina geranium	Hoary cress	Prostrate pigweed	Tumbleweed
Carpetweed	Hoary plantain	Prostrate spurge	Velvet leaf
Catchweed bedstraw	Hoary vervain	Prostrate vervain	Venice mallow
Catsear	Honeysuckle	Puncture vine	Veronica
Catnip	Hop clover	Purslane	Vervain
Chickweed	Horsenettle	Ragweed	Vetch
Chicory	Horsetail	Red clover	Virginia buttonweed
Cinquefoil	Indiana mallow	Redroot pigweed	Virginia creeper
Clover	Ironweed	Red sorrel	Virginia pepperweed
Cockle	Jewelweed	Redstem filaree	Wavyleaf bullthistle
Cocklebur	Jimsonweed	Rough cinquefoil	Western clematis
Coffeebean	Kochia	Rough fleabane	Western salsify
Coffeeweed	Knawel	Roundleafed mangold	White clover
Common chickweed	Knotweed	Rush	White mustard
Common mullein	Lambsquarter	Russian pigweed	Wild mustard
Common sowthistle	Lespedeza	Russian thistle	Wild aster
Corn chamomile	Locoweed	St. Johnswort	Wild buckwheat
Creeping Jenny	Lupine	Scarlet pimpernel	Wild carrot
Crimson clover, croton	Mallow	Scotch thistle	Wild four-o'clock,
Cudweed	Marshelder	Sheep sorrel	Wild geranium

11/14

Wild lettuce	Wild strawberry	Woolly morningglory	Yarrow
Wild marigold	Wild sweet potato	Woodsorrel	Yellow Rocket
Wild parsnip	Wild vetch	Woolly croton	Yellow flower pepperweed
Wild radish	Willow	Woolly plantain	and other broadleaf weeds
Wild rape	Witchweed	Wormseed	

USE TIPS

1. Spray during the growing season when weeds are growing actively.
2. For best results, do not mow for 2 days before or after treating.
3. Do not treat when air temperatures exceed 90°F or damage to grass may occur.
4. For best results, do not water lawn within 24 hours after treatment. Poor weed control may result if rainfall occurs within 24 hours of application.
5. Shake well before using.
6. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
7. Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees. Clover will be damaged if sprayed.
8. Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.
9. If certain tough weeds are not dead within 3 weeks, repeat treatment. Broadcast application is not necessary; follow spot treatment directions.

HOW MUCH TO USE

SPOT TREATMENT

Pressure (Trigger) Sprayer: Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

ENTIRE LAWN

Hose-End Sprayer: Use a hose-end sprayer designed for lawns. Pour recommended amount of product (Column 2) in bottle and add water to indicated level (Column 3), and spray on area shown in Column 4. For example, to spray 2,500 square feet measure 20 fluid ounces (40 Tablespoons) in jar, add water to the 10 gallon level and mix. Turn on garden hose and spray as evenly as possible over 2,500 square feet.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

P/14

NOTE: Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

ORNAMENTAL LAWNS AND TURF

Pressure Sprayer: Measure indicated amount of this product, pour into tank, and add proper amount of water according to the following chart. Use a spray pattern with coarse spray droplets without any fine mist.

AMOUNT OF CONCENTRATE REQUIRED TO TREAT 1 ACRE		
GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER
Cool Season Grasses Bluegrass Fescue Ryegrass	2.75 Gallons (350 ounces)	175 Gallons
Warm Season Grasses Bahagrass Common Bermudagrass Zoysiagrass	1.375 Gallons (175 ounces)	175 Gallons
Warm Season Grasses Centipedegrass St. Augustinegrass*	0.6875 Gallons (87.5 ounces)	175 Gallons

*Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

NON-TURF AREAS

For roadsides (including aprons and guardrails), rights-of-way, drainage ditchbanks, and other similar non-crop areas mix this product at a rate of 0.75 to 4.5 gallons per 50 to 300 gallons of water. This will cover 1 acre. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until this product has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.

The maximum seasonal application rate to non-crop sites is 4 pounds 2,4-D acid equivalent per acre per application site.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep from freezing. to be stored in original container and placed in areas inaccessible to children.

PESTICIDE DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning, if burned, stay out of smoke.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Undesirable plant injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Nufarm or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Nufarm and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. NUFARM MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Nufarm, and Buyer assumes the risk of any such use.

To the extent allowed by law, Nufarm or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF NUFARM AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF NUFARM OR SELLER, THE REPLACEMENT OF PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

(New)

Triplet and The Optical Advantage are Registered Trademarks of Nufarm Americas, Inc.

[OPTIONAL BOX]

PRODUCT FACTS	
What it Does: Kills all Major Broadleaf Weeds including Chickweed, Clover and Dandelion (See inside for complete list)	
Where to Use: (Icon weeds) Lawns: Kills the Weeds not the Lawn	
When to Use: Do not apply when daytime temperatures are above 90°F as lawn injury can occur.	
Questions and Comments: (Icon phone) (icon computer) Call xxxxxxxx or visit our website at www.xxxxxxxx	

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING HYPE - FRONT PANEL

- Patented Low Odor Formulation
- Low Odor Formulation
- Now With Patented Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds
- Controls Over 250 Weeds
- Use When Weeds Are Actively Growing
- May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
- Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
- Overnight Results
- Works in 24 Hours
- Kills the Weeds, Not the Lawn
- Rainproof in Hours - Won't Wash Away
- Colorless Formulation
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Contains Triplet Low Odor Formula
- Easy to Use

14/14

For Thick, Weed-Free Turf
Three Way Control
Apply to a Moist Lawn
Contains The Three Way Power of Triple Low Odor
One Quart Covers Up To 10,000 Square Feet (One XXX covers up to XX,XXX square feet)
Easy to Mix
Just Add Water
New Formulation
Specially formulated for residential use.

ALTERNATE HOSE-END SPRAYER INSTRUCTIONS

How To Use: (Illustrations)

1. Attach garden hose to spray nozzle.
2. Turn on water supply.
3. Bend safety tab down and turn control to "water"
4. Point nozzle toward lawn and turn control to "on". Product will automatically mix with water. Walk back and forth at a steady pace.