



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

SEP 19 2008

Nufarm Americas, Inc.  
150 Harvester Dr., Suite 200  
Burr Ridge, IL 60527

Subject: 2,4-D RED Label Amendment  
Triplet Low Odor Premium Lawn Weed Killer Concentrate  
EPA Reg. No. 228-489

The labeling referred to above is in compliance with the 2,4-D RED provided that you:

1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following changes to the product labeling:

The following comments apply to the Residential labeling:

1. Revise the first paragraph of the Environmental Hazards section to the following:  
"This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate."
2. Under Use Precautions, add the statement "Do not apply as a fine mist because of potential for injury to desirable plants."
3. Under Use Precautions, add the statement "Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only."
4. Under Use Precautions, add the statements "Do not apply more than twice per year. A minimum of 30 days is required between treatments."
5. Under the Storage and Disposal section, include directions for "Container Disposal" in accordance with PR Notice 2007-4 for household/residential uses.

The following comments apply to the **Commercial** labeling:

6. Revise the PPE section to the following:

“All mixers, loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks,
- chemical-resistant gloves (except for applicators using ground boom equipment), and
- chemical resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.”

7. Under the User Safety Requirements, add the statement “Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product’s concentrate. Do not reuse them.”

8. The first sentence under the Engineering Control Statements section must be deleted from the label. In addition, the mechanical transfer text (the remainder of the first paragraph) may be removed from the label.

9. Revise the first sentence of the Environmental Hazards section to read “This pesticide is toxic to fish and aquatic invertebrates” and also delete the text “except as noted on appropriate labels” from this section.

10. With the exception of drift-related text appearing in the Environmental Hazards (“Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas”) and General Precautions and Restrictions (“Do not apply this product in a way that will contact workers or other persons, either directly or through drift”), all drift text appearing on the label must be placed together and be located below the Spray Drift Management text. Any conflicting text must be deleted from the label.

11. Include the following Spray Drift Management statement in the Use Precautions section.

### **SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind directions, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

### **Droplet Size**

Use a medium or coarser spray nozzles according to (ASAE standard 572) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

**Wind Speed**

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

**Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

**Susceptible Plants**

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

**Other State and Local Requirements**

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

**Equipment**

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

- 12. Under Directions for Use, revise the statement "Do not enter...until dusts have settled." to "Do not enter...until sprays have dried."
- 13. Under Directions for Use, include the statement "Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes."
- 14. Add the following statement to the labeling:  
  

"Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA Web Site: <http://www.epa.gov/espp>."

4/18

Page 4


EPA Reg. No. 228-489

15. Under Limitations for use on Ornamental Turf, revise the maximum per application rate of 2.5 gallons per acre to 2.4 gallons per acre, and revise the maximum seasonal rate to 4.8 gallons per acre. Also, delete the phrase "excluding spot treatments". Add the statement "A minimum of 30 days is required between applications."
16. Under Storage and Disposal, revise the heading "Pesticide Container Disposal" to "Container Disposal" and provide directions in accordance with PR Notice 2007-4.

Please Note: Final product reregistration cannot be considered until after all active ingredients in this product are eligible for reregistration.

A stamped copy of your labeling is enclosed for your records. Submit one (1) copy of the revised final printed label for the record before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions

Sincerely,



Joanne I. Miller  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

Enclosures

5/18

RESIDENTIAL

# TRIPLET<sup>®</sup> LOW ODOR PREMIUM LAWN WEED KILLER CONCENTRATE

SELECTIVE BROADLEAF WEED CONTROL RESIDENTIAL TURF WHEN USING A HOSE-END SPRAYER, JUST CONNECT HOSE, TURN ON WATER AND SPRAY.

1 QUART COVERS 8,000 SQUARE FEET

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

GET THE OPTICAL ADVANTAGE<sup>®</sup>

**ACTIVE INGREDIENTS:**

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	13.24%
Dimethylamine Salt of Mecoprop-p,**‡	2.29%
Dicamba Acid***	0.64%
<b>OTHER INGREDIENTS:</b>	<b>83.83%</b>
<b>TOTAL:</b>	<b>100.00%</b>

Isomer Specific Method, Equivalent to:

*2,4-D Acid	7.10%	0.618 lbs./gal.
**Mecoprop-p Acid	1.89%	0.164 lbs./gal.
***Dicamba Acid	0.64%	0.056 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-489  
EPA EST. NO. 228-IL-1

MANUFACTURED BY  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



NET CONTENTS      GALS.

000228-00489.20071107.EPA24D.Pending

**ACCEPTED**  
with **COMMENTS**  
In EPA Letter Dated:  
SEP 19 2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-489

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if inhaled. Avoid breathing spray mist. Wear long pants, long-sleeved shirt, socks, shoes, protective eyewear and chemical-resistant gloves when handling or applying this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Remove saturated clothing as soon as possible and shower.

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

**ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

**USE PRECAUTIONS**

**IMPORTANT:** Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. If necessary, contact your local Extension Agent.

WEEDS CONTROLLED

Alder  
 Annual yellow sweet clover  
 Artichoke  
 Aster  
 Austrian fieldcress  
 Bedstraw  
 Beggartick  
 Bident  
 Bindweed  
 Bird vetch  
 Bitterweed  
 Bitter wintercress  
 Black-eyed Susan  
 Black medic  
 Black mustard  
 Blackseed plantain  
 Blessed thistle  
 Blue lettuce  
 Blue vervain  
 Box elder  
 Bracted plantain  
 Brassbuttons  
 Bristly oxtongue  
 Broadleaf dock  
 Broadleaf plantain  
 Broomweed  
 Buckhorn  
 Buckhorn plantain  
 Bulbous buttercup  
 Bull nettle  
 Bull thistle  
 Burdock  
 Burning nettle  
 Bur ragweed  
 Burweed  
 Buttercup  
 Canada thistle  
 Carolina geranium  
 Carpetweed  
 Catchweed bedstraw  
 Catsear  
 Catnip  
 Chickweed  
 Chicory  
 Cinquefoil  
 Clover  
 Cockle  
 Cocklebur  
 Coffeebean  
 Coffeeweed  
 Common chickweed  
 Common mullein  
 Common sowthistle  
 Corn chamomile  
 Creeping Jenny  
 Crimson clover, croton  
 Cudweed  
 Curly dock  
 Curly indigo  
 Dandelion  
 Dead Nettle  
 Dock

Dogbane  
 Dogfennel  
 Elderberry  
 English daisy  
 Fall dandelion  
 False dandelion  
 False flax  
 False sunflower  
 Fiddleneck  
 Field bindweed  
 Field pansy  
 Flea bane (Daisy)  
 Flixweed  
 Florida betony  
 Florida pusley  
 Frenchweed  
 Galinsoga  
 Garlic mustard  
 Goathead  
 Goatsbeard  
 Goldenrod  
 Ground ivy  
 Gumweed  
 Hairy bittercress  
 Hairy fleabane  
 Hawkweed  
 Healall  
 Heartleaf drymary  
 Hedge bindweed  
 Hedge mustard  
 Hemp  
 Henbit  
 Hoary cress  
 Hoary plantain  
 Hoary vervain  
 Honeysuckle  
 Hop clover  
 Horsenettle  
 Horsetail  
 Indiana mallow  
 Ironweed  
 Jewelweed  
 Jimsonweed  
 Kochia  
 Knawel  
 Knotweed  
 Lambsquarter  
 Lespedeza  
 Locoweed  
 Lupine  
 Mallow  
 Marshelder  
 Matchweed  
 Mexicanweed  
 Milk vetch  
 Milkweed bloodflower  
 Mugwort  
 Morningglory  
 Moussear chickweed  
 Musk thistle  
 Mustard  
 Narrowleaf plantain

Narrowleaf vetch  
 Nettle  
 Nutgrass  
 Orange hawkweed  
 Oxalis  
 Oxeye daisy  
 Parsley-piert  
 Parsnip  
 Pearlwort  
 Pennygrass  
 Pennywort  
 Peppergrass  
 Pepperweed  
 Pigweed  
 Pineywoods bedstraw  
 Plains coeopsis  
 Plantain  
 Poison hemlock  
 Poison ivy  
 Poison oak  
 Pokeweed  
 Poorjoe  
 Povertyweed  
 Prickly lettuce  
 Prickly sida  
 Primrose  
 Prostrate knotweed  
 Prostrate pigweed  
 Prostrate spurge  
 Prostrate vervain  
 Puncture vine  
 Purslane  
 Ragweed  
 Red clover  
 Redroot pigweed  
 Red sorrel  
 Redstem filaree  
 Rough cinquefoil  
 Rough fleabane  
 Roundleaved marigold  
 Rush  
 Russian pigweed  
 Russian thistle  
 St. Johnswort  
 Scarlet pimpernel  
 Scotch thistle  
 Sheep sorrel  
 Shepherdspurse  
 Slender plantain  
 Smallflower galinsoga  
 Smartweed  
 Smooth dock  
 Smooth pigweed  
 Sneezeweed southern wild rose  
 Sowthistle  
 Spanishneedle  
 Spatterdock  
 Speedwell  
 Spiny amaranth  
 Spiny cocklebur  
 Spotted catsear  
 Spotted knapweed

Spotted spurge  
 Spurge  
 Spurweed  
 Stinging nettle  
 Stinkweed  
 Stichwort  
 Strawberry clover  
 Sumac  
 Sunflower  
 Sweet clover  
 Tall nettle  
 Tall vervain  
 Tansy mustard  
 Tansy ragwort  
 Tanweed  
 Tarweed  
 Thistle  
 Tick trefoil  
 Toadflax  
 Trailing crown vetch  
 Tumble mustard  
 Tumble pigweed  
 Tumbleweed  
 Velvet leaf  
 Venice mallow  
 Veronica  
 Vervain  
 Vetch  
 Virginia buttonweed  
 Virginia creeper  
 Virginia pepperweed  
 Wavyleaf bullthistle  
 Western clematis  
 Western salsify  
 White clover  
 White mustard  
 Wild mustard  
 Wild aster  
 Wild buckwheat  
 Wild carrot  
 Wild four-o'clock  
 Wild geranium  
 Wild lettuce  
 Wild marigold  
 Wild parsnip  
 Wild radish  
 Wild rape  
 Wild strawberry  
 Wild sweet potato  
 Wild vetch  
 Willow  
 Witchweed  
 Woolly morningglory  
 Woodsorrel  
 Woolly croton  
 Woolly plantain  
 Wormseed  
 Yarrow  
 Yellow Rocket  
 Yellow flower pepperweed  
 and other broadleaf weeds

**USE TIPS**

1. Spray during the growing season when weeds are growing actively.
2. For best results, do not mow for 2 days before or after treating.
3. Do not treat when air temperatures exceed 90°F or damage to grass may occur.
4. For best results, do not water lawn within 24 hours after treatment. Poor weed control may result if rainfall occurs within 24 hours of application.
5. Shake well before using.
6. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
7. Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees. Clover will be damaged if sprayed.
8. Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.
9. If certain tough weeds are not dead within 3 weeks, repeat treatment. Broadcast application is not necessary; follow spot treatment directions.

**HOW MUCH TO USE**

**SPOT TREATMENT**

**Pressure (Trigger) Sprayer:** Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
<b>Cool Season Grasses</b> Bluegrass Fescue Ryegrass	4 Fluid Ounces (8 Tablespoons)	1 Gallon	1,000 Square Feet
<b>Warm Season Grasses</b> Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	2 Fluid Ounce (4 Tablespoons)	1 Gallon	1,000 Square Feet

**ENTIRE LAWN**

**Hose-End Sprayer:** Use a hose-end sprayer designed for lawns. Pour recommended amount of product (Column 2) in bottle and add water to indicated level (Column 3), and spray on area shown in Column 4. For example in cool season grasses, to spray 4,000 square feet measure 16 fluid ounces (32 Tablespoons) in jar, add water to the 4 gallon level and mix. Turn on garden hose and spray as evenly as possible over 4,000 square feet.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
<b>Cool Season Grasses</b> Bluegrass Fescue Ryegrass	4 Fluid Ounces (8 Tablespoons)	1 Gallon	1,000 Square Feet
<b>Warm Season Grasses</b> Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	2 Fluid Ounce (4 Tablespoons)	1 Gallon	1,000 Square Feet

**NOTE:** Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.



**Pressure Sprayer:** Measure indicated amount of this product, pour into tank, and add proper amount of water according to the following chart. Use a spray pattern with coarse spray droplets without any fine mist.

AMOUNT OF CONCENTRATE IN 1 GALLON OF WATER REQUIRED TO TREAT 1,000 SQUARE FEET	
GRASS TYPE	AMOUNT OF CONCENTRATE
<b>Cool Season Grasses</b> Bluegrass Fescue Ryegrass	4 Fluid Ounces (8 Tablespoons)
<b>Warm Season Grasses</b> Bahiagrass Common Bermudagrass Zoysiagrass	2 Fluid Ounces (4 Tablespoons)
<b>Warm Season Grasses</b> Centipedegrass St. Augustinegrass*	1 Fluid Ounce (2 Tablespoons)

\*Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep from freezing. to be stored in original container and placed in areas inaccessible to children.

**PESTICIDE DISPOSAL:** **If empty** - Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled** - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(110707A)

[OPTIONAL BOX]

PRODUCT FACTS

**What it Does:**  
 Kills all Major Broadleaf Weeds including Chickweed, Clover and Dandelion (See inside for complete list)

**Where to Use: (Icon weeds)**  
 Lawns: Kills the Weeds not the Lawn

**When to Use:**  
 Do not apply when daytime temperatures are above 90°F as lawn injury can occur.

**Questions and Comments: (Icon phone) (icon computer)**  
 Call xxxxxxxx or visit our website at www.xxxxxxxx

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon.	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING HYPE - FRONT PANEL

- Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds
- Controls Over 250 Weeds
- Use When Weeds Are Actively Growing
- May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
- Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
- Overnight Results
- Works in 24 Hours
- Kills the Weeds, Not the Lawn
- Rainproof in Hours - Won't Wash Away
- Colorless Formulation
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Contains Triplet Low Odor Formula
- Easy to Use
- For Thick, Weed-Free Turf
- Three Way Control
- Apply to a Moist Lawn
- Contains The Three Way Power of Triple Low Odor
- One Gallon Covers 32,000 Square Feet (One XXX covers XX,XXX square feet)
- Easy to Mix
- Just Add Water
- New Formulation
- Specially formulated for residential use.

ALTERNATE HOSE-END SPRAYER INSTRUCTIONS

How To Use: (Illustrations)

1. Attach garden hose to spray nozzle.
2. Turn on water supply.
3. Bend safety tab down and turn control to "water"
4. Point nozzle toward lawn and turn control to "on". Product will automatically mix with water. Walk back and forth at a steady pace.

11/18

COMMERCIAL

# TRIPLLET® LOW ODOR PREMIUM LAWN WEED KILLER CONCENTRATE

SELECTIVE BROADLEAF WEED CONTROL FOR ORNAMENTAL TURFGRASSES. TO CONTROL DANDELION, CLOVER, HENBIT, PLANTAINS, WILD ONION, AND MANY OTHER BROADLEAF WEEDS.

ALSO FOR HIGHWAYS, RIGHTS-OF-WAY AND OTHER SIMILAR NON-CROP AREAS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

1 GALLON TREATS (COVERS) 32,000 SQ. FT.

GET THE OPTICAL ADVANTAGE®

**ACTIVE INGREDIENTS:**

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	13.24%
Dimethylamine Salt of Mecoprop-p,**†	2.29%
Dicamba Acid***	0.64%

**OTHER INGREDIENTS:**

TOTAL: 83.83%  
100.00%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	7.10%	0.618 lbs./gal.
**Mecoprop-p Acid	1.89%	0.104 lbs./gal.
***Dicamba Acid	0.04%	0.056 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-489  
EPA EST. NO. 228-IL-1

MANUFACTURED BY  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



NET CONTENTS      GALS.

**ACCEPTED**  
with **COMMENTS**  
In EPA Letter Dated:  
SEP 19 2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

000228-00489.20071107.EPA24D.Pending

228-489

12/18

COMMERCIAL

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if inhaled. Avoid breathing spray mist.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**All loaders, applicators, and other handlers must wear:**

- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical resistant gloves, when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry

**Engineering Control Statements:**

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE may be reduced or modified as specified in the WPS.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	
<ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>	

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

<b>HOT LINE NUMBER</b>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled.

This product can be used on Ornamental Turf Lawns (Residential, Industrial and Institutional) Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs), also along Highways, Rights-of-Way, Drainage Ditches and other similar turf areas.

**USE PRECAUTIONS**

**IMPORTANT:** Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperatures over 90°F, turf injury may occur. When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80°F. When air temperatures exceed 80°F, limit application to spot treatment only. Make new dilution for each use. This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

**WEEDS CONTROLLED**

- |                            |                        |                   |                   |
|----------------------------|------------------------|-------------------|-------------------|
| Alder                      | Bulbous buttercup      | Cudweed           | Gumweed           |
| Annual yellow sweet clover | Bull nettle            | Curly dock        | Hairy bittercress |
| Artichoke                  | Bull thistle           | Curly indigo      | Hairy fleabane    |
| Aster                      | Burdock                | Dandelion         | Hawkweed          |
| Austrian fieldcress        | Burning nettle         | Dead Nettle       | Healall           |
| Bedstraw                   | Bur ragweed            | Dock              | Heartleaf drymary |
| Beggartick                 | Burweed                | Dogbane           | Hedge bindweed    |
| Biden                      | Buttercup              | Dogfennel         | Hedge mustard     |
| Bindweed                   | Canada thistle         | Elderberry        | Hemp              |
| Bird vetch                 | Carolina geranium      | English daisy     | Henbit            |
| Bitterweed                 | Carpetweed             | Fall dandelion    | Hoary cress       |
| Bitter wintercress         | Catchweed bedstraw     | False dandelion   | Hoary plantain    |
| Black-eyed Susan           | Catsear                | False flax        | Hoary vervain     |
| Black medic                | Catnip                 | False sunflower   | Honeysuckle       |
| Black mustard              | Chickweed              | Fiddleneck        | Hop clover        |
| Blackseed plantain         | Chicory                | Field bindweed    | Horsenettle       |
| Blessed thistle            | Cinquefoil             | Field pansy       | Horsetail         |
| Blue lettuce               | Clover                 | Flea bane (Daisy) | Indiana mallow    |
| Blue vervain               | Cockle                 | Flixweed          | Ironweed          |
| Box elder                  | Cocklebur              | Florida betony    | Jewelweed         |
| Bracted plantain           | Coffeebean             | Florida pusley    | Jimsonweed        |
| Brassbuttons               | Coffeeweed             | Frenchweed        | Kochia            |
| Bristly oxtongue           | Common chickweed       | Galinsoga         | Knawel            |
| Broadleaf dock             | Common mullein         | Garlic mustard    | Knotweed          |
| Broadleaf plantain         | Common sowthistle      | Goathead          | Lambsquarter      |
| Broomweed                  | Corn chamomile         | Goatsbeard        | Lespedeza         |
| Buckhorn                   | Creeping Jenny         | Goldenrod         | Locoweed          |
| Buckhorn plantain          | Crimson clover, croton | Ground ivy        |                   |

(Continued)

14/18

**WEEDS CONTROLLED (Continued)**

Lupine	Poorjoe	Spanishneedle	Vetch
Mallow	Povertyweed	Spatterdock	Virginia buttonweed
Marshelder	Prickly lettuce	Speedwell	Virginia creeper
Matchweed	Prickly sida	Spiny amaranth	Virginia pepperweed
Mexicanweed	Primrose	Spiny cocklebur	Wavyleaf bullthistle
Milk vetch	Prostrate knotweed	Spotted catsear	Western clematis
Milkweed bloodflower	Prostrate pigweed	Spotted knapweed	Western salsify
Mugwort	Prostrate spurge	Spotted spurge	White clover
Morningglory	Prostrate vervain	Spurge	White mustard
Mouseear chickweed	Puncture vine	Spurweed	Wild mustard
Musk thistle	Purslane	Stinging nettle	Wild aster
Mustard	Ragweed	Stinkweed	Wild buckwheat
Narrowleaf plantain	Red clover	Stichwort	Wild carrot
Narrowleaf vetch	Redroot pigweed	Strawberry clover	Wild four-o'clock,
Nettle	Red sorrel	Sumac	Wild geranium
Nutgrass	Redstem filaree	Sunflower	Wild lettuce
Orange hawkweed	Rough cinquefoil	Sweet clover	Wild marigold
Oxalis	Rough fleabane	Tall nettle	Wild parsnip
Oxeye daisy	Roundleaved marigold	Tall vervain	Wild radish
Parsley-piert	Rush	Tansy mustard	Wild rape
Parsnip	Russian pigweed	Tansy ragwort	Wild strawberry
Pearlwort	Russian thistle	Tanweed	Wild sweet potato
Pennycress	St. Johnswort	Tarweed	Wild vetch
Pennywort	Scarlet pimpernel	Thistle	Willow
Peppergrass	Scotch thistle	Tick trefoil	Witchweed
Pepperweed	Sheep sorrel	Toadflax	Woolly morningglory
Pigweed	Shepherdspurse	Trailing crown vetch	Woodsorrel
Pineywoods bedstraw	Slender plantain	Tumble mustard	Woolly croton
Plains coeopsis	Smallflower galinsoga	Tumble pigweed	Woolly plantain
Plantain	Smartweed	Tumbleweed	Wormseed
Poison hemlock	Smooth dock	Velvet leaf	Yarrow
Poison ivy	Smooth pigweed	Venice mallow	Yellow Rocket
Poison oak	Sneezeweed southern wild rose	Veronica	Yellow flower pepperweed
Pokeweed	Sowthistle	Vervain	and other broadleaf weeds

**USE TIPS**

1. Spray during the growing season when weeds are growing actively.
2. For best results, do not mow for 2 days before or after treating.
3. Do not treat when air temperatures exceed 90°F or damage to grass may occur.
4. For best results, do not water lawn within 24 hours after treatment. Poor weed control may result if rainfall occurs within 24 hours of application.
5. Shake well before using.
6. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
7. Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees. Clover will be damaged if sprayed.
8. Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.
9. If certain tough weeds are not dead within 3 weeks, repeat treatment. Broadcast application is not necessary; follow spot treatment directions.

15/18

### HOW MUCH TO USE

#### SPOT TREATMENT

**Pressure (Trigger) Sprayer:** Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
<b>Cool Season Grasses</b> Bluegrass Fescue Ryegrass	4 Fluid Ounces (8 Tablespoons)	1 Gallon	1,000 Square Feet
<b>Warm Season Grasses</b> Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	2 Fluid Ounce (4 Tablespoons)	1 Gallon	1,000 Square Feet

#### ENTIRE LAWN

**Hose-End Sprayer:** Use a hose-end sprayer designed for lawns. Pour recommended amount of product (Column 2) in bottle and add water to indicated level (Column 3), and spray on area shown in Column 4. For example in cool season grasses, to spray 4,000 square feet measure 16 fluid ounces (32 Tablespoons) in jar, add water to the 4 gallon level and mix. Turn on garden hose and spray as evenly as possible over 4,000 square feet.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
<b>Cool Season Grasses</b> Bluegrass Fescue Ryegrass	4 Fluid Ounces (8 Tablespoons)	1 Gallon	1,000 Square Feet
<b>Warm Season Grasses</b> Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	2 Fluid Ounce (4 Tablespoons)	1 Gallon	1,000 Square Feet

**NOTE:** Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

16/18

ORNAMENTAL LAWNS AND TURF

Pressure Sprayer: Measure indicated amount of this product, pour into tank, and add proper amount of water according to the following chart. Use a spray pattern with coarse spray droplets without any fine mist.

AMOUNT OF CONCENTRATE REQUIRED TO TREAT 1 ACRE		
GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER
<b>Cool Season Grasses</b> Bluegrass Fescue Ryegrass	1.375 Gallons (175 ounces)	43.5 Gallons
<b>Warm Season Grasses</b> Bahiagrass Common Bermudagrass Zoysiagrass	0.6875 Gallons (87.5 ounces)	43.5 Gallons
<b>Warm Season Grasses</b> Centipedegrass St. Augustinegrass*	0.3475 Gallons (43.75 ounces)	43.5 Gallons

\*Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

Limitations for Use on Ornamental Turf: The maximum rate per application is 2.5 gallons per acre, limited to 2 applications per year. The maximum seasonal rate is 5 gallons per acre, excluding spot treatments.

NON-TURF AREAS

For roadsides (including aprons and guardrails), rights-of-way, drainage ditchbanks, and other similar non-crop areas mix this product at a rate of 0.75 to 4.5 gallons per 50 to 300 gallons of water. This will cover 1 acre. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until this product has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.

Limitations for Non-Turf Areas:

For annual and perennial weeds, the maximum rate per application is 3.25 gallons per acre, limited to 2 applications per year. Minimum of 30 days between applications. For woody plants, the maximum rate per application is 6.5 gallons per acre, limited to 1 application per year.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep from freezing. to be stored in original container and placed in areas inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

PESTICIDE CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



17/18

COMMERCIAL

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(110707A)

Triplet and The Optical Advantage are Registered Trademarks of Nufarm Americas, Inc.

[OPTIONAL BOX]

PRODUCT FACTS

**What it Does:**  
 Kills all Major Broadleaf Weeds including Chickweed, Clover and Dandelion (See inside for complete list)

**Where to Use: (Icon weeds)**  
 Lawns: Kills the Weeds not the Lawn

**When to Use:**  
 Do not apply when daytime temperatures are above 90°F as lawn injury can occur.

**Questions and Comments: (Icon phone) (icon computer)**  
 Call xxxxxxxx or visit our website at www.xxxxxxxx

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING HYPE - FRONT PANEL

- Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds
- Controls Over 250 Weeds
- Use When Weeds Are Actively Growing
- May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
- Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
- Overnight Results
- Works in 24 Hours
- Kills the Weeds, Not the Lawn
- Rainproof in Hours - Won't Wash Away
- Colorless Formulation
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Contains Triplet Low Odor Formula
- Easy to Use
- For Thick, Weed-Free Turf
- Three Way Control
- Apply to a Moist Lawn
- Contains The Three Way Power of Triple Low Odor
- One Gallon Covers 32,000 Square Feet (One XXX covers XX,XXX square feet)
- Easy to Mix
- Just Add Water
- New Formulation
- Specially formulated for residential use.

ALTERNATE HOSE-END SPRAYER INSTRUCTIONS

- How To Use: (Illustrations)
1. Attach garden hose to spray nozzle.
  2. Turn on water supply.
  3. Bend safety tab and turn control to "water"
  4. Point nozzle toward lawn and turn control to "on". Product will automatically mix with water. Walk back and forth at a steady pace.