



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

228-492

Date of Issuance:

SEP - 8 2006

Term of Issuance:

Conditional

Name of Pesticide Product:

Triplet Low Odor Premium Granular Weed Killer

NOTICE OF PESTICIDE:

[x] Registration
[] Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Nufarm Americas Inc.
150 Harvester Drive Suite 200
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

- 1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit within one year of the date of this registration notice, storage stability (830.6317) and corrosion characteristics (830.6320) studies.
3. Make the following label changes before you release the product for shipment:
a. Add "Wear protective eyewear" to the PPE statements under the signal word CAUTION.
b. The Pesticide Disposal Statement (for the Commercial label) must have the following statement "Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility."
c. The Pesticide Container Disposal Statement (for the Commercial Label) must have the following statement "Paper and Plastic Bags: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke."

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Handwritten signature of Joanne I. Miller

Date:

SEP - 8 2006

2/12

4. Before the label is released for shipment, revise the EPA Registration Number to read, EPA Reg. No. 228-492.
5. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

3/12

(RESIDENTIAL LABEL)

TRIPLET[®] LOW ODOR PREMIUM GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE WHICH CONTROLS A WIDE SPECTRUM OF
BROADLEAF WEEDS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

WEEDS CONTACTED BY THIS PRODUCT GRADUALLY DIE AND DISAPPEAR FROM THE LAWN.

TREATS (OR COVERS) UP TO 5,000 SQUARE FEET (50' x 100')

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	1.6726%
Dimethylamine Salt of Mecoprop-p Acid**‡	0.2712%
Dicamba Acid***	0.0896%

OTHER INGREDIENTS:	<u>97.9666%</u>
TOTAL:	100.0000%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	0.8967%
**Mecoprop-p Acid	0.2241%
***Dicamba Acid	0.0896%

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

SEP - 8 2006
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

EPA REG. NO. 228-
EPA EST. NO. 228-IL-1

228-492

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS 20 LBS.

4/12

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water; also before eating, drinking, chewing gum, or using tobacco. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is a stable mixture of three herbicides. It is for use on residential turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WEEDS CONTROLLED

Alder	Bull thistle	Dead Nettle	Hedge bindweed
Annual yellow sweet clover	Burdock	Dock	Hedge mustard
Artichoke	Burning nettle	Dogbane	Hemp
Aster	Bur ragweed	Dogfennel	Henbit
Austrian fieldcress	Burweed	Elderberry	Hoary cress
Bedstraw	Buttercup	English daisy	Hoary plantain
Beggartick	Canada thistle	Fall dandelion	Hoary vervain
Biden	Carolina geranium	False dandelion	Honeysuckle
Bindweed	Carpetweed	False flax	Hop clover
Bird vetch	Catchweed bedstraw	False sunflower	Horsenettle
Bitterweed	Catsear	Fiddleneck	Horsetail
Bitter wintercress	Catnip	Field bindweed	Indiana mallow
Black-eyed Susan	Chickweed	Field pansy	Ironweed
Black medic	Chicory	Flea bane (Daisy)	Jewelweed
Black mustard	Cinquefoil	Flixweed	Jimsonweed
Blackseed plantain	Clover	Florida betony	Kochia
Blessed thistle	Cockle	Florida pusley	Knawel
Blue lettuce	Cocklebur	Frenchweed	Knotweed
Blue vervain	Coffeebean	Galinsoga	Lambsquarter
Box elder	Coffeeweed	Garlic mustard	Lespedeza
Bracted plantain	Common chickweed	Goathead	Locoweed
Brassbuttons	Common mullein	Goatsbeard	Lupine
Bristly oxtongue	Common sowthistle	Goldenrod	Mallow
Broadleaf dock	Corn chamomile	Ground ivy	Marshelder
Broadleaf plantain	Creeping Jenny	Gumweed	Matchweed
Broomweed	Crimson clover, croton	Hairy bittercress	Mexicanweed
Buckhorn	Cudweed	Hairy fleabane	Milk vetch
Buckhorn plantain	Curly dock	Hawkweed	Milkweed bloodflower
Bulbous buttercup	Curly indigo	Healall	Mugwort
Bull nettle	Dandelion	Heartleaf drymary	Morningglory

5/12

Mouseear chickweed	Prostrate spurge	Spotted catsear	Virginia pepperweed
Musk thistle	Prostrate vervain	Spotted knapweed	Wavyleaf bullthistle
Mustard	Puncture vine	Spotted spurge	Western clematis
Narrowleaf plantain	Purslane	Spurge	Western salsify
Narrowleaf vetch	Ragweed	Spurweed	White clover
Nettle	Red clover	Stinging nettle	White mustard
Nutgrass	Redroot pigweed	Stinkweed	Wild mustard
Orange hawkweed	Red sorrel	Stichwort	Wild aster
Oxalis	Redstem filaree	Strawberry clover	Wild buckwheat
Oxeye daisy	Rough cinquefoil	Sumac	Wild carrot
Parsley-piert	Rough fleabane	Sunflower	Wild four-o'clock,
Parsnip	Roundleafed marigold	Sweet clover	Wild geranium
Pearlwort	Rush	Tall nettle	Wild lettuce
Pennycress	Russian pigweed	Tall vervain	Wild marigold
Pennywort	Russian thistle	Tansy mustard	Wild parsnip
Peppergrass	St. Johnswort	Tansy ragwort	Wild radish
Pepperweed	Scarlet pimpernel	Tanweed	Wild rape
Pigweed	Scotch thistle	Tarweed	Wild strawberry
Pineywoods bedstraw	Sheep sorrel	Thistle	Wild sweet potato
Plains coeopsis	Shepherdspurse	Tick trefoil	Wild vetch
Plantain	Slender plantain	Toadflax	Willow
Poison hemlock	Smallflower galinsoga	Trailing crown vetch	Witchweed
Poison ivy	Smartweed	Tumble mustard	Woolly morningglory
Poison oak	Smooth dock	Tumble pigweed	Woodsorrel
Pokeweed	Smooth pigweed	Tumbleweed	Woolly croton
Poorjoe	Sneezeweed southern wild rose	Velvet leaf	Woolly plantain
Povertyweed	Sowthistle	Venice mallow	Wormseed
Prickly lettuce	Spanishneedle	Veronica	Yarrow
Prickly sida	Spatdock	Vervain	Yellow Rocket
Primrose	Speedwell	Vetch	Yellow flower pepperweed
Prostrate knotweed	Spiny amaranth	Virginia buttonweed	and other broadleaf weeds
Prostrate pigweed	Spiny cocklebur	Virginia creeper	

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

**SPREADER SETTINGS
ONE PASS APPLICATION:**

<u>MAKE</u>	<u>SETTING</u>
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all materials from surfaces immediately after application, because certain ingredients can cause permanent

6/12

stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(022006)

Triplet and The Optical Advantage are registered trademarks of Nufarm Americas Inc.

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

*NOTE: Guaranteed Spreader Settings will vary among various commercial manufacturers.

OPTIONAL MARKETING HYPE - FRONT PANEL

- Patented Low Odor Formulation
- Low Odor Formulation
- Now With Patented Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds

7/12

Controls Over 250 Weeds

Use When Weeds Are Actively Growing

May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass

Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More

Overnight Results

Works in 24 Hours

Kills the Weeds, Not the Lawn

Rainproof in Hours - Won't Wash Away

Colorless Formulation

Won't Harm the Home Lawn

Illustration(s) of various weeds listed on label

Contains Triplet Low Odor Formula

Easy to Use

For Thick, Weed-Free Turf

Three Way Control

Apply to a Moist Lawn

Contains The Three Way Power of Triple Low Odor

20 pound bag Covers Up To 5,000 Square Feet (XXX pound bag covers up to XX,XXX square feet.)

New Formulation

Specially formulated for residential use.

8/12

(COMMERCIAL LABEL)

TRIPLET[®] LOW ODOR PREMIUM GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE WHICH CONTROLS A WIDE SPECTRUM OF
BROADLEAF WEEDS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

WEEDS CONTACTED BY THIS PRODUCT GRADUALLY DIE AND DISAPPEAR FROM THE LAWN.

TREATS (OR COVERS) UP TO 5,000 SQUARE FEET (50' x 100')

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	1.6726%
Dimethylamine Salt of Mecoprop-p Acid** ‡	0.2712%
Dicamba Acid***	0.0896%

OTHER INGREDIENTS:	<u>97.9666%</u>
TOTAL:	100.0000%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	0.8967%
**Mecoprop-p Acid	0.2241%
***Dicamba Acid	0.0896%

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

SEP - 8 2006

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

EPA REG. NO. 228-
EPA EST. NO. 228-IL-1

228-492

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866



NET CONTENTS 20 LBS.

9/12

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water; also before eating, drinking, chewing gum, or using tobacco. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WEEDS CONTROLLED

Alder	Bull thistle	Dead Nettle	Hedge bindweed
Annual yellow sweet clover	Burdock	Dock	Hedge mustard
Artichoke	Burning nettle	Dogbane	Hemp
Aster	Bur ragweed	Dogfennel	Henbit
Austrian feldcress	Burweed	Elderberry	Hoary cress
Bedstraw	Buttercup	English daisy	Hoary plantain
Beggartick	Canada thistle	Fall dandelion	Hoary vervain
Biden	Carolina geranium	False dandelion	Honeysuckle
Bindweed	Carpetweed	False flax	Hop clover
Bird vetch	Catchweed bedstraw	False sunflower	Horsenettle
Bitterweed	Catsear	Fiddleneck	Horsetail
Bitter wintercress	Catnip	Field bindweed	Indiana mallow
Black-eyed Susan	Chickweed	Field pansy	Ironweed
Black medic	Chicory	Flea bane (Daisy)	Jewelweed
Black mustard	Cinquefoil	Flixweed	Jimsonweed
Blackseed plantain	Clover	Florida betony	Kochia
Blessed thistle	Cockle	Florida pusley	Knawel
Blue lettuce	Cocklebur	Frenchweed	Knotweed
Blue vervain	Coffeebean	Galinsoga	Lambsquarter
Box elder	Coffeeweed	Garlic mustard	Lespedeza
Bracted plantain	Common chickweed	Goathead	Locoweed
Brassbuttons	Common mullein	Goatsbeard	Lupine
Bristly oxtongue	Common sowthistle	Goldenrod	Mallow
Broadleaf dock	Corn chamomile	Ground ivy	Marshelder
Broadleaf plantain	Creeping Jenny	Gumweed	Matchweed
Broomweed	Crimson clover, croton	Hairy bittercress	Mexicanweed
Buckhorn	Cudweed	Hairy fleabane	Milk vetch
Buckhorn plantain	Curly dock	Hawkweed	Milkweed bloodflower
Bulbous buttercup	Curly indigo	Healall	Mugwort
Bull nettle	Dandelion	Heartleaf drymary	Morningglory

10/12

Mouseear chickweed	Prostrate spurge	Spotted catsear	Virginia pepperweed
Musk thistle	Prostrate vervain	Spotted knapweed	Wavyleaf bullthistle
Mustard	Puncture vine	Spotted spurge	Western clematis
Narrowleaf plantain	Purslane	Spurge	Western salsify
Narrowleaf vetch	Ragweed	Spurweed	White clover
Nettle	Red clover	Stinging nettle	White mustard
Nutgrass	Redroot pigweed	Stinkweed	Wild mustard
Orange hawkweed	Red sorrel	Stichwort	Wild aster
Oxalis	Redstem filaree	Strawberry clover	Wild buckwheat
Oxeye daisy	Rough cinquefoil	Sumac	Wild carrot
Parsley-piert	Rough fleabane	Sunflower	Wild four-o'clock,
Parsnip	Roundleaved marigold	Sweet clover	Wild geranium
Pearlwort	Rush	Tall nettle	Wild lettuce
Pennycress	Russian pigweed	Tall vervain	Wild marigold
Pennywort	Russian thistle	Tansy mustard	Wild parsnip
Peppergrass	St. Johnswort	Tansy ragwort	Wild radish
Pepperweed	Scarlet pimpernel	Tanweed	Wild rape
Pigweed	Scotch thistle	Tarweed	Wild strawberry
Pineywoods bedstraw	Sheep sorrel	Thistle	Wild sweet potato
Plains coeopsis	Shepherdspurse	Tick trefoil	Wild vetch
Plantain	Slender plantain	Toadflax	Willow
Poison hemlock	Smallflower galinsoga	Trailing crown vetch	Witchweed
Poison ivy	Smartweed	Tumble mustard	Woolly morningglory
Poison oak	Smooth dock	Tumble pigweed	Woodsorrel
Pokeweed	Smooth pigweed	Tumbleweed	Woolly croton
Poorjoe	Sneezeweed southern wild rose	Velvet leaf	Woolly plantain
Povertyweed	Sowthistle	Venice mallow	Wormseed
Prickly lettuce	Spanishneedle	Veronica	Yarrow
Prickly sida	Spatardock	Vervain	Yellow Rocket
Primrose	Speedwell	Vetch	Yellow flower pepperweed
Prostrate knotweed	Spiny amaranth	Virginia buttonweed	and other broadleaf weeds
Prostrate pigweed	Spiny cocklebur	Virginia creeper	

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed, raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

**SPREADER SETTINGS
ONE PASS APPLICATION:**

<u>MAKE</u>	<u>SETTING</u>
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all materials from surfaces immediately after application, because certain ingredients can cause permanent

11/12

stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: *If empty:* Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled:* If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(022006)

Triplet and The Optical Advantage are registered trademarks of Nufarm Americas Inc.

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

***NOTE:** Guaranteed Spreader Settings will vary among various commercial manufacturers.

OPTIONAL MARKETING HYPE - FRONT PANEL

- Patented Low Odor Formulation
- Low Odor Formulation
- Now With Patented Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds

12/12

Controls Over 250 Weeds
Use When Weeds Are Actively Growing
May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
Overnight Results
Works in 24 Hours
Kills the Weeds, Not the Lawn
Rainproof in Hours - Won't Wash Away
Colorless Formulation
Won't Harm the Home Lawn
Illustration(s) of various weeds listed on label
Contains Triplet Low Odor Formula
Easy to Use
For Thick, Weed-Free Turf
Three Way Control
Apply to a Moist Lawn
Contains The Three Way Power of Triple Low Odor
20 pound bag Covers Up To 5,000 Square Feet (XXX pound bag covers up to XX,XXX square feet.)
New Formulation