

228-492

09/17/2008

1/14



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

SEP 17 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Nufarm Americas Inc.
150 Harvester Drive
Suite 200
Burr Ridge, IL 60527

Subject: EPA Reg. 228-492 Triplet Low Odor Premium Granular Weed Killer / 2,4-D RED
Label Amendment

The labeling referred to above is in compliance with the 2,4-D RED provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- 2) Please note: Final product reregistration can't be considered until after all active ingredients in this product are eligible for reregistration.

Comment on the Commercial Label:

- 1) Per the 2,4-D, Dicamba and MCPP-p REDs, "chemical-resistant gloves made of any waterproof material" must be added to the handler PPE section.
- 2) The first 2 sentences of the Environmental Hazards section must be revised to read:

"This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark."

3) To the label add "Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes."

4) Per the product chemistry review, the Pesticide Disposal section must be revised to read:

"Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance."

5) Per the MCPP-p RED, a 30-day minimum retreatment interval must be added to the application restriction section of the label.

Page 2
EPA Reg. 228-492

- 6) To page 2, delete “and similar turf areas”.
- 7) To the label add “Do not apply more than two applications per year, including broadcast applications in combination with spot treatments.”

Comment on the Homeowner Label:

1) The first paragraph of the Environmental Hazards section of the label must be revised to read:

“This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate.”

2) Per the MCPP-p RED, a 30-day minimum retreatment interval must be added to the application restriction section of the label.


3) Per the MCPP-p RED, the following text must be added to the application restriction section of the label:

“Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.”

4) The text “Limited to 2 applications per year, including broadcast applications in combination with spot treatments.” must be added to the label.

Submit one copy of the revised final printed label for the record. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,



Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

3/14

COMMERCIAL

TRIPLER[®] LOW ODOR PREMIUM GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE
WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

WEEDS CONTACTED BY THIS PRODUCT GRADUALLY DIE AND DISAPPEAR FROM THE LAWN.

TREATS (OR COVERS) UP TO 5,000 SQUARE FEET (50' x 100')

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	1.6726%
Dimethylamine Salt of Mecoprop-p Acid**†	0.2712%
Dicamba Acid***	0.0896%

OTHER INGREDIENTS:	<u>97.9666%</u>
TOTAL:	100.0000%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	0.8967%
**Mecoprop-p Acid	0.2241%
***Dicamba Acid	0.0896%

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:
SEP 17 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-492

EPA REG. NO. 228-492
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS 20 LBS.

4/14

COMMERCIAL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WEEDS CONTROLLED

- | | | | |
|----------------------------|--------------------|--------------------|--------------------|
| Alder | Black-eyed Susan | Broadleaf plantain | Canada thistle |
| Annual yellow sweet clover | Black medic | Broomweed | Carolina geranium |
| Artichoke | Black mustard | Buckhorn | Carpetweed |
| Aster | Blackseed plantain | Buckhorn plantain | Catchweed bedstraw |
| Austrian fieldcress | Blessed thistle | Bulbous buttercup | Catsear |
| Bedstraw | Blue lettuce | Bull nettle | Catnip |
| Beggartick | Blue vervain | Bull thistle | Chickweed |
| Biden | Box elder | Burdock | Chicory |
| Bindweed | Bracted plantain | Burning nettle | Cinquefoil |
| Bird vetch | Brassbuttons | Bur ragweed | Clover |
| Bitterweed | Bristly oxtongue | Burweed | Cockle |
| Bitter wintercress | Broadleaf dock | Buttercup | Cocklebur |

(Continued)

COMMERCIAL

WEEDS CONTROLLED (Continued)

Coffeebean	Hop clover	Prickly sida	Tansy mustard
Coffeeweed	Horsenettle	Primrose	Tansy ragwort
Common chickweed	Horsetail	Prostrate knotweed	Tanweed
Common mullein	Indiana mallow	Prostrate pigweed	Tarweed
Common sowthistle	Ironweed	Prostrate spurge	Thistle
Corn chamomile	Jewelweed	Prostrate vervain	Tick trefoil
Creeping Jenny	Jimsonweed	Puncture vine	Toadflax
Crimson clover, croton	Kochia	Purslane	Trailing crown vetch
Cudweed	Knawel	Ragweed	Tumble mustard
Curly dock	Knotweed	Red clover	Tumble pigweed
Curly indigo	Lambsquarter	Redroot pigweed	Tumbleweed
Dandelion	Lespedeza	Red sorrel	Velvet leaf
Dead Nettle	Locoweed	Redstem filaree	Venice mallow
Dock	Lupine	Rough cinquefoil	Veronica
Dogbane	Mallow	Rough fleabane	Vervain
Dogfennel	Marshelder	Roundleafed marigold	Vetch
Elderberry	Matchweed	Rush	Virginia buttonweed
English daisy	Mexicanweed	Russian pigweed	Virginia creeper
Fall dandelion	Milk vetch	Russian thistle	Virginia pepperweed
False dandelion	Milkweed bloodflower	St. Johnswort	Wavyleaf bullthistle
False flax	Mugwort	Scarlet pimpernel	Western clematis
False sunflower	Morningglory	Scotch thistle	Western salsify
Fiddleneck	Mouseear chickweed	Sheep sorrel	White clover
Field bindweed	Musk thistle	Shepherdspurse	White mustard
Field pansy	Mustard	Slender plantain	Wild mustard
Flea bane (Daisy)	Narrowleaf plantain	Smallflower galinsoga	Wild aster
Flixweed	Narrowleaf vetch	Smartweed	Wild buckwheat
Florida betony	Nettle	Smooth dock	Wild carrot
Florida pusley	Nutgrass	Smooth pigweed	Wild four-o'clock
Frenchweed	Orange hawkweed	Sneezeweed southern wild rose	Wild geranium
Galinsoga	Oxalis	Sowthistle	Wild lettuce
Garlic mustard	Oxeye daisy	Spanishneedle	Wild marigold
Goathead	Parsley-piert	Spatterdock	Wild parsnip
Goatsbeard	Parsnip	Speedwell	Wild radish
Goldenrod	Pearlwort	Spiny amaranth	Wild rape
Ground ivy	Pennycress	Spiny cocklebur	Wild strawberry
Gumweed	Pennywort	Spotted catsear	Wild sweet potato
Hairy bittercress	Peppergrass	Spotted knapweed	Wild vetch
Hairy fleabane	Pepperweed	Spotted spurge	Willow
Hawkweed	Pigweed	Spurge	Witchweed
Healall	Pineywoods bedstraw	Spurweed	Woolly morningglory
Heartleaf drymary	Plains coeopsis	Stinging nettle	Woodsorrel
Hedge bindweed	Plantain	Stinkweed	Woolly croton
Hedge mustard	Poison hemlock	Stichwort	Woolly plantain
Hemp	Poison ivy	Strawberry clover	Wormseed
Henbit	Poison oak	Sumac	Yarrow
Hoary cress	Pokeweed	Sunflower	Yellow Rocket
Hoary plantain	Poorjoe	Sweet clover	Yellow flower pepperweed
Hoary vervain	Povertyweed	Tall nettle	and other broadleaf weeds
Honeysuckle	Prickly lettuce	Tall vervain	

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

6/14

COMMERCIAL

HOW TO APPLY: This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

**SPREADER SETTINGS
ONE PASS APPLICATION:**

MAKE	SETTING
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all materials from surfaces immediately after application, because certain ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Limitations: The maximum rate per application is 4 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 8 pounds per 1,000 square feet, excluding spot treatments.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay away from smoke.

7/14

COMMERCIAL

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(110807A)

Triplet and The Optical Advantage is a registered trademark of Nufarm Americas Inc.

8/14

COMMERCIAL

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

NOTE: Guaranteed Spreader Settings will vary among various commercial manufacturers.

OPTIONAL MARKETING HYPE - FRONT PANEL

Patented Low Odor Formulation
 Low Odor Formulation
 Now With Patented Low Odor Formulation
 Kills Over 250 Weed Types
 Kills 250+ Weed Types
 Kills Over 250 Weed Listed
 Kills Over 250 Weeds
 Kills All Major Broadleaf Weeds
 Kills Dandelion, Clover and Other Broadleaf Weeds
 Kills Dandelion, Clover and Other Listed Broadleaf Weeds
 Controls Over 250 Weeds
 Use When Weeds Are Actively Growing
 May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
 Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
 Overnight Results
 Works in 24 Hours
 Kills the Weeds, Not the Lawn
 Rainproof in Hours - Won't Wash Away
 Colorless Formulation
 Won't Harm the Home Lawn
 Illustration(s) of various weeds listed on label
 Contains Triplet Low Odor Formula
 Easy to Use
 For Thick, Weed-Free Turf
 Three Way Control
 Apply to a Moist Lawn
 Contains The Three Way Power of Triple Low Odor
 20 pound bag Covers Up To 5,000 Square Feet (XXX pound bag covers up to XX,XXX square feet.)
 New Formulation

9/14

RESIDENTIAL

TRIPLET[®] LOW ODOR PREMIUM GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE
WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

WEEDS CONTACTED BY THIS PRODUCT GRADUALLY DIE AND DISAPPEAR FROM THE LAWN.

TREATS (OR COVERS) UP TO 5,000 SQUARE FEET (50' x 100')

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	1.6726%
Dimethylamine Salt of Mecoprop-p Acid**‡	0.2712%
Dicamba Acid***	0.0896%

OTHER INGREDIENTS:	97.9666%
TOTAL:	100.0000%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	0.8967%
**Mecoprop-p Acid	0.2241%
***Dicamba Acid	0.0896%

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**


SEP 17 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-492

EPA REG. NO. 228-492
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS 20 LBS.

000228-00492.20071112.res.EPA24D.Pending

10/14

RESIDENTIAL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear, long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water; also before eating, drinking, chewing gum, or using tobacco.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

This product is a stable mixture of three herbicides. It is for use on residential turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WEEDS CONTROLLED

Alder	Broadleaf plantain	Coffebean	Field pansy.
Annual yellow sweet clover	Broomweed	Coffeeweed	Flea bane (Daisy)
Artichoke	Buckhorn	Common chickweed	Flixweed
Aster	Buckhorn plantain	Common mullein	Florida betony
Austrian fieldcress	Bulbous buttercup	Common sowthistle	Florida pusley
Bedstraw	Bull nettle	Corn chamomile	Frenchweed
Beggartick	Bull thistle	Creeping Jenny	Galinsoga
Biden	Burdock	Crimson clover, croton	Garlic mustard
Bindweed	Burning nettle	Cudweed	Goathead
Bird vetch	Bur ragweed	Curly dock	Goatsbeard
Bitterweed	Burweed	Curly indigo	Goldenrod
Bitter wintercress	Buttercup	Dandelion	Ground ivy
Black-eyed Susan	Canada thistle	Dead Nettle	Gumweed
Black medic	Carolina geranium	Dock	Hairy bittercress
Black mustard	Carpetweed	Dogbane	Hairy fleabane
Blackseed plantain	Catchweed bedstraw	Dogfennel	Hawkweed
Blessed thistle	Catsear	Elderberry	Healall
Blue lettuce	Catnip	English daisy	Heartleaf drymary
Blue vervain	Chickweed	Fall dandelion	Hedge bindweed
Box elder	Chicory	False dandelion	Hedge mustard
Bracted plantain	Cinquefoil	False flax	Hemp
Brassbuttons	Clover	False sunflower	Herbit
Bristly oxtongue	Cockle	Fiddleneck	Hoary cress
Broadleaf dock	Cocklebur	Field bindweed	Hoary plantain

(Continued)

11/14

RESIDENTIAL


WEEDS CONTROLLED (Continued)

Hoary vervain	Peppergrass	Smallflower galinsoga	Veronica
Honeysuckle	Pepperweed	Smartweed	Vervain
Hop clover	Pigweed	Smooth dock	Vetch
Horsenettle	Pineywoods bedstraw	Smooth pigweed	Virginia buttonweed
Indiana mallow	Plains coeopsis	Sneezeweed southern wild rose	Virginia creeper
Ironweed	Plantain	Sowthistle	Virginia pepperweed
Jewelweed	Poison hemlock	Spanishneedle	Wavyleaf bullthistle
Jimsonweed	Poison ivy	Spatterdock	Western clematis
Kochia	Poison oak	Speedwell	Western salsify
Knawel	Pokeweed	Spiny amaranth	White clover
Knotweed	Poorjoe	Spiny cocklebur	White mustard
Lambsquarter	Povertyweed	Spotted catsear	Wild mustard
Lespedeza	Prickly lettuce	Spotted knapweed	Wild aster
Locoweed	Prickly sida	Spotted spurge	Wild buckwheat
Lupine	Primrose	Spurge	Wild carrot
Mallow	Prostrate knotweed	Spurweed	Wild four-o'clock
Marshelder	Prostrate pigweed	Stinging nettle	Wild geranium
Matchweed	Prostrate spurge	Stinkweed	Wild lettuce
Mexicanweed	Prostrate vervain	Stichwort	Wild marigold
Milk vetch	Puncture vine	Strawberry clover	Wild parsnip
Milkweed bloodflower	Purslane	Sumac	Wild radish
Mugwort	Ragweed	Sunflower	Wild rape
Morningglory	Red clover	Sweet clover	Wild strawberry
Mouseear chickweed	Redroot pigweed	Tall nettle	Wild sweet potato
Musk thistle	Red sorrel	Tall vervain	Wild vetch
Mustard	Redstem filaree	Tansy mustard	Willow
Narrowleaf plantain	Rough cinquefoil	Tansy ragwort	Witchweed
Narrowleaf vetch	Rough fleabane	Tanweed	Woolly morningglory
Nettle	Roundleafed marigold	Tarweed	Woodsorrel
Nutgrass	Rush	Thistle	Woolly croton
Orange hawkweed	Russian pigweed	Tick trefoil	Woolly plantain
Oxalis	Russian thistle	Toadflax	Wormseed
Oxeye daisy	St. Johnswort	Trailing crown vetch	Yarrow
Parsley-piert	Scarlet pimpernel	Tumble mustard	Yellow Rocket
Parsnip	Scotch thistle	Tumble pigweed	Yellow flower pepperweed
Pearlwort	Sheep sorrel	Tumbleweed	and other broadleaf weeds
Pennycress	Shepherdspurse	Velvet leaf	
Pennywort	Slender plantain	Venice mallow	

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.



12/14

RESIDENTIAL

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

SPREADER SETTINGS ONE PASS APPLICATION:

MAKE	SETTING
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all materials from surfaces immediately after application, because certain ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Environmental Benefits

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL


Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.



14/14

RESIDENTIAL

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

NOTE: Guaranteed Spreader Settings will vary among various commercial manufacturers.

OPTIONAL MARKETING HYPE - FRONT PANEL

Patented Low Odor Formulation
 Low Odor Formulation
 Now With Patented Low Odor Formulation
 Kills Over 250 Weed Types
 Kills 250+ Weed Types
 Kills Over 250 Weed Listed
 Kills Over 250 Weeds
 Kills All Major Broadleaf Weeds
 Kills Dandelion, Clover and Other Broadleaf Weeds
 Kills Dandelion, Clover and Other Listed Broadleaf Weeds
 Controls Over 250 Weeds
 Use When Weeds Are Actively Growing
 May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
 Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
 Overnight Results
 Works in 24 Hours
 Kills the Weeds, Not the Lawn
 Rainproof in Hours - Won't Wash Away
 Colorless Formulation
 Won't Harm the Home Lawn
 Illustration(s) of various weeds listed on label
 Contains Triplet Low Odor Formula
 Easy to Use
 For Thick, Weed-Free Turf
 Three Way Control
 Apply to a Moist Lawn
 Contains The Three Way Power of Triple Low Odor
 20 pound bag Covers Up To 5,000 Square Feet (XXX pound bag covers up to XX,XXX square feet.)
 New Formulation
 Specially formulated for residential use.

