

228-493

11/21/2008

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

NOV 21 2008

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Nufarm Americas Inc.  
150 Harvester Drive  
Suite 200  
Burr Ridge, IL 60527

Subject: EPA Reg. 228-493 Triplet Low Odor LA Granular Weed Killer / 2,4-D RED Label Amendment

The labeling referred to above is in compliance with the 2,4-D RED and is amended under the RED provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- 2) Please note: Final product reregistration can't be considered until after all active ingredients in this product are eligible for reregistration.

**residential-use label:**

1) The following text must be added to the end of the first sentence under the Environmental Hazards section of the label:

“...and may adversely affect non-target plants.”

Also in this section, delete the text “except as noted on appropriate labels.”

2) The following text must be added to the label:

“Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.”

3) Under the How to Apply section, add the following text:

“The maximum rate per application is 4.4 pounds per 1000 sq.ft.  
Limited to 2 applications per year.”

Minimum of 30 days between applications.  
The maximum seasonal rate is 8.8 pounds per 1000 sq. ft.”

- 4) On page 3 delete “recommended” from “recommended rate”.

**commercial-use label:**

- 1) The PPE section must be revised to read as follows:

“Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are” (*registrant inserts correct chemical-resistant material*). “If you want more options, follow the instructions for category” [*registrant inserts A,B,C,D,E,F,G,or H*] “on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks, and
- chemical-resistant gloves (except for applicators using groundboom equipment)”

- 2) The Environmental Hazards text currently on the label must be revised to delete the text “except as noted on appropriate labels.”

- 3) Per the product chemistry review, the Storage and Disposal (Pesticide Disposal subheading) must include the following statement:

“Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.”

- 4) Add the following statement to the directions for use:

“Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.”

- 5) Under the How to Apply section, revise the “Limitations” text to read as follows:

“The maximum rate per application is 4.4 pounds per 1000 sq.ft.  
Limited to 2 applications per year.  
Minimum of 30 days between applications.


Page 3  
EPA Reg. 228-493

The maximum seasonal rate is 8.8 pounds per 1000 sq. ft.”

6) Delete “and similar turf areas” from page 2.

Submit one copy of the revised final printed label for the record. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,



Joanne Miller  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

4/15

COMMERCIAL

# TRIPLET<sup>®</sup> LOW ODOR L.A. GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE  
WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) up to 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE<sup>®</sup>

**ACTIVE INGREDIENTS:**

Trisopropanolamine Salt of 2,4-D Acid*	1.4420%
Dimethylamine Salt of Mecoprop-p Acid**†	0.2338%
Dicamba Acid***	0.0772%

**OTHER INGREDIENTS:**

TOTAL: 98.2470%  
100.0000%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	0.7728%
**Mecoprop-p Acid	0.1932%
***Dicamba Acid	0.0772%

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-493  
EPA EST. NO. 228-IL-1

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
NOV 21 2008**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.


228-493

MANUFACTURED BY  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



NET CONTENTS 20 LBS.

000228-00493.20071108.EPA24D.Pending



5/15

COMMERCIAL

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**All loaders, applicators, and other handlers must wear:**

- long-sleeved shirt and long pants,
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	
<ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>	

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

**ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.


**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.



## WEEDS CONTROLLED

Alder	Dogbane	Narrowleaf vetch	Spotted spurge
Annual yellow sweet clover	Dogfennel	Nettle	Spurge
Artichoke	Elderberry	Nutgrass	Spurweed
Aster	English daisy	Orange hawkweed	Stinging nettle
Austrian fieldcress	Fall dandelion	Oxalis	Stinkweed
Bedstraw	False dandelion	Oxeye daisy	Stichwort
Beggartick	False flax	Parsley-piert	Strawberry clover
Biden	False sunflower	Parsnip	Sumac
Bindweed	Fiddleneck	Pearlwort	Sunflower
Bird vetch	Field bindweed	Pennycress	Sweet clover
Bitterweed	Field pansy	Pennywort	Tall nettle
Bitter wintercress	Flea bane (Daisy)	Peppergrass	Tall vervain
Black-eyed Susan	Flixweed	Pepperweed	Tansy mustard
Black medic	Florida betony	Pigweed	Tansy ragwort
Black mustard	Florida pusley	Pineywoods bedstraw	Tanweed
Blackseed plantain	Frenchweed	Plains coeopsis	Tarweed
Blessed thistle	Galinsoga	Plantain	Thistle
Blue lettuce	Garlic mustard	Poison hemlock	Tick trefoil
Blue vervain	Goathead	Poison ivy	Toadflax
Box elder	Goatsbeard	Poison oak	Trailing crown vetch
Bracted plantain	Goldenrod	Pokeweed	Tumble mustard
Brassbuttons	Ground ivy	Poorjoe	Tumble pigweed
Bristly oxtongue	Gumweed	Povertyweed	Tumbleweed
Broadleaf dock	Hairy bittercress	Prickly lettuce	Velvet leaf
Broadleaf plantain	Hairy fleabane	Prickly sida	Venice mallow
Broomweed	Hawkweed	Primrose	Veronica
Buckhorn	Healall	Prostrate knotweed	Vervain
Buckhorn plantain	Heartleaf drymary	Prostrate pigweed	Vetch
Bulbous buttercup	Hedge bindweed	Prostrate spurge	Virginia buttonweed
Bull nettle	Hedge mustard	Prostrate vervain	Virginia creeper
Bull thistle	Hemp	Puncture vine	Virginia pepperweed
Burdock	Henbit	Purslane	Wavyleaf bullthistle
Burning nettle	Hoary cress	Ragweed	Western clematis
Bur ragweed	Hoary plantain	Red clover	Western salsify
Burweed	Hoary vervain	Redroot pigweed	White clover
Buttercup	Honeysuckle	Red sorrel	White mustard
Canada thistle	Hop clover	Redstem filaree	Wild mustard
Carolina geranium	Horsenettle	Rough cinquefoil	Wild aster
Carpetweed	Horsetail	Rough fleabane	Wild buckwheat
Catchweed bedstraw	Indiana mallow	Roundleaved marigold	Wild carrot
Catsear	Ironweed	Rush	Wild four-o'clock,
Catnip	Jewelweed	Russian pigweed	Wild geranium
Chickweed	Jimsonweed	Russian thistle	Wild lettuce
Chicory	Kochia	St. Johnswort	Wild marigold
Cinquefoil	Knawel	Scarlet pimpernel	Wild parsnip
Clover	Knotweed	Scotch thistle	Wild radish
Cockle	Lambsquarter	Sheep sorrel	Wild rape
Cocklebur	Lespedeza	Shepherdspurse	Wild strawberry
Coffeebean	Locoweed	Slender plantain	Wild sweet potato
Coffeeweed	Lupine	Smallflower galinsoga	Wild vetch
Common chickweed	Mallow	Smartweed	Willow
Common mullein	Marshelder	Smooth dock	Witchweed
Common sowthistle	Matchweed	Smooth pigweed	Woolly morningglory
Corn chamomile	Mexicanweed	Sneezeweed southern wild rose	Woodsorrel
Creeping Jenny	Milk vetch	Sowthistle	Woolly croton
Crimson clover, croton	Milkweed bloodflower	Spanishneedle	Woolly plantain
Cudweed	Mugwort	Spatterdock	Wormseed
Curly dock	Morningglory	Speedwell	Yarrow
Curly indigo	Mouseear chickweed	Spiny amaranth	Yellow Rocket
Dandelion	Musk thistle	Spiny cocklebur	Yellow flower pepperweed
Dead Nettle	Mustard	Spotted catsear	and other broadleaf weeds
Dock	Narrowleaf plantain	Spotted knapweed	

7/15

COMMERCIAL

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

**SPREADER SETTINGS  
ONE PASS APPLICATION\*:**

<u>MAKE</u>	<u>SETTING</u>
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all materials from surfaces immediately after application, because certain ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

**Limitations:** The maximum rate per application is 4.5 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 9 pounds per 1,000 square feet, excluding spot treatments.

**Environmental Benefits**

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

8/15

COMMERCIAL

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(110807A)

Triplet and The Optical Advantage is a registered trademark of Nufarm Americas Inc.



9/15

COMMERCIAL

**OPTIONAL SPREADER INSTRUCTIONS:**

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

**\*NOTE:** Guaranteed Spreader Settings will vary among various commercial manufacturers.

**OPTIONAL MARKETING HYPE - FRONT PANEL**

Patented Low Odor Formulation  
Low Odor Formulation  
Now With Patented Low Odor Formulation  
Kills Over 250 Weed Types  
Kills 250+ Weed Types  
Kills Over 250 Weed Listed  
Kills Over 250 Weeds  
Kills All Major Broadleaf Weeds  
Kills Dandelion, Clover and Other Broadleaf Weeds  
Kills Dandelion, Clover and Other Listed Broadleaf Weeds  
Controls Over 250 Weeds  
Use When Weeds Are Actively Growing  
May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass  
Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More  
Overnight Results  
Works in 24 Hours  
Kills the Weeds, Not the Lawn  
Rainproof in Hours - Won't Wash Away  
Won't Harm the Home Lawn  
Illustration(s) of various weeds listed on label  
Contains Triplet Low Odor Formula  
Easy to Use  
For Thick, Weed-Free Turf  
Three Way Control  
Apply to a Moist Lawn  
Contains The Three Way Power of Triplet Low Odor  
20 lb. bag covers up to 5,000 Square Feet (XXX pound bag covers up to XX,XXX square feet.)  
New Formulation

10/15

RESIDENTIAL

# TRIPLET<sup>®</sup> LOW ODOR L.A. GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE  
WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) up to 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE<sup>®</sup>

**ACTIVE INGREDIENTS:**

Triisopropanolamine Salt of 2,4-D Acid*	1.4420%
Dimethylamine Salt of Mecoprop-p Acid**‡	0.2338%
Dicamba Acid***	0.0772%

<b>OTHER INGREDIENTS:</b>	<b>98.2470%</b>
<b>TOTAL:</b>	<b>100.0000%</b>

Isomer Specific Method, Equivalent to:

*2,4-D Acid	0.7728%
**Mecoprop-p Acid	0.1932%
***Dicamba Acid	0.0772%

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-493  
EPA EST. NO. 228-IL-1

**ACCEPTED  
with COMMENTS**

In EPA Letter Dated:  
NOV 21 2000

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.


MANUFACTURED BY  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



228-493

NET CONTENTS 20 LBS.

000228-00493.20071112.EPA24D.Pending



11/15

RESIDENTIAL

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear, pants, shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water; also before eating, drinking, chewing gum, or using tobacco.

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

**ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

This product is a stable mixture of three herbicides. It is for use on residential turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. **DO NOT USE** on Dichondra and Carpet grass. **DO NOT USE** this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

**WEEDS CONTROLLED**

- |                            |                    |                        |                   |
|----------------------------|--------------------|------------------------|-------------------|
| Alder                      | Brassbuttons       | Chickweed              | Dogfennel         |
| Annual yellow sweet clover | Bristly oxtongue   | Chicory                | Elderberry        |
| Artichoke                  | Broadleaf dock     | Cinquefoil             | English daisy     |
| Aster                      | Broadleaf plantain | Clover                 | Fall dandelion    |
| Austrian fieldcress        | Broomweed          | Cockle                 | False dandelion   |
| Bedstraw                   | Buckhorn           | Cocklebur              | False flax        |
| Beggarick                  | Buckhorn plantain  | Coffeebean             | False sunflower   |
| Biden                      | Bulbous buttercup  | Coffeeweed             | Fiddleneck        |
| Bindweed                   | Bull nettle        | Common chickweed       | Field bindweed    |
| Bird vetch                 | Bull thistle       | Common mullein         | Field pansy       |
| Bitterweed                 | Burdock            | Common sowthistle      | Flea bane (Daisy) |
| Bitter wintercress         | Burning nettle     | Corn chamomile         | Flixweed          |
| Black-eyed Susan           | Bur ragweed        | Creeping Jenny         | Florida betony    |
| Black medic                | Burweed            | Crimson clover, croton | Florida pusley    |
| Black mustard              | Buttercup          | Cudweed                | Frenchweed        |
| Blackseed plantain         | Canada thistle     | Curly dock             | Galinsoga         |
| Blessed thistle            | Carolina geranium  | Curly indigo           | Garlic mustard    |
| Blue lettuce               | Carpetweed         | Dandelion              | Goathead          |
| Blue vervain               | Catchweed bedstraw | Dead Nettle            | Goatsbeard        |
| Box elder                  | Catsear            | Dock                   | Goldenrod         |
| Bracted plantain           | Catnip             | Dogbane                | Ground ivy        |

(Continued)

**WEEDS CONTROLLED (Continued)**

Gumweed	Nettle	Russian thistle	Tumble pigweed
Hairy bittercress	Nutgrass	St. Johnswort	Tumbleweed
Hairy fleabane	Orange hawkweed	Scarlet pimpernel	Velvet leaf
Hawkweed	Oxalis	Scotch thistle	Venice mallow
Healall	Oxeye daisy	Sheep sorrel	Veronica
Heartleaf drymary	Parsley-piert	Shepherdspurse	Vervain
Hedge bindweed	Parsnip	Slender plantain	Vetch
Hedge mustard	Pearlwort	Smallflower galinsoga	Virginia buttonweed
Hemp	Pennycress	Smartweed	Virginia creeper
Henbit	Pennywort	Smooth dock	Virginia pepperweed
Hoary cress	Peppergrass	Smooth pigweed	Wavyleaf bullthistle
Hoary plantain	Pepperweed	Sneezeweed southern wild rose	Western clematis
Hoary vervain	Pigweed	Sowthistle	Western salsify
Honeysuckle	Pineywoods bedstraw	Spanishneedle	White clover
Hop clover	Plains coeopsis	Spatterdock	White mustard
Horsenettle	Plantain	Speedwell	Wild mustard
Horsetail	Poison hemlock	Spiny amaranth	Wild aster
Indiana mallow	Poison ivy	Spiny cocklebur	Wild buckwheat
Ironweed	Poison oak	Spotted catsear	Wild carrot
Jewelweed	Pokeweed	Spotted knapweed	Wild four-o'clock,
Jimsonweed	Poorjoe	Spotted spurge	Wild geranium
Kochia	Povertyweed	Spurge	Wild lettuce
Knawel	Prickly lettuce	Spurweed	Wild marigold
Knotweed	Prickly sida	Stinging nettle	Wild parsnip
Lambsquarter	Primrose	Stinkweed	Wild radish
Lespedeza	Prostrate knotweed	Stichwort	Wild rape
Locoweed	Prostrate pigweed	Strawberry clover	Wild strawberry
Lupine	Prostrate spurge	Sumac	Wild sweet potato
Mallow	Prostrate vervain	Sunflower	Wild vetch
Marshelder	Puncture vine	Sweet clover	Willow
Matchweed	Purslane	Tall nettle	Witchweed
Mexicanweed	Ragweed	Tall vervain	Woolly morningglory
Milk vetch	Red clover	Tansy mustard	Woodsorrel
Milkweed bloodflower	Redroot pigweed	Tansy ragwort	Woolly croton
Mugwort	Red sorrel	Tanweed	Woolly plantain
Morningglory	Redstem filaree	Tarweed	Wormseed
Mouseear chickweed	Rough cinquefoil	Thistle	Yarrow
Musk thistle	Rough fleabane	Tick trefoil	Yellow Rocket
Mustard	Roundleafed marigold	Toadflax	Yellow flower pepperweed
Narrowleaf plantain	Rush	Trailing crown vetch	and other broadleaf weeds
Narrowleaf vetch	Russian pigweed	Tumble mustard	

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

13/15

RESIDENTIAL

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

SPREADER SETTINGS  
ONE PASS APPLICATION:

MAKE	SETTING
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all materials from surfaces immediately after application, because certain ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

**Environmental Benefits**

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- √ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

**STORAGE AND DISPOSAL**


Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

**PESTICIDE DISPOSAL: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.





15/15

**OPTIONAL SPREADER INSTRUCTIONS:**

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

**\*NOTE:** Guaranteed Spreader Settings will vary among various commercial manufacturers.

**OPTIONAL MARKETING HYPE - FRONT PANEL**

- Patented Low Odor Formulation
- Low Odor Formulation
- Now With Patented Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds
- Controls Over 250 Weeds
- Use When Weeds Are Actively Growing
- May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
- Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
- Overnight Results
- Works in 24 Hours
- Kills the Weeds, Not the Lawn
- Rainproof in Hours - Won't Wash Away
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Contains Triplet Low Odor Formula
- Easy to Use
- For Thick, Weed-Free Turf
- Three Way Control
- Apply to a Moist Lawn
- Contains The Three Way Power of Triplet Low Odor .
- 20 lb. bag covers up to 5,000 Square Feet (XXX pound bag covers up to XX,XXX square feet.)
- New Formulation
- Specially formulated for residential use.

