

228-498

4/11/2011

1 of 10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 11, 2011

Matthew Granahan
Regulatory Manager
Nufarm Americas, Inc.
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527

Subject: Amendment – Adding Application Rates
Bifenthrin 0.2% Granule Insecticide
EPA Reg. No. 228-498
Your submission dated November 10, 2010

Dear Mr. Granahan:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable subject to the comments listed below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

1. On the front panel replace "For use to control ants (including imported fire ants), mole crickets and other insect pests on lawns ... etc" with "For use to control ants (including imported fire ants), mole crickets and other listed insect pests on lawns ... etc".
2. Replace "Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco" with "Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet" in the *Precautionary Statement* section.
3. Under *Lawn and Perimeter Applications*, last paragraph, replace "May be applied at up to 0.4 pound a.i. per acre per application" with "Applied at up to 0.4 pound a.i. per acre per application".
4. Under Application Rates, first paragraph, replace "See specific pest for recommended level" with "See specific pest for application level".
5. Replace instance of "Food utensils such as spoons and measuring cups should not be used for food purposes after use in measuring pesticides" with "Food utensils such as spoons

and measuring cups must not be used for food purposes after use in measuring pesticides". Also, replace "Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticide" with "Food utensils such as teaspoons and measuring cups must not be used for food purposes after use with insecticide".

6. On page 5, second paragraph, replace, "See section under **Lawn Applications** for additional rate recommendations" with "See section under Lawn Applications for additional rate applications".
7. On page 5, replace the table title "**Recommended Spreader Settings For Broadcast Applications**" with "**Spreader Settings For Broadcast Applications**". Also replace the first sentence in the *Note* following the table, "These recommended setting should be used on starting points ... etc" with "These setting should be used on starting points ... etc".

If you have any questions regarding this action, please contact BeWanda Alexander at Alexander.bewanda@epa.gov or (703) 305-7460.

Sincerely,

A handwritten signature in cursive script that reads "BeWanda Alexander for".

Richard Gebken
Product Manager
Insecticide Branch
Registration Division (7505P)

Enclosure

BIFENTHRIN 0.2% GRANULE INSECTICIDE

FOR USE TO CONTROL ANTS (INCLUDING IMPORTED FIRE ANTS), MOLE
CRICKETS AND OTHER INSECT PESTS ON LAWNS IN LANDSCAPED AREAS
AND PERIMETERS AROUND RESIDENTIAL, INSTITUTIONAL, PUBLIC,
COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL
AREAS AND ATHLETIC FIELDS.

ACTIVE INGREDIENT:

Bifenthrin* 0.2%

OTHER INGREDIENTS: 99.8%

TOTAL: 100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with COMMENTS
In EPA Letter Dated
APR 11 2011
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
228-498

NOTE: Spanish language is optional

EPA REG. NO. 228-498
EPA EST. NO.

Manufactured For
NUFARM AMERICAS INC.
150 Harvester Drive
Burr Ridge, IL 60527



NET WEIGHT: Lbs. (Kg)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION / PRECAUCION**

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN This product is a pyrethroid. If large amounts have been ingested, milk, cream or other digestible fats and oils may increase absorption and so should be avoided.	

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run-off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

Do not broadcast apply more than 0.4 pounds a.i. (200 pounds of this product) per acre per year [0.008 pounds a.i. (4 pounds of this product) per 1,000 sq. ft. per year]. However, for residential lawn use, do not apply more than 0.2 pounds a.i. per acre per application [0.004 pounds a.i. (2 pounds of this product) per 1,000 sq. ft. per application].

Not for use on golf courses, sod farms, nurseries, in commercial greenhouses, or on grass grown for seed.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run-off.

Do not make applications during rain.

LAWN AND PERIMETER APPLICATIONS

Lawn Application: Broadcast this product with suitable application equipment to ensure uniform coverage over the treatment area.

Do not apply more than 0.2 pound active ingredient (a.i.) per acre per application [0.004 pounds a.i. (2 pounds of this product) per 1,000 sq. ft. per application] on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 0.4 pound a.i. per acre per application [0.008 pounds a.i. (4 pounds of this product) per 1,000 sq. ft. per year] on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

APPLICATION RATES

The application rates listed in the following table will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, this product may be applied at up to 0.4 pounds a.i./acre [0.008 pounds a.i. (4 pounds of this product)/1,000 sq. ft.] on non-residential use sites to control each of the pests listed in the table. See specific pest for recommended level. Irrigation or watering by 0.1 to 0.5 inches immediately after application generally improves the effectiveness of this product.

PEST	ACTIVE INGREDIENT POUNDS PER ACRE	ACTIVE INGREDIENT POUNDS PER 1,000 SQ. FT.	APPLICATION RATE OF THIS PRODUCT	
Armyworms ¹ Cutworms ¹ Sod Webworm ¹	0.1 pound a.i. per acre	0.002 pound a.i. per 1,000 sq. ft.	50 pounds per acre	1.15 pounds per 1,000 sq. ft.
Annual Bluegrass Weevils (<i>Hyperodes</i>) (Adult) ² Billbugs (Adult) ³ Black Turfgrass Ataenius (Adult) ⁴ Leafhoppers Mealybugs	0.1 to 0.2 pound a.i. per acre	0.002 to 0.004 pound a.i. per 1,000 sq. ft.	50 to 100 pounds per acre	1.15 to 2.3 pounds per 1,000 sq. ft.
PEST	ACTIVE INGREDIENT POUNDS PER ACRE	ACTIVE INGREDIENT POUNDS PER 1,000 SQ. FT.	APPLICATION RATE OF THIS PRODUCT	
European Crane Flies	0.15 to 0.3 pound a.i. per acre	0.003 to 0.006 pound a.i. per 1,000 sq. ft.	75 to 150 pounds per acre	1.75 to 3.5 pounds per 1,000 sq. ft.
Ants Centipedes Chinch Bugs ⁵ Fleas (Larvae) ⁶ Fleas (Adult) Imported Fire Ants (Adult) ⁷ Imported Fire Ants (Mounds) ⁷ Millipedes Mole Cricket (Adult) ⁸ Mole Cricket (Nymph) ⁸ Scorpions ¹⁰ Ticks ¹¹	0.2 to 0.4 pound a.i. per acre	0.004 to 0.008 pound a.i. per 1,000 sq. ft.	100 to 200 pounds per acre	2.3 to 4.6 pounds per 1,000 sq. ft.
Pillbugs Sowbugs	0.4 pound a.i. per acre	0.008 pound a.i. per 1,000 sq. ft.	200 pounds per acre	4.6 pounds per 1,000 sq. ft.

In New York State, this product may NOT be applied to any grass, or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

¹**Armyworms, Cutworms, and Sod Webworms:** To ensure optimum control, irrigate the treated area with up to 0.1 inches of water immediately after application to activate (release from the granule) the insecticide.

²**Annual Bluegrass Weevil (*Hyperodes*) adults:** Applications should be timed to control adult weevils as they leave their over-wintering sites and move into grass areas. This movement generally begins when *Forsythia* is in full bloom and concludes when Flowering Dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

³**Billbug Adults:** Applications should be made when adult Billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting Billbug adults will also provide control of over-wintered Chinch Bugs.

⁴**Black Turfgrass Ataenius Adults:** Applications should be made during May and July to control the first and second generation of Black Turfgrass Ataenius adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and Horse Chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus synacus*).

⁹Chinch Bugs: Chinch Bugs infest the base of grass plants and are often found in the thatch layer. Irrigate the treated area with up to 0.25 inches of water immediately after application to activate (release from the granule) the insecticide. Chinch Bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

⁹Flea Larvae: Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Irrigate the treated area with up to 0.5 inches of water immediately after application to activate (release from the granule) the insecticide.

¹Imported Fire Ants: Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound treatments that will control existing colonies. If the soil is not moist, then it is important to irrigate before application. Broadcast treatments should apply 0.4 pound a.i./acre [0.008 pounds a.i. (4 pounds of this product)/1,000 sq. ft.]. Mounds should be treated by applying 1/2 cup of this product per mound and then drenching the mound with 1 to 2 gallons of water. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the Ant tunnels. A four-foot diameter circle around the mound should also be treated. For best results, apply in cool weather (65 to 80°F) or in early morning or late evening hours. Food utensils such as spoons and measuring cups should not be used for food purposes after use in measuring pesticides. Application of 125 lbs./acre [2.85 lbs./1,000 sq. ft.] of this product will provide 6 months residual activity for the control of foraging fire ants and newly mated fly-in queens.

⁹Mole Cricket Adults: Achieving acceptable control of adult Mole Crickets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the Mole Crickets closer to the soil surface where contact with the insecticide will be maximized. Grass areas that receive pressure from adult Mole Crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

⁹Mole Cricket Nymphs: Grass areas that received intense adult Mole Cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher application rates and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the Mole Crickets closer to the soil surface where contact with the insecticide will be maximized.

¹⁰Scorpions: To ensure optimum control, treat the building perimeter at dusk to prevent outdoor scorpions from entering the building. Scorpions reside hidden in cracks and voids during the day and are active at night at temperatures above 77°F/25°C. Resting areas can be identified at night using a blacklight (UV bulb) as scorpions will fluoresce. These areas should also be treated. Prior to treatment, to increase treatment efficacy, remove trash, debris, or firewood that scorpions may use as resting sites.

¹¹Ticks (Including ticks that may transmit Lyme Disease and Rocky Mountain Spotted fever): Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher application rates when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Do not allow public use of treated areas during application.

Deer ticks (ixodes sp.) have a complicated life cycle that ranges over a two-year period and involves four life stages. Applications should be made in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter.

American dog ticks may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid spring to early fall to control American dog tick larvae, nymphs and adults.

Perimeter Application: Apply to ground area 5 to 10 feet wide around and adjacent to the structure.

Boundary Applications: Apply to soil areas, ornamental gardens, pavement crevices, fencelines, planter areas, vegetation areas and mulch areas.

APPLICATION RATE POUNDS PER 1,000 SQ. FT.	BAND WIDTH	POUNDS OF THIS PRODUCT FOR PERIMETER APPLICATIONS LINEAR DISTANCE (FEET) AROUND STRUCTURE									
		25	50	100	150	200	250	300	400	500	1,000
2.3	5 feet	0.3	0.6	1.3	1.9	2.5	3.0	3.6	4.8	5.9	11.7
2.3	10 feet	0.7	1.5	3.0	4.2	5.3	6.5	7.6	9.9	12.2	23.7
3.0	5 feet	0.4	0.8	1.7	2.5	3.3	3.9	4.7	6.3	7.7	15.2
3.0	10 feet	0.9	1.9	3.9	5.4	6.9	8.5	9.9	12.9	15.9	30.9
3.5	5 feet	0.5	1.0	2.0	2.9	3.8	4.5	5.4	7.2	8.9	17.5
3.5	10 feet	1.1	2.2	4.5	6.3	8.0	9.8	11.4	14.9	18.3	35.5
4.6	5 feet	0.6	1.3	2.6	3.8	5.0	6.0	7.2	9.6	11.8	23.4
4.6	10 feet	1.3	3.0	6.0	8.4	10.6	13.0	15.2	19.8	24.4	47.4

Note: The amounts of this product listed in the above table are based on the total area being treated. This area is determined by multiplying the linear distance by the band width and then adding to that value the area of a circle whose radius is equal to the band width (to account for the area treated at the four corners of a rectangular structure).

Fire Ant Mound Treatment: Apply 1/2 cup of this product per mound and then drench the mound with 1 to 2 gallons of water. The mound should be drenched with sufficient force to break its apex and allow the insecticide to flow into the ant tunnels. For best results, apply in cool weather (65 to 80°F) or in early morning or late evening hours. Treat new mounds as they appear. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticide.

INDUSTRIAL NON-AGRICULTURAL APPLICATIONS
(Including Industrial Sites, Parks, Ball Fields and Cemeteries)

Broadcast Spot Applications: For the control of actively foraging ants (including Imported Fire Ants) broadcast uniformly 2.3 to 4.6 pounds of this product per 1,000 square feet (0.2 to 0.4 pounds a.i./acre). See section under **Lawn Applications** for additional rate recommendations.

Pad Mounted Electrical Sites: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 1.0 to 1.5 tablespoons (0.06 to 0.09 pounds) of this product per 4 square feet of mounting pad. Where existing transformers are completely or partially filled with soil excavated by resident ants, remove any excess fill and level the remaining soil prior to application. **Food utensils such as tablespoons must not be used for food purposes after use in measuring pesticides.**

Note: Electrical transformers appear to be more attractive to ants during the cooler months. For best results treat in the fall or early spring.

RECOMMENDED SPREADER SETTINGS FOR BROADCAST APPLICATIONS

EQUIPMENT	Application Rate lbs./1,000 square feet			PATTERN SETTING	WIDTH
	1.15	2.30	4.60		
LESCO Broadcast	F	G	I	-	10
Prizelawn CBR-III	E 1/2	F 1/2	H 1/2	-	10
Scotts RBA	G 1/2	H 1/2	J 1/2	3	10
Scotts SR2000	H 1/2	I 1/2	L 1/2	8	12
Scotts Handy Green	2	3	4	-	10
Scotts Accupro	4-1/2	4-3/4	5	8	12
Spyker	3-1/2	3-3/4	4-1/4	-	10
Vicon 402*	25	28-1/2	38	-	40
Lely W*	3-3/4	4-1/2	5	III-B	52
Cycloneway	1.5	3	5-3/4	-	4
Earthway 2400	8-1/2	10-1/2	12-1/2	-	10

*Speed applied: Vicon 5.6 mph, Lely 4.5 mph

Note: These recommended settings should be used as starting points when calibrating a spreader to achieve the desired application rate of this product. After calibration, the technician should regularly observe the amount of this product that is applied to a known area and make further setting adjustments, as necessary, to maintain the appropriate application rate. Spreaders should be calibrated periodically to ensure that equipment wear or other factors have not altered the flow rate of this product. Calibration instructions are included in the Owner's Manual that was provided by the equipment manufacturer at the time of purchase.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

CONTAINER DISPOSAL: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV111010)

Optional Language:

Controls Ticks!

Broadcast or Mount Treatment

LABEL HISTORY

FILE NAME	REVISION MARK	COMMENTS
000228-00494.20100408.EPA.Notification	RV040810N	PRN2007-4; 98-01 & Pyrethroid homeowner products
000228-00494.20100623.MASTER	RV062310	EPA Approval of changes from RV040810N
000228-00494.20100624.EPA.Notification	RV062410	PRN98-01
000228-00494.20100929.MASTER	RV092910	EPA Approval of changes from RV062410
000228-00494.20101110.EPA.Notification	RV111010	PRN98-01