22	8-506	10/23/	2006	Pape 17	7
A UNITED STATES Office		MENTAL PROTECTIC AGENCY Pesticide Programs	DN EPA Reg. Number: 228-506	Date of Issuance:	
The PROTECTION	Arie 1200 Pen	Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460		OCT 2 3 2006	
			Term of Issuance:	Conditional	
	NOTICE OF PESTICIDE: <u>x</u> Registration		Name of Pesticide F		
	Reregistration (under FIFRA, as amended)	Triplet Low Premium W (18/5)	Odor L.A. eed and Feed		
Name and Address Nufarm Ameri 150 Harvester Burr Ridge, IL	Drive Suite 200	IP Code):			
Note: Changes in labelir	ng differing in substance from t	hat accepted in connection with	his registration must be submitted us product diways felse to the above	to and accepted by the	
right to exclusive use of This product is provided that yo 1. Subm the A	the name or to its use if it has b s conditionally regist u: .it and/or cite all data gency requires all re	tered in accordance w a required for registrat gistrants of similar pro	ith FIFRA sec. 3(c)(7)(ith FIFRA sec. 3(c)(7)(cion/reregistration of yc oducts to submit such c tration notice, storage s	A) and (B) our product when lata.	
and c	_	tics (830.6320) studie			
	0	•	o read, "EPA Reg. No.	228-506.	
	nit one copy of the re uct for shipment.	evised final printed la	bel for the record befor	e you release the	
Signature of Approving	Official:		Date:		
Joanne I. Miller Product Manage Herbicide Branc Registration Div	r 23 Joann	. J. mille	r l	CT 2 3 2005	

page 2 EPA Reg. No. 228-506

287

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505P)

TRIPLET® LOW ODOR L.A. WEED AND FEED (18/5)

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) up to 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS: Triisopropanolamine Sall of 2,4-D Acid* Dimethylamine Sall of Mccoprop-p Acid**‡ Dicamba Acid***	
OTHER INGREDIENTS:	<u>98.5541%</u> TOTAL: 100.0000%
Isomer Specific Method, Equivalent to: *2,4-D Acid **Mecoprop-p Acid **Dicamba Acid \$\$CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.	
GUARANTEED ANALYSIS:° Total Nitrogen (N) X% Ammoniacal Nitrogen X% Urea Nitrogen Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from (List sources) Chlorine (CI) Maximum The urea nitrogen in this product has been coated to provide X% c Fertilizer guaranteed by (Name of fertilizer registrant)	

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-EPA EST. NO. 228-IL-1 MANUFACTURED BY NUFARM AMERICAS INC. BURR RIDGE, IL 60527-0866



ACCEPTED with COMMENTS In EPA Letter Dated:

OCT 2 3 2006 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

228-506

NET CONTENTS 18 LBS.

000228-00000.20060410.NUP 7K 04

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear pants, shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water; also before eating, drinking, chewing gum, or using tobacco. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

	FIRST AID
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
	HOT LINE NUMBER container or label with you when calling a poison control center or doctor, or going for treatment. You may also i-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is a stable mixture of three herbicides. It is for use on residential turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WEEDS CONTROLLED

Alder Annual yellow sweet clover Artichoke Aster Austrian fieldcress Bedstraw Beggartick Biden Bindweed Bird vetch Bitterweed Bitter wintercress Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Blue lettuce Blue vervain Box elder Bracted plantain Brassbuttons Bristly oxtongue Broadleaf dock Broadleaf plantain Broomweed Buckhorn Buckhorn plantain Bulbous buttercup Bull nettle **Bull thistle**

Burdock Burning nettle Bur ragweed Burweed Buttercup Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catsear Cathip Chickweeed Chicory Cinquetoil Clover Cockle Cocklebur Coffeebean Coffeeweed Common chickweeed Common mullein Common sowthistle Corn chamomile Creeping Jenny Crimson clover, croton Cudweed Curly dock Curly indigo Dandelion Dead Nettle Dock

Dogbane Dogfennel Elderberry English daisy Fall dandelion False dandelion False flax False sunflower Fiddleneck Field bindweed Field pansy Flea bane (Daisy) Flixweed Florida betony Florida pusley Frenchweed Galinsoga Garlic mustard Goathead Goatsbeard Goldenrod Ground ivv Gumweed Hairy bittercress Hairy fleabane Hawkweed Healall Heartleaf drymary Hedge bindweed Hedge mustard Нетр

Henbit Hoary cress Hoary plantain Hoary vervain Honeysuckle Hop clover Horsenettle Horsetail Indiana mallow Ironweed Jewelweed Jimsonweed Kochia Knawel Knotweed Lambsquarter Lespedeza Locoweed Lupine Mallow Marshelder Matchweed Mexicanweed Milk vetch Milkweed bloodflower Mugwort Morningglory Mouseear chickweed Musk thistle Mustard Narrowleal plantain

Narrowleaf vetch Nettle Nutgrass Orange hawkweed Oxalis Oxeye daisy Parsley-piert Parsnip Pearlwort Pennycress Pennywort Peppergrass Pepperweed Pigweed Pineywoods bedstraw Plains coeopsis Plantain Poison hemlock Poison ivv Poison oak Pokeweed Poorjoe Povertyweed Prickly lettuce Prickly sida Primrose Prostrate knolweeed Prostrate pigweed Prostrate spurge Prostrate vervain Puncture vine

Purslane Ragweed Red clover Redroot pigweed Red sorrel Redstern filaree Rough cinquefoil Rough fleabane Roundleafed marigold Rush Russian pigweed Russian thistle St. Johnswort Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smartweed Smooth dock Smooth pigweed Sneezeweed southern wild rose Sowthistle Spanishneedle Spatterdock Speedwell Spiny amaranth Spiny cocklebur Spotted catsear Spotted knapweed

Spotted spurge Spurge Spurweed Stinging nettle Stinkweed Stichwort Strawberry clover Sumac Sunflower Sweet clover Tall nettle Tall vervain Tansy mustard Tansy ragwort Tanweed Tarweed Thistle Tick trefoil Toadflax Trailing crown vetch Tumble mustard Tumble pigweed Tumbleweed Velvet leaf Venice mallow Veronica Vervain Vetch Virginia buttonweeed Virginia creeper Virginia pepperweed

Wavyleaf bullthistle Western clematis Western salsify White clover White mustard Wild mustard Wild aster Wild buckwheat Wild carrot Wild four-o'clock, Wild geranium Wild lettuce Wild marigold Wild parsnip Wild radish Wild rape Wild strawberry Wild sweet potato Wild vetch Willow Witchweed Wooily morningglory Woodsorrel Woolly croton Woolly plantain Wormseed Yarrow Yellow Rocket Yellow flower pepperweed and other broadleaf weeds

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, overapplication may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 18 pound bag should be used on 5,000 square feet. Apply eventy with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 3.6 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and uneveness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square leet and adjust to apply at the rate of .9 pound per 250 square feet.

SPREADER SETTINGS ONE PASS APPLICATION':

MAKE	<u>SETTING</u>
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all tertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen/1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (No. of Ibs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂0 levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer

program.

*Based on inert fertilizer claimed.

Environmental Benefits

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

(Grasscycling Logo)

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more

readily so nutrients go back into your

- A 50' x 50' lawn produces enough oygen for a family of 4 every day
- v Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & Other Pollutants
- ✓ Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

lawn.

PESTICIDE STORAGE: Always use original container to store pesticides and keep any bags containing unused porduct in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(NEW)

7

Triplet and The Optical Advantage are registered trademarks of Nularm Americas Inc.

OPTIONAL SPREADER INSTRUCTIONS:

- (Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.
- (Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following note will not be part of the actual label copy.

'NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State fertilizer regulation.

7 7 7

OPTIONAL MARKETING HYPE - FRONT PANEL Patented Low Odor Formulation Low Odor Formulation Now With Patented Low Odor Formulation Kills Over 250 Weed Types Kills 250+ Weed Types Kills Over 250 Weed Listed Kills Over 250 Weeds Kills All Major Broadleaf Weeds Kills Dandelion, Clover and Other Broadleat Weeds Kills Dandelion, Clover and Other Listed Broadleaf Weeds Controls Over 250 Weeds Use When Weeds Are Actively Growing May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More **Overnight Results** Works in 24 Hours Kills the Weeds, Not the Lawn Rainproof in Hours - Won't Wash Away Won't Harm the Home Lawn Illustration(s) of various weeds listed on label Contains Triplet Low Odor Formula Easy to Use For Thick, Weed-Free Turf Three Way Control Apply to a Moist Lawn Contains The Three Way Power of Triple Low Odor 18 lb. Bag Treats Up To 5,000 Square Feet (XXX bag covers up to XX,XXX square feet.) New Formulation Specially formulated for residential use.