

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

SEP - 8 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Nufarm Americas Inc. 150 Harvester Drive Suite 200 Burr Ridge, IL 60527

Subject: EPA Reg. 228-506 (Triplet Low Odor LA Premium Weed and Feed)/ 2,4-D RED Label Amendment

The labeling referred to above is in compliance with the 2,4-D RED provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- 2) Please note: Final product reregistration can't be considered until after all active ingredients in this product are eligible for reregistration.
- 3) The first paragraph of the Environmental Hazard section must be revised to read: "This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate."
- 4) The following text must be added to the label: "Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated areas(s) to the point of runoff or apply when raining or when rain is expected that day."
- 5) To the Directions for Use add "Do not apply more than 2 times per year, including broadcast applications in combination with spot treatments. Applications must be at least 30 days apart."
- 6) The Hazards to Humans and Domestic Animals must be revised to read as specified in the bold type below: "CAUTION Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, shoes and socks. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water, also before eating, drinking, chewing gum, or using tobacco."

Page 2 EPA Reg. 228-506

Submit one copy of the revised final printed label for the record. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

Joanne Miller

Product Manager 23

Herbicide Branch

Registration Division (7505P)

J. Miller

TRIPLET[®] LOW ODOR L.A. PREMIUM WEED AND FEED (18/5)

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) up to 5,000 Square Feet (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:	
Triisopropanolamine Salt of 2,4-D Acid*	
Dimethylamine Salt of Mecoprop-p Acid**‡	
Dicamba Acid***	
OTHER INGREDIENTS:	
	TOTAL: 100.0000%
Isomer Specific Method, Equivalent to:	
*2,4-D Acid	0.6376%
**Mecoprop-p Acid	0.1594%
***Dicamba Acid	
‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.	
GUARANTEED ANALYSIS.°	
Total Nitrogen (N)	X%
X% Ammoniacal Nitrogen	
X% Urea Nitrogen	•
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Primary Plant Foods derived from (List sources)	•
Chlorine (CI) Maximum	
The urea nitrogen in this product has been coated to provide X% coated	
Fertilizer guaranteed by (Name of fertilizer registrant)	-
3	

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED with COMMENTS

EPA REG. NO. 228-506 EPA EST. NO. 228-IL-1 In EPA Letter Dated: MANUFACTURED BY NUFARM AMERICAS INC

SEP - 8 2008 150 HARVESTER DRIVE
Under the Federal Insecticide.
BURR RIDGE, IL 60527
Fungicide, and Rodenticide Act
as amended, for the pesticide

NET CONTENTS

18 LBS.

228-506

registered under EPA Reg. No.



000228-00506.20071112.EPA24D.Pending: 20

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear pants, shirt, socks and shoes when applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water; also before eating, drinking, chewing gum, or using tobacco.

FIRST AID				
IF IN EYES	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 			
	HOT LINE NUMBER container or label with you when calling a poison control center or doctor, or going for treatment. You may also -1840 for emergency medical treatment information.			

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

This product is a stable mixture of three herbicides. It is for use on residential turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WEEDS CONTROLLED

Alder Annual yellow sweet clover Artichoke Aster Austrian fieldcress Bedstraw Beggartick Biden Bindweed Bird vetch Bitterweed Bitter wintercress Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Blue lettuce Blue vervain Box elder Bracted plantain	Broadleaf dock Broadleaf plantain Broomweed Buckhorn Buckhorn plantain Bulbous buttercup Bull nettle Bull thistle Burdock Burning nettle Bur ragweed Burweed Buttercup Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catsear Catnip Chickweeed Chicory	Cockle Cocklebur Coffeebean Coffeeweed Common chickweeed Common mullein Common sowthistle Corn chamomile Creeping Jenny Crimson clover, croton Cudweed Curly dock Curly indigo Dandelion Dead Nettle Dock Dogbane Dogfennel Elderberry English daisy Fall dandelion	False sunflower Fiddleneck Field bindweed Field pansy Flea bane (Daisy) Flixweed Florida betony Florida pusley Frenchweed Galinsoga Garlic mustard Goathead Goatsbeard Goldenrod Ground ivy Gumweed Hairy bittercress Hairy fleabane Hawkweed Healall Heartleaf drymary
Bracted plantain	Chicory	Fall dandelion	Heartleaf drymary
Brassbuttons Bristly oxtongue	Cinquefoil Clover	False dandelion False flax	Hedge bindweed Hedge mustard

(Continued)

WEEDS CONTROLLED (Continued)

Hemp Henbit Hoary cress Hoary plantain Hoary vervain Honeysuckle Hop clover Horsenettle Horsetail Indiana mallow Ironweed Jewelweed Jimsonweed Kochia Knawel Knotweed Lambsquarter Lespedeza Locoweed Lupine Mallow

Lupine
Mallow
Marshelder
Matchweed
Mexicanweed
Milk vetch
Milkweed bloodflower

Mugwort
Morningglory
Mouseear chickweed
Musk thistle
Mustard

Narrowleaf plantain Narrowleaf vetch

Nettle Nutgrass Orange hawkweed

Oxalis
Oxeye daisy
Parsley-piert

Parsnip
Pearlwort
Pennycress
Pennywort
Peppergrass
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coeopsis
Plantain
Poison hemlock

Poison nemico Poison ivy Poison oak Pokeweed Poorjoe Povertyweed Prickly lettuce Prickly sida Primrose

Prostrate knotweeed
Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncture vine
Purslane
Ragweed
Red clover
Redroot pigweed

Red sorrel

Redstem filaree Rough cinquefoil Rough fleabane Roundleafed marigold Rush

Russian pigweed Russian thistle St. Johnswort Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smartweed Smooth dock

Smooth pigweed Sneezeweed southern wild rose

Sowthistle
Spanishneedle
Spatterdock
Speedwell
Spiny amaranth
Spiny cocklebur
Spotted catsear
Spotted knapweed
Spotted spurge
Spurge
Spurweed
Stinging nettle

Stinkweed Stichwort Strawberry clover Sumac Sunflower Sweet clover Tall nettle Tall vervain Tansy mustard Tansy ragwort

Tansy ragwort
Tanweed
Tarweed
Thistle
Tick trefoil
Toadflax
Trailing crown vetch

Tumble mustard
Tumble pigweed
Tumbleweed

Velvet leaf Venice mallow Veronica Vervain Vetch

Virginia buttonweeed Virginia creeper Virginia pepperweed Wavyleaf bullthistle Western clematis Western salsify White clover White mustard Wild mustard Wild aster Wild buckwheat Wild carrot Wild four-o'clock, Wild geranium Wild lettuce Wild marigold Wild parsnip

Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Willow

Witchweed Woolly morningglory Woodsorrel Woolly croton Woolly plantain

Woolly plantain Wormseed Yarrow Yellow Rocket

Yellow flower pepperweed and other broadleaf weeds

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to southern grasses - under conditions of inadequate moisture, high temperature and high humidity, overapplication may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 18 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 3.6 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and uneveness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.9 pound per 250 square feet.

SPREADER SETTINGS ONE PASS APPLICATION:

MAKE	<u>SETTING</u>
Earthway	5
Scotts	3-3/4
Cyclone	5
Republic	3
Sears	5
Red Devil (Precision)	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

(The following paragraph is mandatory in Kentucky when the nitrogen analysis does not provide 1 pound nitrogen per 1,000 square feet; otherwise, this information is optional.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂0 levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on inert fertilizer claimed.

Environmental Benefits

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

(Grasscycling Logo)

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- √ A 50' x 50' lawn produces enough oygen for a family of 4 every day
- √ Reduces Glare & Reduces Noise
- √ Traps Dust, Smoke Particles & Other Pollutants
- √ Purifies Water & Modifies Temperature
- √ Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides and keep any bags containing unused porduct in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides. Do not stack more than two pallets high.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(111207A)

Triplet and The Optical Advantage are registered trademarks of Nufarm Americas Inc.

OPTIONAL SPREADER INSTRUCTIONS:

(Illustration) For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn,

overlapping very slightly and shutting off spreader when turning.

(Illustration) For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the

center, overlapping very slightly.

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer

manufacturers. Chlorine declaration and fertilizer guarantee are optional except where mandated by State

fertilizer regulation.

OPTIONAL MARKETING HYPE - FRONT PANEL

Patented Low Odor Formulation

Low Odor Formulation

Now With Patented Low Odor Formulation

Kills Over 250 Weed Types

Kills 250+ Weed Types

Kills Over 250 Weed Listed

Kills Over 250 Weeds

Kills All Major Broadleaf Weeds

Kills Dandelion, Clover and Other Broadleaf Weeds

Kills Dandelion, Clover and Other Listed Broadleaf Weeds

Controls Over 250 Weeds

Use When Weeds Are Actively Growing

May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass

Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More

Overnight Results

Works in 24 Hours

Kills the Weeds, Not the Lawn

Rainproof in Hours - Won't Wash Away

Won't Harm the Home Lawn

Illustration(s) of various weeds listed on label

Contains Triplet Low Odor Formula

Easy to Use

For Thick, Weed-Free Turf

Three Way Control

Apply to a Moist Lawn

Contains The Three Way Power of Triple Low Odor

18 lb. Bag Treats Up To 5,000 Square Feet (XXX bag covers up to XX,XXX square feet.)

New Formulation

Specially formulated for residential use.