



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

SEP - 5 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Nufarm Americas Inc.
150 Harvester Drive
Suite 200
Burr Ridge, IL 60527

Subject: EPA Reg. 228-508 (Triplet Low Odor Premium 8000 Lawn Weed Killer Concentrate)/
2,4-D RED Label Amendment

The labeling referred to above is in compliance with the 2,4-D RED provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- 2) Please note: Final product reregistration can't be considered until after all active ingredients in this product are eligible for reregistration.

Commercial Use Label

- 1) To the label add "Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes".
- 2) The Hazards to Humans and Domestic Animals must be revised to read: "Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing."
- 3) The PPE section must be revised to read:

"All mixers, loaders, applicators, and other handlers must wear:

Long-sleeved shirt and long pants,
Shoes and socks,

Chemical-resistant gloves, except for applicators using groundboom equipment, and
Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise
exposed to the concentrate."

- 4) The mechanical transfer engineering control text is no longer needed and may be deleted from the label.
- 5) The first two sentences of the Environmental Hazard section must be revised to read: "This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below

the mean high water mark.”

6) Spray drift text must be added to the label and should read:

“Spray Drift Management

A variety of factors including weather conditions (e.g., wind directions, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Use a medium or coarser spray nozzles according to (ASAE standard 572) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.”

7) The text "Do not enter or allow people (or pets) to enter the treated area until dusts have settled" currently appearing on the label must be corrected to read "Do not enter or allow entry until sprays have dried."

8) The following text must be added to the label:

"Minimum of 30 days between applications. Do not make more than 2 applications per year including broadcast applications in combination with spot treatments"

Residential Use Label:

1) The first three sentences of the Hazards to Humans and Domestic Animals must be revised to read: "Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing."

2) The first paragraph of the Environmental Hazard section must be revised to read: "This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate."

3) The text "Use a spray pattern with coarse spray droplets without any fine mist" currently on the label must be revised to read "Do not apply as a fine mist because of potential for injury to desirable plants."

4) The text below must be added to the label: "Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated areas(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only."

5) The following rate restrictions must be added to the label: "The maximum application rate is 2.5 fluid ounces per 250 square feet. Do not apply more than 2 applications per year, including broadcast applications in combination with spot treatments. There must be a minimum of 30 days between applications."


Submit one copy of the revised final printed label for the record. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A

4/19

Page 4
EPA Reg. 228-508

stamped copy of the label is enclosed for your records. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,



Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

5/19

RESIDENTIAL

TRIPLET[®] LOW ODOR PREMIUM 8000 LAWN WEED KILLER Concentrate

FOR SELECTIVE BROADLEAF WEED CONTROL IN RESIDENTIAL TURF.
WHEN USING A HOSE-END SPRAYER, JUST CONNECT HOSE, TURN ON WATER AND SPRAY.

ONE QUART COVERS UP TO 8,000 SQUARE FEET

(SEE DIRECTIONS FOR DIFFERENT GRASSES.)

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid*	10.41%
Dimethylamine Salt of Mecoprop-p,**‡	1.79%
Dicamba Acid***	0.50%

OTHER INGREDIENTS:	87.30%
TOTAL:	100.00%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	5.58%, 0.481 lbs./gal.
**Mecoprop-p Acid	1.48%, 0.128 lbs./gal.
***Dicamba Acid	0.50%, 0.043 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

SEP - 5 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



EPA REG. NO. 228-508
EPA EST. NO. 228-IL-1

NET CONTENTS GALS.

228-508

000228-00508.20071109.EPA24D.Pending
NUP6K04

6/19

RESIDENTIAL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Harmful if inhaled. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse. Remove saturated clothing as soon as possible and shower. Wear long pants, long-sleeved shirt, socks, shoes, protective eyewear and chemical-resistant gloves when handling or applying this product.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.** Shake well before using.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. If necessary, contact your local Extension Agent.

RESIDENTIAL

WEEDS CONTROLLED

- Alder
- Annual yellow sweet clover
- Artichoke
- Aster
- Austrian fieldcress
- Bedstraw
- Beggartick
- Biden
- Bindweed
- Bird vetch
- Bitterweed
- Bitter wintercress
- Black-eyed Susan
- Black medic
- Black mustard
- Blackseed plantain
- Blessed thistle
- Blue lettuce
- Blue vervain
- Box elder
- Bracted plantain
- Brassbuttons
- Bristly oxtongue
- Broadleaf dock
- Broadleaf plantain
- Broomweed
- Buckhorn
- Buckhorn plantain
- Bulbous buttercup
- Bull nettle
- Bull thistle
- Burdock
- Burning nettle
- Bur ragweed
- Burweed
- Buttercup
- Canada thistle
- Carolina geranium
- Carpetweed
- Catchweed bedstraw
- Catsear
- Catnip
- Chickweed
- Chicory
- Cinquefoil
- Clover
- Cockle
- Cocklebur
- Coffeebean
- Coffeeweed
- Common chickweed
- Common mullein
- Common sowthistle
- Corn chamomile
- Creeping Jenny
- Crimson clover, croton
- Cudweed
- Curly dock
- Curly indigo
- Dandelion
- Dead Nettle
- Dock
- Dogbane
- Dogfennel
- Elderberry
- English daisy
- Fall dandelion
- False dandelion
- False flax
- False sunflower
- Fiddleneck
- Field bindweed
- Field pansy
- Flea bane (Daisy)
- Flixweed
- Florida betony
- Florida pusley
- Frenchweed
- Galinsoga
- Garlic mustard
- Goathead
- Goatsbeard
- Goldenrod
- Ground ivy
- Gumweed
- Hairy bittercress
- Hairy fleabane
- Hawkweed
- Healall
- Heartleaf drymary
- Hedge bindweed
- Hedge mustard
- Hemp
- Henbit
- Hoary cress
- Hoary plantain
- Hoary vervain
- Honeysuckle
- Hop clover
- Horsenettle
- Horsetail
- Indiana mallow
- Ironweed
- Jewelweed
- Jimsonweed
- Kochia
- Knawel
- Knotweed
- Lambsquarter
- Lespedeza
- Locoweed
- Lupine
- Mallow
- Marshelder
- Matchweed
- Mexicanweed
- Milk vetch
- Milkweed bloodflower
- Mugwort
- Morningglory
- Mouseear chickweed
- Musk thistle
- Mustard
- Narrowleaf plantain
- Narrowleaf vetch
- Nettle
- Nutgrass
- Orange hawkweed
- Oxalis
- Oxeye daisy
- Parsley-piert
- Parsnip
- Pearlwort
- Pennycress
- Pennywort
- Peppergrass
- Pepperweed
- Pigweed
- Pineywoods bedstraw
- Plains coeopsis
- Plantain
- Poison hemlock
- Poison ivy
- Poison oak
- Pokeweed
- Poorjoe
- Povertyweed
- Prickly lettuce
- Prickly sida
- Primrose
- Prostrate knotweed
- Prostrate pigweed
- Prostrate spurge
- Prostrate vervain
- Puncture vine
- Purslane
- Ragweed
- Red clover
- Redroot pigweed
- Red sorrel
- Redstem filaree
- Rough cinquefoil
- Rough fleabane
- Roundleafed marigold
- Rush
- Russian pigweed
- Russian thistle
- St. Johnswort
- Scarlet pimpernel
- Scotch thistle
- Sheep sorrel
- Shepherdspurse
- Slender plantain
- Smallflower galinsoga
- Smartweed
- Smooth dock
- Smooth pigweed
- Sneezeweed southern wild rose
- Sowthistle
- Spanishneedle
- Spatterdock
- Speedwell
- Spiny amaranth
- Spiny cocklebur
- Spotted catsear
- Spotted knapweed
- Spotted spurge
- Spurge
- Spurweed
- Stinging nettle
- Stinkweed
- Stichwort
- Strawberry clover
- Sumac
- Sunflower
- Sweet clover
- Tall nettle
- Tall vervain
- Tansy mustard
- Tansy ragwort
- Tanweed
- Tarweed
- Thistle
- Tick trefoil
- Toadflax
- Trailing crown vetch
- Tumble mustard
- Tumble pigweed
- Tumbleweed
- Velvet leaf
- Venice mallow
- Veronica
- Vervain
- Vetch
- Virginia buttonweed
- Virginia creeper
- Virginia pepperweed
- Wavyleaf bullthistle
- Western clematis
- Western salsify
- White clover
- White mustard
- Wild mustard
- Wild aster
- Wild buckwheat
- Wild carrot
- Wild four-o'clock,
- Wild geranium
- Wild lettuce
- Wild marigold
- Wild parsnip
- Wild radish
- Wild rape
- Wild strawberry
- Wild sweet potato
- Wild vetch
- Willow
- Witchweed
- Woolly morningglory
- Woodsorrel
- Woolly croton
- Woolly plantain
- Wormseed
- Yarrow
- Yellow Rocket
- Yellow flower pepperweed
- and other broadleaf weeds

USE TIPS

1. Spray during the growing season when weeds are growing actively.
2. For best results, do not mow for 2 days before or after treating.
3. Do not treat when air temperatures exceed 90°F or damage to grass may occur.
4. For best results, do not water lawn within 24 hours after treatment. Poor weed control may result if rainfall occurs within 24 hours of application.
5. Shake well before using.
6. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
7. Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees. Clover will be damaged if sprayed.
8. Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.
9. If certain tough weeds are not dead within 3 weeks, repeat treatment. Broadcast application is not necessary; follow spot treatment directions.

HOW MUCH TO USE

SPOT TREATMENT

Pressure (Trigger) Sprayer: Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahiagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

ENTIRE LAWN

Hose-End Sprayer: Use a hose-end sprayer designed for lawns. Pour recommended amount of product (Column 2 below) in bottle and add water to indicated level (Column 3), and spray on area shown in Column 4. For example on cool season grasses, to spray 2,500 square feet measure 20 fluid ounces (40 Tablespoons) in jar, add water to the 10 gallon level and mix. Turn on garden hose and spray as evenly as possible over 2,500 square feet.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahiagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

NOTE: Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

9/19

RESIDENTIAL

Pressure Sprayer: Measure indicated amount of this product, pour into tank, and dilute with 1 gallon of water according to the following chart. Use a spray pattern with coarse spray droplets without any fine mist.

AMOUNT OF CONCENTRATE IN 1 GALLON OF WATER REQUIRED TO TREAT 250 SQ. FT. OF LAWN	
GRASS TYPE	AMOUNT OF CONCENTRATE
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)
Warm Season Grasses Bahigrass Common Bermudagrass Zoysiagrass	1 Fluid Ounce (2 Tablespoon)
Warm Season Grasses Centipedegrass St. Augustinegrass*	1/2 Fluid Ounce (1 Tablespoon)

*Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep from freezing. to be stored in original container and placed in areas inaccessible to children.

PESTICIDE DISPOSAL: If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(110907A)

Triplet and The Optical Advantage is a registered trademark of Nufarm Americas Inc.

10/19

RESIDENTIAL

[OPTIONAL BOX]

PRODUCT FACTS

What it Does:
 Kills all Major Broadleaf Weeds including Chickweed, Clover and Dandelion (See inside for complete list)

Where to Use: (icon weeds)
 Lawns: Kills the Weeds not the Lawn

When to Use:
 Do not apply when daytime temperatures are above 90°F as lawn injury can occur.

Questions and Comments: (icon phone) (icon computer)
 Call xxxxxxxx or visit our website at www.xxxxxxxx

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING HYPE - FRONT PANEL

- Patented Low Odor Formulation
- Low Odor Formulation
- Now With Patented Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds
- Controls Over 250 Weeds
- Use When Weeds Are Actively Growing
- May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
- Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
- Overnight Results
- Works in 24 Hours
- Kills the Weeds, Not the Lawn
- Rainproof in Hours - Won't Wash Away
- Colorless Formulation
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Contains Triplet Low Odor Formula
- Easy to Use
- For Thick, Weed-Free Turf
- Three Way Control
- Apply to a Moist Lawn
- Contains The Three Way Power of Triple Low Odor
- One Quart Covers Up To 8,000 Square Feet (One XXX covers up to XX,XXX square feet.)
- Easy to Mix
- Just Add Water
- New Formulation
- Specially formulated for residential use.

11/19

RESIDENTIAL

ALTERNATE HOSE-END SPRAYER INSTRUCTIONS

How To Use:

(Illustrations)

1. Attach garden hose to spray nozzle.
2. Turn on water supply.
3. Bend safety tab down and turn control to "water"
4. Point nozzle toward lawn and turn control to "on". Product will automatically mix with water. Walk back and forth at a steady pace.

12/19

COMMERCIAL

TRIPLER[®] LOW ODOR PREMIUM 8000 LAWN WEED KILLER Concentrate

SELECTIVE BROADLEAF WEED CONTROL FOR ORNAMENTAL TURFGRASSES.
TO CONTROL DANDELION, CLOVER, HENBIT, PLANTAINS, WILD ONION, AND
MANY OTHER BROADLEAF WEEDS.

ALSO FOR HIGHWAYS, RIGHT-OF-WAYS AND OTHER SIMILAR NONCROP AREAS

CONTAINS 2,4-D, MECOPROP-P AND DICAMBA

1 QUART TREATS (COVERS) 8,000 SQ. FT.

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Triisopropanolamine Salt of 2,4 D Acid*	10.41%
Dimethylamine Salt Mecoprop-p Acid**†	1.79%
Dicamba Acid***	0.50%

OTHER INGREDIENTS:	87.30%
TOTAL:	100.00%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	5.58%, 0.481 lbs./gal.
**Mecoprop-p Acid	1.48%, 0.128 lbs./gal.
***Dicamba Acid	0.50%, 0.043 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with COMMENTS
In EPA Letter Dated:

SEP - 5 2008

EPA REG. NO. 228-508
EPA EST. NO. 228-IL-1

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. **150**

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS GALS.

228-508

000228-00508.20071109.EPA24D.Pending
NUP6K04

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks.
- chemical resistant gloves, when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry

Engineering Control Statements:

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

14/19

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperature over 90°F, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80°F. When air temperatures exceed 80°F, limit application to spot treatment only.

Make new dilution for each use. This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

- | | | | |
|----------------------------|------------------------|-------------------|----------------------|
| Alder | Carpetweed | Florida pusley | Matchweed |
| Annual yellow sweet clover | Catchweed bedstraw | Frenchweed | Mexicanweed |
| Artichoke | Catsear | Galinsoga | Milk vetch |
| Aster | Catnip | Garlic mustard | Milkweed bloodflower |
| Austrian fieldcress | Chickweed | Goathead | Mugwort |
| Bedstraw | Chicory | Goatsbeard | Morningglory |
| Beggartick | Cinquefoil | Goldenrod | Mouseear chickweed |
| Biden | Clover | Ground ivy | Musk thistle |
| Bindweed | Cockle | Gumweed | Mustard |
| Bird vetch | Cocklebur | Hairy bittercress | Narrowleaf plantain |
| Bitterweed | Coffeebean | Hairy fleabane | Narrowleaf vetch |
| Bitter wintercress | Coffeeweed | Hawkweed | Nettle |
| Black-eyed Susan | Common chickweed | Healall | Nutgrass |
| Black medic | Common mullein | Heartleaf drymary | Orange hawkweed |
| Black mustard | Common sowthistle | Hedge bindweed | Oxalis |
| Blackseed plantain | Corn chamomile | Hedge mustard | Oxeye daisy |
| Blessed thistle | Creeping Jenny | Hemp | Parsley-piert |
| Blue lettuce | Crimson clover, croton | Henbit | Parsnip |
| Blue vervain | Cudweed | Hoary cress | Pearlwort |
| Box elder | Curly dock | Hoary plantain | Pennycress |
| Bracted plantain | Curly indigo | Hoary vervain | Pennywort |
| Brassbuttons | Dandelion | Honeysuckle | Peppergrass |
| Bristly oxtongue | Dead Nettle | Hop clover | Pepperweed |
| Broadleaf dock | Dock | Horsenettle | Pigweed |
| Broadleaf plantain | Dogbane | Horsetail | Pineywoods bedstraw |
| Broomweed | Dogfennel | Indiana mallow | Plains coeopsis |
| Buckhorn | Elderberry | Ironweed | Plantain |
| Buckhorn plantain | English daisy | Jewelweed | Poison hemlock |
| Bulbous buttercup | Fall dandelion | Jimsonweed | Poison ivy |
| Bull nettle | False dandelion | Kochia | Poison oak |
| Bull thistle | False flax | Knawel | Pokeweed |
| Burdock | False sunflower | Knotweed | Poorjoe |
| Burning nettle | Fiddleneck | Lambsquarter | Povertyweed |
| Bur ragweed | Field bindweed | Lespedeza | Prairie |
| Burweed | Field pansy | Locoweed | Prickly lettuce |
| Buttercup | Flea bane (Daisy) | Lupine | Prickly sida |
| Canada thistle | Flixweed | Mallow | Primrose |
| Carolina geranium | Florida betony | Marshelder | Prostrate knotweed |

(Continued)

WEEDS CONTROLLED (Continued)

Prostrate pigweed	Smooth pigweed	Tanweed	Wild carrot
Prostrate spurge	Sneezeweed southern wild rose	Tarweed	Wild four-o'clock,
Prostrate vervain	Sowthistle	Thistle	Wild geranium
Puncture vine	Spanishneedle	Tick trefoil	Wild lettuce
Purslane	Spatterdock	Toadflax	Wild marigold
Ragweed	Speedwell	Trailing crown vetch	Wild parsnip
Red clover	Spiny	Tumble mustard	Wild radish
Redroot pigweed	Spiny amaranth	Tumble pigweed	Wild rape
Red sorrel	Spiny cocklebur	Tumbleweed	Wild strawberry
Redstem filaree	Spotted catsear	Velvet leaf	Wild sweet potato
Rough cinquefoil	Spotted knapweed	Venice mallow	Wild vetch
Rough fleabane	Spotted spurge	Veronica	Willow
Roundleafed marigold	Spurge	Vervain	Witchweed
Rush	Spurweed	Vetch	Woolly morningglory
Russian pigweed	Stinging nettle	Virginia buttonweed	Woodsorrel
Russian thistle	Stinkweed	Virginia creeper	Woolly croton
St. Johnswort	Stichwort	Virginia pepperweed	Woolly plantain
Scarlet pimpernel	Strawberry clover	Wavyleaf bullthistle	Wormseed
Scotch thistle	Sumac	Western clematis	Yarrow
Sheep sorrel	Sunflower	Western salsify	Yellow Rocket
Shepherdspurse	Sweet clover	White clover	Yellow flower pepperweed
Slender plantain	Tall nettle	White mustard	and other broadleaf weeds
Smallflower galinsoga	Tall vervain	Wild mustard	
Smartweed	Tansy mustard	Wild aster	
Smooth dock	Tansy ragwort	Wild buckwheat	

USE TIPS

1. Spray during the growing season when weeds are growing actively.
2. For best results, do not mow for 2 days before or after treating.
3. Avoid broadcast applications when air temperature exceeds 90 degrees.
4. For best results, do not water lawn within 24 hours after treatment. Poor weed control may result if rainfall occurs within 24 hours of application.
5. Shake well before using.
6. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
7. Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees. Clover will be damaged if sprayed.
8. Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.
9. If certain tough weeds are not dead within 3 weeks, repeat treatment. Broadcast application is not necessary; follow spot treatment directions.

16/19

HOW MUCH TO USE

SPOT TREATMENT

Pressure (Trigger) Sprayer: Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

ENTIRE LAWN

Hose-End Sprayer: Use a hose-end sprayer designed for lawns. Pour recommended amount of product (Column 2 below) in bottle and add water to indicated level (Column 3), and spray on area shown in Column 4. For example on cool season grasses, to spray 2,500 square feet measure 20 fluid ounces (40 Tablespoons) in jar, add water to the 10 gallon level and mix. Turn on garden hose and spray as evenly as possible over 2,500 square feet.

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

NOTE: Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

17/19

COMMERCIAL

Pressure Sprayer: Measure indicated amount of this product, pour into tank, and dilute with 1 gallon of water according to the following chart. Use a spray pattern with coarse spray droplets without any fine mist.

AMOUNT OF CONCENTRATE IN 1 GALLON OF WATER REQUIRED TO TREAT 250 SQ. FT. OF LAWN	
GRASS TYPE	AMOUNT OF CONCENTRATE
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)
Warm Season Grasses Bahigrass Common Bermudagrass Zoysiagrass	1 Fluid Ounce (2 Tablespoon)
Warm Season Grasses Centipedegrass St. Augustinegrass*	1/2 Fluid Ounce (1 Tablespoon)

*Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

Limitations: The maximum rate per application is 2.5 fluid ounces per 250 square feet, limited to 2 applications per year. The maximum seasonal rate is 5 fluid ounces per 250 square feet, excluding spot treatments.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Keep from freezing. To be stored in original container and placed in areas inaccessible to children.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Triplet rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(110807A)

Triplet and The Optical Advantage is a registered trademark of Nufarm Americas Inc.

18/19

OPTIONAL CONTAINER LANGUAGE:

Containers: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank, and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal.

Once clean, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer.

Returnable-Refillable Containers: Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

[OPTIONAL BOX]

PRODUCT FACTS	
What it Does: Kills all Major Broadleaf Weeds including Chickweed, Clover and Dandelion (See inside for complete list)	
Where to Use: (Icon weeds) Lawns: Kills the Weeds not the Lawn	
When to Use: Avoid broadcast applications when air temperature exceeds 90 degrees.	
Questions and Comments: (Icon phone) (Icon computer) Call xxxxxxxx or visit our website at www.xxxxxxxxxx	

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING HYPE - FRONT PANEL

- Patented Low Odor Formulation
- Low Odor Formulation
- Now With Patented Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds
- Controls Over 250 Weeds
- Use When Weeds Are Actively Growing
- May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass
- Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More
- Overnight Results
- Works in 24 Hours
- Kills the Weeds, Not the Lawn
- Rainproof in Hours - Won't Wash Away
- Colorless Formulation
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Contains Triplet Low Odor Formula
- Easy to Use
- For Thick, Weed-Free Turf
- Three Way Control

19/19

COMMERCIAL

Apply to a Moist Lawn
Contains The Three Way Power of Triple Low Odor
One Quart Covers Up To 8,000 Square Feet, (One XXX covers up to XX,XXX square feet)
Easy to Mix
Just Add Water
New Formulation

ALTERNATE HOSE-END SPRAYER INSTRUCTIONS

How To Use: (Illustrations)

1. Attach garden hose to spray nozzle.
2. Turn on water supply.
3. Bend safety tab down and turn control to "water".
4. Point nozzle toward lawn and turn control to "on". Product will automatically mix with water. Walk back and forth at a steady pace.