

228-508

6/13/2012

1/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr George Meindl
Nufarm Americas Inc
150 Harvester Dr , Suite 200
Burr Ridge, IL 60527

JUN 13 2012

Subject Notification per PR Notice 2007-4 and 2008-1
Triplet Low Odor Premium 8000 Lawn Weed Killer Concentrate
EPA Reg No 228-508
Application dated March 27, 2012

Dear Mr Meindl

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 and 2008-1 of the subject product The Registration Division (RD) has conducted a review of this request for its applicability under PRN 2007-4 and 2008-1 and finds that the action requested falls within the scope of 2007-4 and 2008-1 The label submitted with the application has been date-stamped "Notification" and will be placed in our records

If you have any questions, please contact Hope Johnson at 703-305-5410

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)



Please read instructions on reverse before completing form

Form Approved OMB No 2070 0060

<p>United States Environmental Protection Agency Washington DC 20460</p>	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
---	--	-----------------------

Application for Pesticide - Section I

1 Company/Product Number 228-508	2 EPA Product Manager Kathryn Montague	3 Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4 Company/Product (Name) Triplet Low Odor Premium 8000 Lawn Weed Killer Concentrate	PM# 23	
5 Name and Address of Applicant (Include ZIP Code) Nufarm Americas Inc 150 Harvester Drive Suite 200 Burr Ridge IL 60527 <input type="checkbox"/> Check if this is a new address	6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(i) my product is similar or identical in composition and labeling to <div style="text-align: center; font-weight: bold; font-size: 1.2em;">NOTIFICATION</div> EPA Reg No _____ JUN 13 2012 Product Name _____	

Section - II

<input type="checkbox"/> Amendment Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

Explanation Use additional page(s) if necessary (For section I and Section II)

Notification of label change per PR Notice 2007 4 and 2008 1 see cover letter for detailed explanation This notification is consistent with the guidance in PR Notice 2007 4 and 2008 1 and of EPA's regulations at 40 CFR 156 10 156 140 156 144 156 146 and 156 156 No other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make false statements to EPA I further understand that if the amended label is not consistent with the requirements of 40 CFR 156 10 156 140 156 144 156 146 and 156 156 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

Section - III

1 Material This Product Will Be Packaged In				2 Type of Container	
Child Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No			
* Certification must be submitted			If "Yes" Unit Packaging wgt _____ No per container _____	If "Yes" Package wgt _____ No per container _____	<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
3 Location of Net Contents Information		4 Size(s) Retail Container		5 Location of Label Directions	
<input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		1 pint 1 quart 1 2.5 and 5 gallons containers		<input checked="" type="checkbox"/>	
6 Manner in Which Label is Affixed to Product			<input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____		

Section - IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)		
Name George Meindl george.meindl@us.nufarm.com	Title Sr Regulatory Affairs Manager	Telephone No (Include Area Code) 630 455 2017
Certification I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		3 Date Application Received (Stamped)
2 Signature 	3 Title Sr Regulatory Affairs Manager	
4 Typed Name George Meindl	5 Date 3/27/2012	

3/19



Nufarm Americas, Inc
George Meindl
Sr Regulatory Affairs Manager
150 Harvester Drive Suite 200
Burr Ridge IL 60527
Phone 630 455 2017 Fax 630 455 2030
george.meindl@us.nufarm.com

March 27 2012

Via Overnight Courier

Kathryn Montague (PM-23)
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U S Environmental Protection Agency
Room S4900, One Potomac Yard
2777 S Crystal Drive
Arlington, VA 22202

NOTIFICATION

JUN 13 2012

Subject Triplet Low Odor Premium 8000 Lawn Weed Killer Concentrate
EPA Reg No 228-508
Label Notification per PR Notice 2007-4 and 2008-1

Dear Ms Montague

Nufarm Americas Inc would like to update the subject product s labeling to be in compliance with PR notice 2007-4 and 2008-1 We have adopted the exact wording set forth in the regulations which is also provided again in the PRN2007-4 and 2008-1 for both the residential label and commercial label versions of this registration

To process this request please find enclosed the following

- Application for Pesticide Registration (EPA form 8570-1)
- Revised labeling with areas of change clearly identified (1 copy)
- Revised labeling – clean (1 copy)
- Certification with Respect to Label Integrity
- CD containing the proposed labeling file name 000228-00508 20120327 PRN2007-4 and 2008-1 update (residential) pdf and file name 000228-00508 20120327 PRN2007-4 and 2008-1 update (commercial) pdf

If you should have any questions regarding this matter please feel free to contact me at 630 455 2017 or email at george.meindl@us.nufarm.com

Sincerely

George Meindl
Sr Regulatory Affairs Manager
Nufarm Americas, Inc

4/19/12

4/19

RESIDENTIAL

TRIPL[®] LOW ODOR PREMIUM 8000 LAWN WEED KILLER Concentrate

FOR SELECTIVE BROADLEAF WEED CONTROL IN RESIDENTIAL TURF
WHEN USING A HOSE END SPRAYER JUST CONNECT HOSE TURN ON WATER AND SPRAY

ONE QUART COVERS UP TO 8 000 SQUARE FEET
(SEE DIRECTIONS FOR DIFFERENT GRASSES)
CONTAINS 2 4 D MECOPROP p AND DICAMBA
GET THE OPTICAL ADVANTAGE[®]

NOTIFICATION

JUN 13 2012

ACTIVE INGREDIENTS

Trisopropanolamine Salt of 2 4 Dichlorophenoxyacetic Acid	10 41%
Dimethylamine Salt of Mecoprop p ‡	1 79%
Dicamba Acid	0 50%

OTHER INGREDIENTS

TOTAL

87 30%
100 00%

Isomer Specific Method Equivalent to

2 4 D Acid

5 58% 0 481 lbs /gal

Mecoprop p Acid

1 48% 0 128 lbs /gal

Dicamba Acid

0 50% 0 043 lbs /gal

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP p

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill Leak Fire or Exposure Call CHEMTREC (800) 424 9300
For Medical Emergencies Only Call (877) 325 1840

EPA REG NO 228 508
EPA EST NO 228 IL 1

MANUFACTURED BY
NUFARM AMERICAS INC
150 HARVESTER DRIVE
BURR RIDGE IL 60527



NET CONTENTS GALS

000228 00508 20120327 PRN2007-4 and 2008 † update
NUP6K04

5/19

RESIDENTIAL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed Causes moderate eye irritation Avoid contact with eyes or clothing Wash thoroughly with soap and water after handling and before eating drinking chewing gum or using tobacco Remove and wash contaminated clothing before reuse Remove saturated clothing as soon as possible and shower Wear long pants long sleeved shirt socks shoes protective eyewear and chemical resistant gloves when handling or applying this product

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 to 20 minutes • Call a poison control center or doctor for treatment advice
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible • Call a poison control center or doctor for further treatment advice
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 877 325 1840 for emergency medical treatment information	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non target plants To protect the environment do not allow pesticide to enter or run into storm drains drainage ditches gutters or surface waters Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems

This chemical has properties and characteristics associated with chemicals detected in groundwater The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination Application around a cistern or well may result in contamination of drinking water or groundwater

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling READ ENTIRE LABEL BEFORE USING THIS PRODUCT USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS Shake well before using

Do not apply this product in a way that will contact any person or pet either directly or through drift Keep people and pets out of the area during application Do not allow people or pets to enter the treated area until sprays have dried

USE PRECAUTIONS

IMPORTANT Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings Do not spray exposed roots of desirable plants Do not spray on Dichondra Lippia nor on grasses where desirable Clovers are present Do not apply to newly seeded grasses until they have been mowed 3 times Do not reseed for 3 to 4 weeks after use If lawn needs watering water thoroughly before application For best results do not water lawn within 24 hours after treatment Watering will wash off weed killing material Do not apply if rainfall is expected within 24 hours Avoid fine mists Use a lawn type sprayer with a coarse spray as wind drift is less likely Spray entire lawn wetting weeds and lawn grasses Do not exceed recommended rates because damage to turf may occur Spray when air is calm to avoid spray drift that might injure desirable ornamental plants Do not apply when temperature exceeds 90 F Make new dilution for each use This product can be mixed with some liquid fertilizers or liquid iron materials Because liquid fertilizer and liquid iron differ in pH free ammonia content density salt concentration and percentage of water a compatibility test is recommended prior to mixing in application equipment If necessary contact your local Extension Agent

6/19

RESIDENTIAL

WEEDS CONTROLLED

Alder	Dogbane	Narrowleaf vetch	Spotted spurge
Annual yellow sweet clover	Dogfennel	Nettle	Spurge
Artichoke	Elderberry	Nutgrass	Spurweed
Aster	English daisy	Orange hawkweed	Stinging nettle
Austrian fieldcress	Fall dandelion	Oxalis	Stinkweed
Bedstraw	False dandelion	Oxeye daisy	Stichwort
Beggartick	False flax	Parsley piert	Strawberry clover
Biden	False sunflower	Parsnip	Sumac
Bindweed	Fiddleneck	Pearlwort	Sunflower
Bird vetch	Field bindweed	Pennycress	Sweet clover
Bitterweed	Field pansy	Pennywort	Tall nettle
Bitter wintercress	Flea bane (Daisy)	Peppergrass	Tall vervain
Black eyed Susan	Flixweed	Pepperweed	Tansy mustard
Black medic	Florida betony	Pigweed	Tansy ragwort
Black mustard	Florida pusley	Pineywoods bedstraw	Tanweed
Blackseed plantain	Frenchweed	Plains coeopsis	Tarweed
Blessed thistle	Galinsoga	Plantain	Thistle
Blue lettuce	Garlic mustard	Poison hemlock	Tick trefoil
Blue vervain	Goathead	Poison ivy	Toadflax
Box elder	Goatsbeard	Poison oak	Trailing crown vetch
Bracted plantain	Goldenrod	Pokeweed	Tumble mustard
Brassbuttons	Ground ivy	Poorjoe	Tumble pigweed
Bristly oxtongue	Gumweed	Povertyweed	Tumbleweed
Broadleaf dock	Hairy bittercress	Prickly lettuce	Velvet leaf
Broadleaf plantain	Hairy fleabane	Prickly sida	Venice mallow
Broomweed	Hawkweed	Primrose	Veronica
Buckhorn	Healall	Prostrate knotweed	Vervain
Buckhorn plantain	Heartleaf drymary	Prostrate pigweed	Vetch
Bulbous buttercup	Hedge bindweed	Prostrate spurge	Virginia buttonweed
Bull nettle	Hedge mustard	Prostrate vervain	Virginia creeper
Bull thistle	Hemp	Puncture vine	Virginia pepperweed
Burdock	Henbit	Purslane	Wavyleaf bullthistle
Burning nettle	Hoary cress	Ragweed	Western clematis
Bur ragweed	Hoary plantain	Red clover	Western safsify
Burweed	Hoary vervain	Redroot pigweed	White clover
Buttercup	Honeysuckle	Red sorrel	White mustard
Canada thistle	Hop clover	Redstem filaree	Wild mustard
Carolina geranium	Horsenettle	Rough cinquefoil	Wild aster
Carpetweed	Horsetail	Rough fleabane	Wild buckwheat
Catchweed bedstraw	Indiana mallow	Roundleafed margold	Wild carrot
Catsear	Ironweed	Rush	Wild four o'clock
Catnip	Jewelweed	Russian pigweed	Wild geranium
Chickweed	Jimsonweed	Russian thistle	Wild lettuce
Chicory	Kochia	St Johnswort	Wild margold
Cinquefoil	Knawel	Scarlet pimpernel	Wild parsnip
Clover	Knotweed	Scotch thistle	Wild radish
Cockle	Lambsquarter	Sheep sorrel	Wild rape
Cocklebur	Lespedeza	Shepherdspurse	Wild strawberry
Coffeebean	Locoweed	Slender plantain	Wild sweet potato
Coffeeweed	Lupine	Smallflower galinsoga	Wild vetch
Common chickweed	Mallow	Smartweed	Willow
Common mullein	Marshelder	Smooth dock	Witchweed
Common sowthistle	Matchweed	Smooth pigweed	Woolly morningglory
Corn chamomile	Mexicanweed	Sneezweed southern wild rose	Woodsorrel
Creeping Jenny	Milk vetch	Sowthistle	Woolly croton
Crimson clover croton	Milkweed bloodflower	Spanishneedle	Woolly plantain
Cudweed	Mugwort	Spatterdock	Wormseed
Curly dock	Morningglory	Speedwell	Yarrow
Curly indigo	Mouseear chickweed	Spiny amaranth	Yellow Rocket
Dandelion	Musk thistle	Spiny cocklebur	Yellow flower pepperweed
Dead Nettle	Mustard	Spotted catsear	and other broadleaf weeds
Dock	Narrowleaf plantain	Spotted knapweed	

USE TIPS

- 1 Spray during the growing season when weeds are growing actively
- 2 For best results do not mow for 2 days before or after treating
- 3 Do not treat when air temperatures exceed 90 F or damage to grass may occur
- 4 For best results do not water lawn within 24 hours after treatment Poor weed control may result if rainfall occurs within 24 hours of application
- 5 Shake well before using
- 6 On new lawns wait until the grass has reached a height of 2 inches before making an overall spray treatment However a spot treatment may be made at any time with a small pressure sprayer Wait 3 weeks before seeding or planting stolons plugs or sprigs
- 7 Do not allow spray to get on ornamental plants or vegetables since injury may result Do not spray around roots of ornamentals and trees Clover will be damaged if sprayed
- 8 Whether using a hose end sprayer or pressure sprayer avoid sprays which are too fine because they can drift and damage other plants
- 9 If certain tough weeds are not dead within 3 weeks repeat treatment Broadcast application is not necessary follow spot treatment directions

HOW MUCH TO USE

SPOT TREATMENT

Pressure (Trigger) Sprayer Adjust spray nozzle to give coarse spray Aim at center of weed and spray to wet One application should be sufficient Effects begin to show after a few days and weeds gradually die A repeat application may be required in 3 to 4 weeks for hard to kill (or control) weeds Do not use a hose end sprayer for spot treatment

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

ENTIRE LAWN

Hose End Sprayer Use a hose end sprayer designed for lawns Pour recommended amount of product (Column 2 below) in bottle and add water to indicated level (Column 3) and spray on area shown in Column 4 For example on cool season grasses to spray 2 500 square feet measure 20 fluid ounces (40 Tablespoons) in jar add water to the 10 gallon level and mix Turn on garden hose and spray as evenly as possible over 2 500 square feet

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

NOTE Do not apply to Floratam variety of St Augustine grass in Florida Contact your local County Extension Agent if you are not sure as to what type of grass you have

8/19

RESIDENTIAL

Pressure Sprayer Measure indicated amount of this product pour into tank and dilute with 1 gallon of water according to the following chart Do not apply as a fine mist because of potential for injury to desirable plants

AMOUNT OF CONCENTRATE IN 1 GALLON OF WATER REQUIRED TO TREAT 250 SQ FT OF LAWN	
GRASS TYPE	AMOUNT OF CONCENTRATE
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)
Warm Season Grasses Bahagrass Common Bermudagrass Zoysiagrass	1 Fluid Ounce (2 Tablespoon)
Warm Season Grasses Centipedegrass St Augustinegrass	1/2 Fluid Ounce (1 Tablespoon)

Do not apply to Floratam variety of St Augustine grass in Florida Contact your local County Extension Agent if you are not sure as to what type of grass you have

The maximum application rate is 2.5 fluid ounces per 250 square feet Do not apply more than 2 applications per year including broadcast applications in combination with spot treatments There must be a minimum of 30 days between applications

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

PESTICIDE STORAGE Keep from freezing To be stored in original container and placed in areas inaccessible to children

PESTICIDE DISPOSAL AND CONTAINER DISPOSAL Nonrefillable container Do not reuse or refill this container

If empty Place in trash or offer for recycling if available **If partly filled** Call your local solid waste agency for disposal instructions Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully TO THE EXTENT CONSISTENT WITH APPLICABLE LAW (1) THE GOODS DELIVERED TO YOU ARE FURNISHED AS IS BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES GUARANTEES OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER EITHER EXPRESS OR IMPLIED OR BY USAGE OF TRADE STATUTORY OR OTHERWISE WITH REGARD TO THE PRODUCT SOLD INCLUDING BUT NOT LIMITED TO MERCHANTABILITY FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS OR THE MANNER OF USE OR APPLICATION INCLUDING WEATHER ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD INCLUDING USE APPLICATION HANDLING AND DISPOSAL MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER IF ANY OR FOR ANY DAMAGES OR SUMS OF MONEY CLAIMS OR DEMANDS WHATSOEVER RESULTING FROM OR BY REASON OF OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER USER OR ITS CUSTOMERS BUYER'S OR USER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT

If you do not agree with or do not accept any of directions for use the warranty disclaimers or limitations on liability do not use the product and return it unopened to the Seller and the purchase price will be refunded

(052009)

Triplet and The Optical Advantage is a registered trademark of Nufarm Americas Inc

5

9/19

RESIDENTIAL

[OPTIONAL BOX]

PRODUCT FACTS

What it Does
 Kills all Major Broadleaf Weeds including Chickweed Clover and Dandelion (See inside for complete list)

Where to Use (Icon weeds)
 Lawns Kills the Weeds not the Lawn

When to Use
 Do not apply when daytime temperatures are above 90 F as lawn injury can occur


Questions and Comments (Icon phone) (icon computer)
 Call xxxxxxxx or visit our website at www xxxxxxxxxxxx

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING HYPE FRONT PANEL

Patented Low Odor Formulation
 Low Odor Formulation
 Now With Patented Low Odor Formulation
 Kills Over 250 Weed Types
 Kills 250+ Weed Types
 Kills Over 250 Weed Listed
 Kills Over 250 Weeds
 Kills All Major Broadleaf Weeds
 Kills Dandelion Clover and Other Broadleaf Weeds
 Kills Dandelion Clover and Other Listed Broadleaf Weeds
 Controls Over 250 Weeds
 Use When Weeds Are Actively Growing
 May Be Used on Bahia Common Bermuda Bluegrass Centipede Fescue Ryegrass St Augustine and Zoysia Grass
 Controls Weeds Such As Dandelions Chickweed Henbit Oxalis Plantain And Many More
 Overnight Results
 Works in 24 Hours
 Kills the Weeds Not the Lawn
 Rainproof in Hours Wont Wash Away
 Colorless Formulation
 Wont Harm the Home Lawn
 Illustration(s) of various weeds listed on label
 Contains Triplet Low Odor Formula
 Easy to Use
 For Thick Weed Free Turf
 Three Way Control
 Apply to a Moist Lawn
 Contains The Three Way Power of Triple Low Odor
 One Quart Covers Up To 8 000 Square Feet (One XXX covers up to XX XXX square feet)
 Easy to Mix
 Just Add Water
 New Formulation
 Specially formulated for residential use



10/19


RESIDENTIAL

ALTERNATE HOSE END SPRAYER INSTRUCTIONS

How To Use

(Illustrations)

- 1 Attach garden hose to spray nozzle
- 2 Turn on water supply
- 3 Bend safety tab down and turn control to water
- 4 Point nozzle toward lawn and turn control to on Product will automatically mix with water Walk back and forth at a steady pace



45 11/19/12

11/19

COMMERCIAL

TRIPLET[®] LOW ODOR PREMIUM 8000 LAWN WEED KILLER Concentrate

NOTIFICATION

JUN 13 2012

SELECTIVE BROADLEAF WEED CONTROL FOR ORNAMENTAL TURFGRASSES
TO CONTROL DANDELION CLOVER HENBIT PLANTAINS WILD ONION AND
MANY OTHER BROADLEAF WEEDS

ALSO FOR HIGHWAYS RIGHT OF WAYS AND OTHER SIMILAR NONCROP AREAS

CONTAINS 2 4 D MECOPROP P AND DICAMBA

1 QUART TREATS (COVERS) 8,000 SQ FT

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS

Triisopropanolamine Salt of 2 4 D Acid	10 41%
Dimethylamine Salt Mecoprop p Acid ‡	1 79%
Dicamba Acid	0 50%

OTHER INGREDIENTS

TOTAL	<u>87 30%</u>
	100 00%

Isomer Specific Method Equivalent to

2 4 D Acid	5 58%	0 481 lbs /gal
Mecoprop p Acid	1 48%	0 128 lbs /gal
Dicamba Acid	0 50%	0 043 lbs /gal

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP p

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill Leak Fire or Exposure Call CHEMTREC (800) 424 9300
For Medical Emergencies Only Call (877) 325 1840

EPA REG NO 228 508
EPA EST NO 228 IL 1

MANUFACTURED BY
NUFARM AMERICAS INC
150 HARVESTER DRIVE
BURR RIDGE IL 60527



NET CONTENTS GALS

000228 00508 20120327 PRN2007 4 and 2008 01 update
NUP6K04

12/19

COMMERCIAL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed Causes moderate eye irritation Avoid contact with eyes or clothing

PERSONAL PROTECTIVE EQUIPMENT (PPE)

- All loaders applicators and other handlers must wear
- long sleeved shirt and long pants
 - shoes plus socks
 - chemical resistant gloves except for applicators using groundboom equipment and
 - chemical resistant apron when mixing or loading cleaning up spills or equipment or otherwise exposed to the concentrate

See engineering controls for additional requirements

Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS	
Users Should	
<ul style="list-style-type: none"> • Wash hands before eating drinking chewing gum using tobacco or using the toilet • Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing If pesticide gets on skin wash immediately with soap and water • Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 to 20 minutes • Call a poison control center or doctor for treatment advice
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible • Call a poison control center or doctor for further treatment advice

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 877 325 1840 for emergency medical treatment information

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non target plants To protect the environment do not allow pesticide to enter or run into storm drains drainage ditches gutters or surface waters Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems

This chemical has properties and characteristics associated with chemicals detected in groundwater The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination Application around a cistern or well may result in contamination of drinking water or groundwater

13/19

COMMERCIAL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow entry until sprays have dried.

Not for use on turf grown for sale or other commercial use as sod or for commercial seed protection or for research purposes.

USE PRECAUTIONS

IMPORTANT Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Avoid broadcast applications when air temperature exceeds 90 F. When using small spot treatment applications in temperature over 90 F, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80 F. When air temperatures exceed 80 F, limit application to spot treatment only.

Make new dilution for each use. This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature and relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Use a medium or coarser spray nozzle according to (ASAE standard 572) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed


Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on target deposition and there are not sensitive areas (including but not limited to residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use, or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.



Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2 4 D herbicides Where states have more stringent regulations they must be observed

Equipment


All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates

Do not apply with a nozzle height greater than 4 feet above the crop canopy

WEEDS CONTROLLED

Alder	Carpetweed	Florida pusley	Matchweed
Annual yellow sweet clover	Catchweed bedstraw	Frenchweed	Mexicanweed
Artichoke	Catsear	Galinsoga	Milk vetch
Aster	Catnip	Garlic mustard	Milkweed bloodflower
Austrian fieldcress	Chickweed	Goathead	Mugwort
Bedstraw	Chicory	Goatsbeard	Morningglory
Beggartick	Cinquefoil	Goldenrod	Mouseear chickweed
Biden	Clover	Ground ivy	Musk thistle
Bindweed	Cockle	Gumweed	Mustard
Bird vetch	Cocklebur	Hairy bittercress	Narrowleaf plantain
Bitterweed	Coffebean	Hairy fleabane	Narrowleaf vetch
Bitter wintercress	Coffeeweed	Hawkweed	Nettle
Black eyed Susan	Common chickweed	Healall	Nutgrass
Black medic	Common mullein	Heartleaf drymary	Orange hawkweed
Black mustard	Common sowthistle	Hedge bindweed	Oxalis
Blackseed plantain	Corn chamomile	Hedge mustard	Oxeye daisy
Blessed thistle	Creeping Jenny	Hemp	Parsley piert
Blue lettuce	Crimson clover croton	Henbit	Parsnip
Blue vervain	Cudweed	Hoary cress	Pearlwort
Box elder	Curly dock	Hoary plantain	Pennycress
Bracted plantain	Curly indigo	Hoary vervain	Pennywort
Brassbuttons	Dandelion	Honeysuckle	Peppergrass
Bristly oxtongue	Dead Nettle	Hop clover	Pepperweed
Broadleaf dock	Dock	Horsenettle	Pigweed
Broadleaf plantain	Dogbane	Horsetail	Pineywoods bedstraw
Broomweed	Dogfennel	Indiana mallow	Plains coeopsis
Buckhorn	Elderberry	Ironweed	Plantain
Buckhorn plantain	English daisy	Jewelweed	Poison hemlock
Bulbous buttercup	Fall dandelion	Jimsonweed	Poison ivy
Bull nettle	False dandelion	Kochia	Poison oak
Bull thistle	False flax	Knawel	Pokeweed
Burdock	False sunflower	Knotweed	Poorjoe
Burning nettle	Fiddleneck	Lambsquarter	Povertyweed
Bur ragweed	Field bindweed	Lespedeza	Prairie
Burweed	Field pansy	Locoweed	Prickly lettuce
Buttercup	Flea bane (Daisy)	Lupine	Prickly sida
Canada thistle	Flixweed	Mallow	Primrose
Carolina geranium	Florida betony	Marshelder	Prostrate knotweed

(Continued)



15/19


COMMERCIAL

WEEDS CONTROLLED (Continued)

Prostrate pigweed	Smooth pigweed	Tanweed	Wild carrot
Prostrate spurge	Sneezeweed southern wild rose	Tarweed	Wild four o'clock
Prostrate vervain	Sowthistle	Thistle	Wild geranium
Puncture vine	Spanishneedle	Tick trefoil	Wild lettuce
Purslane	Spatterdock	Toadflax	Wild marigold
Ragweed	Speedwell	Trailing crown vetch	Wild parsnip
Red clover	Spiny	Tumble mustard	Wild radish
Redroot pigweed	Spiny amaranth	Tumble pigweed	Wild rape
Red sorrel	Spiny cocklebur	Tumbleweed	Wild strawberry
Redstem filaree	Spotted catsear	Velvet leaf	Wild sweet potato
Rough cinquefoil	Spotted knapweed	Venice mallow	Wild vetch
Rough fleabane	Spotted spurge	Veronica	Willow
Roundleafed marigold	Spurge	Vervain	Witchweed
Rush	Spurweed	Vetch	Woolly morningglory
Russian pigweed	Stinging nettle	Virginia buttonweed	Woodsorrel
Russian thistle	Stinkweed	Virginia creeper	Woolly croton
St Johnswort	Stichwort	Virginia pepperweed	Woolly plantain
Scarlet pimpernel	Strawberry clover	Wavyleaf bullthistle	Wormseed
Scotch thistle	Sumac	Western clematis	Yarrow
Sheep sorrel	Sunflower	Western salsify	Yellow Rocket
Shepherdspurse	Sweet clover	White clover	Yellow flower pepperweed
Slender plantain	Tall nettle	White mustard	and other broadleaf weeds
Smallflower galinsoga	Tall vervain	Wild mustard	
Smartweed	Tansy mustard	Wild aster	
Smooth dock	Tansy ragwort	Wild buckwheat	

USE TIPS

- 1 Spray during the growing season when weeds are growing actively
- 2 For best results do not mow for 2 days before or after treating
- 3 Avoid broadcast applications when air temperature exceeds 90 degrees
- 4 For best results do not water lawn within 24 hours after treatment Poor weed control may result if rainfall occurs within 24 hours of application
- 5 Shake well before using
- 6 On new lawns wait until the grass has reached a height of 2 inches before making an overall spray treatment However a spot treatment may be made at any time with a small pressure sprayer Wait 3 weeks before seeding or planting stolons plugs or sprigs
- 7 Do not allow spray to get on ornamental plants or vegetables since injury may result Do not spray around roots of ornamentals and trees Clover will be damaged if sprayed
- 8 Whether using a hose end sprayer or pressure sprayer avoid sprays which are too fine because they can drift and damage other plants
- 9 If certain tough weeds are not dead within 3 weeks repeat treatment Broadcast application is not necessary follow spot treatment directions
- 10 Minimum of 30 days between applications Do not make more than 2 applications per year including broadcast applications in combination with spot treatments



16/19

COMMERCIAL

HOW MUCH TO USE

SPOT TREATMENT

Pressure (Trigger) Sprayer Adjust spray nozzle to give coarse spray Aim at center of weed and spray to wet One application should be sufficient Effects begin to show after a few days and weeds gradually die A repeat application may be required in 3 to 4 weeks for hard to kill (or control) weeds Do not use a hose end sprayer for spot treatment

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

ENTIRE LAWN

Hose End Sprayer Use a hose end sprayer designed for lawns Pour recommended amount of product (Column 2 below) in bottle and add water to indicated level (Column 3) and spray on area shown in Column 4 For example on cool season grasses to spray 2 500 square feet measure 20 fluid ounces (40 Tablespoons) in jar add water to the 10 gallon level and mix Turn on garden hose and spray as evenly as possible over 2 500 square feet

GRASS TYPE	AMOUNT OF CONCENTRATE	AMOUNT OF WATER	AREA TREATED
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)	1 Gallon	250 Square Feet
Warm Season Grasses Bahagrass Common Bermudagrass Centipedegrass St Augustinegrass Zoysiagrass	1 Fluid Ounce (2 Tablespoons)	1 Gallon	250 Square Feet

NOTE Do not apply to Floratam variety of St Augustine grass in Florida Contact your local County Extension Agent if you are not sure as to what type of grass you have



17/19

COMMERCIAL

Pressure Sprayer Measure indicated amount of this product pour into tank and dilute with 1 gallon of water according to the following chart Use a spray pattern with coarse spray droplets without any fine mist

AMOUNT OF CONCENTRATE IN 1 GALLON OF WATER REQUIRED TO TREAT 250 SQ. FT. OF LAWN	
GRASS TYPE	AMOUNT OF CONCENTRATE
Cool Season Grasses Bluegrass Fescue Ryegrass	2 Fluid Ounces (4 Tablespoons)
Warm Season Grasses Bahiagrass Common Bermudagrass Zoysiagrass	1 Fluid Ounce (2 Tablespoon)
Warm Season Grasses Centipedegrass St. Augustinegrass	1/2 Fluid Ounce (1 Tablespoon)

Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

Limitations The maximum rate per application is 2.5 fluid ounces per 250 square feet limited to 2 applications per year. The maximum seasonal rate is 5 fluid ounces per 250 square feet excluding spot treatments.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE Keep from freezing. To be stored in original container and placed in areas inaccessible to children.

PESTICIDE DISPOSAL Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL [HANDLING] Nonrefillable Containers 5 Gallons or Less Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable containers larger than 5 gallons Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning not available, puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

OR

Refillable containers larger than 5 gallons Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully TO THE EXTENT CONSISTENT WITH APPLICABLE LAW (1) THE GOODS DELIVERED TO YOU ARE FURNISHED AS IS BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER EITHER EXPRESS OR IMPLIED OR BY USAGE OF TRADE STATUTORY OR OTHERWISE WITH REGARD TO THE

18/19

PRODUCT SOLD INCLUDING BUT NOT LIMITED TO MERCHANTABILITY FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS OR THE MANNER OF USE OR APPLICATION INCLUDING WEATHER ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS


LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD INCLUDING USE APPLICATION HANDLING AND DISPOSAL MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER IF ANY OR FOR ANY DAMAGES OR SUMS OF MONEY CLAIMS OR DEMANDS WHATSOEVER RESULTING FROM OR BY REASON OF OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER USER OR ITS CUSTOMERS BUYER'S OR USER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT

If you do not agree with or do not accept any of directions for use the warranty disclaimers or limitations on liability do not use the product and return it unopened to the Seller and the purchase price will be refunded

(052009)

Triplet and The Optical Advantage is a registered trademark of Nufam Americas Inc



19/19

COMMERCIAL

[OPTIONAL BOX]

PRODUCT FACTS

What it Does
Kills all Major Broadleaf Weeds including Chickweed Clover and Dandelion (See inside for complete list)

Where to Use (Icon weeds)
Lawns Kills the Weeds not the Lawn

When to Use
Avoid broadcast applications when air temperature exceeds 90 degrees

Questions and Comments (Icon phone) (Icon computer)
Call xxxxxxxx or visit our website at www xxxxxxxxxx

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING TYPE FRONT PANEL

- Patented Low Odor Formulation
- Low Odor Formulation
- Now With Patented Low Odor Formulation
- Kills Over 250 Weed Types
- Kills 250+ Weed Types
- Kills Over 250 Weed Listed
- Kills Over 250 Weeds
- Kills All Major Broadleaf Weeds
- Kills Dandelion Clover and Other Broadleaf Weeds
- Kills Dandelion Clover and Other Listed Broadleaf Weeds
- Controls Over 250 Weeds
- Use When Weeds Are Actively Growing
- May Be Used on Bahia Common Bermuda Bluegrass Centipede Fescue Ryegrass St Augustine and Zoysia Grass
- Controls Weeds Such As Dandelions Chickweed Henbit Oxalis Plantain And Many More
- Overnight Results
- Works in 24 Hours
- Kills the Weeds Not the Lawn
- Rainproof in Hours Won't Wash Away
- Colorless Formulation
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Contains Triplet Low Odor Formula
- Easy to Use
- For Thick Weed Free Turf
- Three Way Control
- Apply to a Moist Lawn
- Contains The Three Way Power of Triple Low Odor
- One Quart Covers Up To 8 000 Square Feet (One XXX covers up to XX XXX square feet)
- Easy to Mix
- Just Add Water
- New Formulation

ALTERNATE HOSE END SPRAYER INSTRUCTIONS

How To Use

(Illustrations)

- 1 Attach garden hose to spray nozzle
- 2 Turn on water supply
- 3 Bend safety tab down and turn control to 'water'
- 4 Point nozzle toward lawn and turn control to on Product will automatically mix with water Walk back and forth at a steady pace

