



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

20 MAR 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Nufarm Americas, Inc.
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527

Subject: EPA Reg. 228-511 / Triplet Low Odor Premium Spot Lawn Weed Killer Ready-to-Spray / RED Label Amendment: 2,4-D

The labeling referred to above is in compliance with the 2,4-D RED and is amended under the RED provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- 2) Please note: Final product reregistration can't be considered until after all active ingredients in this product are eligible for reregistration.
- 3) Add "Wear long-sleeved shirt, long pants, and shoes and socks" to the Hazards to Humans and Domestic Animals statements.
- 4) Revise the first paragraph of the Environmental Hazards section to "This pesticide is toxic to fish and aquatic invertebrates and may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinseate."

Submit one copy of the revised final printed label for the record. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Joanne S. Miller".

Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2/7

TRIPL[®] LOW ODOR PREMIUM SPOT LAWN WEED KILLER READY-TO-SPRAY

ACCEPTED
with COMMENTS
in EPA Letter Dated

FOR BROADLEAF WEED CONTROL IN RESIDENTIAL TURF
(SEE DIRECTIONS FOR DIFFERENT GRASSES.)

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA

GET THE OPTICAL ADVANTAGE[®]

20 MAR 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-511

ACTIVE INGREDIENTS:

Triisopropanolamine salt of 2,4-D Acid* (CAS No. 32341-80-3)	0.946%
Dimethylamine Salt of Mecoprop-p,**‡ (CAS No. 66423-09-4)	0.163%
Dicamba Acid*** (CAS no. 1918-00-9)	0.045%

OTHER INGREDIENTS:	<u>98.846%</u>
TOTAL:	100.000%

Isomer Specific Method, Equivalent to:

*2,4-D Acid	0.502%, 0.042 lbs./gal.
**Mecoprop-p Acid	0.135%, 0.011 lbs./gal.
***Dicamba Acid	0.045%, 0.004 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-511
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS GALS.

000228-00511.20061205.
NUP6L04

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if inhaled. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse. Remove saturated clothing as soon as possible and shower.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

Limited to 2 applications per year.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. For best results, do not water lawn within 24 hours after treatment. Watering will wash off weed killing material. Do not apply if rainfall is expected within 24 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F.

WEEDS CONTROLLED

Alder
 Annual yellow sweet clover
 Artichoke
 Aster
 Austrian fieldcress
 Bedstraw
 Beggartick
 Bidens
 Bindweed
 Bird vetch
 Bitterweed
 Bitter wintercress
 Black-eyed Susan
 Black medic
 Black mustard
 Blackseed plantain
 Blessed thistle
 Blue lettuce
 Blue vervain
 Box elder
 Bracted plantain
 Brassbuttons
 Bristly oxtongue
 Broadleaf dock
 Broadleaf plantain
 Broomweed
 Buckhorn
 Buckhorn plantain
 Bulbous buttercup
 Bull nettle
 Bull thistle
 Burdock
 Burning nettle
 Bur ragweed
 Burweed
 Buttercup
 Canada thistle
 Carolina geranium
 Carpetweed
 Catchweed bedstraw
 Catsear
 Catnip
 Chickweed
 Chicory
 Cinquefoil
 Clover
 Cockle
 Cocklebur
 Coffeebean
 Coffeeweed
 Common chickweed
 Common mullein
 Common sowthistle
 Corn chamomile
 Creeping Jenny
 Crimson clover, croton
 Cudweed
 Curly dock
 Curly indigo
 Dandelion
 Dead Nettle
 Dock
 Dogbane

Dogfennel
 Elderberry
 English daisy
 Fall dandelion
 False dandelion
 False flax
 False sunflower
 Fiddleneck
 Field bindweed
 Field pansy
 Flea bane (Daisy)
 Flixweed
 Florida betony
 Florida pusley
 Frenchweed
 Galinsoga
 Garlic mustard
 Goathead
 Goatsbeard
 Goldenrod
 Ground ivy
 Gumweed
 Hairy bittercress
 Hairy fleabane
 Hawkweed
 Healall
 Heartleaf drymary
 Hedge bindweed
 Hedge mustard
 Hemp
 Henbit
 Hoary cress
 Hoary plantain
 Hoary vervain
 Honeysuckle
 Hop clover
 Horsenettle
 Horsetail
 Indiana mallow
 Ironweed
 Jewelweed
 Jimsonweed
 Kochia
 Knawel
 Knotweed
 Lambsquarter
 Lespedeza
 Locoweed
 Lupine
 Mallow
 Marshelder
 Matchweed
 Mexicanweed
 Milk vetch
 Milkweed bloodflower
 Mugwort
 Morningglory
 Mouseear chickweed
 Musk thistle
 Mustard
 Narrowleaf plantain
 Narrowleaf vetch
 Nettle

Nutgrass
 Orange hawkweed
 Oxalis
 Oxeye daisy
 Parsley-piert
 Parsnip
 Pearlwort
 Pennycress
 Pennywort
 Peppergrass
 Pepperweed
 Pigweed
 Pineywoods bedstraw
 Plains coeopsis
 Plantain
 Poison hemlock
 Poison ivy
 Poison oak
 Pokeweed
 Poorjoe
 Povertyweed
 Prickly lettuce
 Prickly sida
 Primrose
 Prostrate knotweed
 Prostrate pigweed
 Prostrate spurge
 Prostrate vervain
 Puncture vine
 Purslane
 Ragweed
 Red clover
 Redroot pigweed
 Red sorrel
 Redstem filaree
 Rough cinquefoil
 Rough fleabane
 Roundleafed marigold
 Rush
 Russian pigweed
 Russian thistle
 St. Johnswort
 Scarlet pimpernel
 Scotch thistle
 Sheep sorrel
 Shepherdspurse
 Slender plantain
 Smallflower galinsoga
 Smartweed
 Smooth dock
 Smooth pigweed
 Sneezeweed southern wild rose
 Sowthistle
 Spanishneedle
 Spatterdock
 Speedwell
 Spiny amaranth
 Spiny cocklebur
 Spotted catsear
 Spotted knapweed
 Spotted spurge
 Spurge
 Spurweed

Stinging nettle
 Stinkweed
 Stichwort
 Strawberry clover
 Sumac
 Sunflower
 Sweet clover
 Tall nettle
 Tall vervain
 Tansy mustard
 Tansy ragwort
 Tanweed
 Tarweed
 Thistle
 Tick trefoil
 Toadflax
 Trailing crown vetch
 Tumble mustard
 Tumble pigweed
 Tumbleweed
 Velvet leaf
 Venice mallow
 Veronica
 Vervain
 Vetch
 Virginia buttonweed
 Virginia creeper
 Virginia pepperweed
 Wavyleaf bullthistle
 Western clematis
 Western salsify
 White clover
 White mustard
 Wild mustard
 Wild aster
 Wild buckwheat
 Wild carrot
 Wild four-o'clock
 Wild geranium
 Wild lettuce
 Wild marigold
 Wild parsnip
 Wild radish
 Wild rape
 Wild strawberry
 Wild sweet potato
 Wild vetch
 Willow
 Witchweed
 Woolly morningglory
 Woodsorrel
 Woolly croton
 Woolly plantain
 Wormseed
 Yarrow
 Yellow Rocket
 Yellow flower pepperweed
 and other broadleaf weeds

USE TIPS

1. Spray during the growing season when weeds are growing actively.
2. For best results, do not mow for 2 days before or after treating.
3. Do not treat when air temperatures exceed 90°F or damage to grass may occur.
4. For best results, do not water lawn within 24 hours after treatment. Do not apply if rain is expected within 24 hours of application.
5. Shake well before using.
6. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Do not plant seed, stolons, plugs, or sprigs within 3 weeks of application.
7. Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not apply over roots of ornamentals and trees. Do not apply to clover or mixtures of grass and clover, unless you wish to kill the clover.
8. If certain weeds are not dead within 3 weeks, repeat treatment. Broadcast application is not necessary; follow spot treatment directions.

HOW MUCH TO USE

SPOT TREATMENT

Pressure (Trigger) Sprayer: Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. To obtain wider coverage raise container.

LAWN GRASSES

Cool Season

- Bluegrass
- Fescue
- Ryegrass

Warm Season

- Bahiagrass
- Centipedegrass
- Common Bermudagrass
- St. Augustinegrass
- Zoysiagrass

NOTE: Do not apply to Floratam variety of St. Augustine grass in Florida. Contact your local County Extension Agent if you are not sure as to what type of grass you have.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep from freezing. To be stored in original container and placed in areas inaccessible to children.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

6/7

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(110907A)

Triplet and The Optical Advantage is a registered trademark of Nufarm Americas Inc.

[OPTIONAL BOX]

PRODUCT FACTS

Kills Broadleaf Weeds including Chickweed, Clover and Dandelion (See inside for complete list)

Where to Use:

Lawns: Kills the Weeds in listed lawn grasses.

When to Use:

Do not apply when daytime temperatures are above 90°F as lawn injury can occur.

Questions and Comments: (Icon phone) (Icon computer)

Call xxxxxxxx or visit our website at www.xxxxxxxxxx

OPTIONAL MARKETING HYPE - FRONT PANEL

Low Odor Formulation

Kills 250+ Weed Types

Kills Over 250 Weeds

Kills Dandelion, Clover and Other Broadleaf Weeds

Kills Dandelion, Clover and Other Listed Broadleaf Weeds

Controls Over 250 Weeds

Use When Weeds Are Actively Growing

May Be Used on Bahia, Common Bermuda, Bluegrass, Centipede, Fescue, Ryegrass, St. Augustine and Zoysia Grass

Controls Weeds Such As Dandelions, Chickweed, Henbit, Oxalis, Plantain, And Many More

Overnight Results

Works in 24 Hours

Kills the Weeds, Not the Lawn

Rainproof in Hours - Won't Wash Away

Colorless Formulation

Won't Harm the Home Lawn

Illustration(s) of various weeds listed on label

Contains Triplet Low Odor Formula

Easy to Use

For Thick, Weed-Free Turf

Three Way Control

Apply to a Moist Lawn

Contains The Three Way Power of Triplet Low Odor