



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

November 13, 2025

Joseph Wolfe, M.S.  
Regulatory Manager, T&O  
Nufarm Americas Inc.  
4020 Aerial Center Parkway  
Morrisville, NC 27560

Subject: Label Amendment - Registration Review Mitigation for Prodiamine  
Product Name: PRODEUCE RTU GRASS & WEED KILLER  
EPA Registration Number: 228-514  
Case Number: 474404  
Application Dates: June 20, 2022

Dear Joseph Wolfe:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Prodiamine Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at 202-566-0636, or via email at [carr.caleb@epa.gov](mailto:carr.caleb@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Julie R. Javier". The signature is fluid and cursive, with the first name "Julie" being the most prominent.

Julie Javier, Team Leader  
Risk Mitigation and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

PRODIAMINE	GROUP	3	HERBICIDE
GLYPHOSATE	GROUP	9	HERBICIDE

# ProDeuce® RTU

## GRASS & WEED KILLER

KILLS AND PREVENTS LISTED WEEDS AND GRASSES FOR UP TO 3 MONTHS  
ONE APPLICATION KILLS OR PREVENTS LISTED WEEDS FOR UP TO 3 MONTHS  
ON SOIL-SURFACED: PATIOS, DRIVEWAYS, SIDEWALKS AND MULCHED AREAS

### ACTIVE INGREDIENTS:

Glyphosate, N-(phosphonomethyl) glycine in the form of its isopropylamine salt\* ..... 2.00%  
Prodiamine\*\* ..... 0.50%

OTHER INGREDIENTS: ..... 97.50%  
TOTAL: ..... 100.00%

\* Contains 20.1 grams per litre or 0.17 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 14.8 grams per litre or 0.12 pounds per U.S. gallon of the acid, glyphosate.

\*\* Contains 5.0 grams per litre of Prodiamine (CAS # 29091-21-1).

### KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

**ACCEPTED**

Nov 13, 2025

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 228-514

EPA REG. NO. 228-514  
EPA EST. NO.

Manufactured For  
NUFARM AMERICAS INC.  
11901 S. Austin Ave.  
Alsip, IL 60803

NET CONTENTS:



000228-00514.20141104.MASTER

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION / PRECAUCION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>HOT LINE NUMBER</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

**ENVIRONMENTAL HAZARDS**

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash water or rinsate.

**PHYSICAL OR CHEMICAL HAZARDS**

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT, OR SPRAY SOLUTIONS OF THIS PRODUCT, IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Stay off treated areas until sprays have dried. Precautions should be taken to prevent tracking product after application on to desirable grasses.

**SHAKE WELL BEFORE USING.**

**WEED RESISTANCE MANAGEMENT**

For resistance management, please note that **ProDeuce**® RTU contains both a Group 3/prodiamine and a Group 9/glyphosate herbicide. Any weed population may contain plants naturally resistant to Group 3 and/or Group 9 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 or 9 herbicides within a season or among seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
  - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
  - (2) a spreading patch of non-controlled plants of a particular weed species;
  - (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, contact Nufarm at 1-800-345-3330.

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### Handheld Technology Applications

- Take precautions to minimize spray drift.

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## PRODUCT INFORMATION

### Where to Use

FOR OUTDOOR USE ONLY. Apply only where no vegetation is desired, such as driveways, landscaped areas, around well-established (at least 6 months old) plants, trees and shrubs, along fences, sidewalks, walkways, patios, curbs, gutters and cracks in sidewalks. DO NOT USE IN VEGETABLE GARDENS. DO NOT APPLY THROUGH ANY TYPE OF IRRIGATION SYSTEM. AERIAL APPLICATION IS PROHIBITED.

### When to Apply

Apply to actively growing weeds. Visible effects to most weeds should occur within 3 to 5 days with complete kill of most weeds in 2 to 4 weeks. If rainfall occurs within 6 hours, repeat treatment may be required. To avoid drift to desirable plants, apply only when air is calm. If leaves of desirable plants are accidentally sprayed, immediately wash off plants with water. Spray when air is calm to prevent drift to desirable plants. For best results, apply during warm, sunny weather (above 60°F).

### USE RESTRICTIONS

- **DO NOT** spray desirable plants.
- **DO NOT** use in lawns or for lawn renovation as this product prevents desirable grass establishment.
- **DO NOT** use for vegetable garden preparation or in and around fruits and vegetables.
- **DO NOT** spray around ornamental plants or in an area that will be planted or seeded 12 months after application if applied at the 1 gallon per 1,220 sq. ft. (1 quart per 305 sq. ft.) (1 pint per 155 sq. ft.) use rate.
- **DO NOT** spray around ornamental plants or in an area that will be planted or seeded 6 months when applying at the 1 gallon per 2,400 sq. ft. (1 quart per 600 sq. ft.) (1 pint per 300 sq. ft.) use rate.

## WEEDS CONTROLLED

### Pre-Emergent:

When used as directed, this product will prevent the following weeds from emerging:

Barnyardgrass	Henbit <sup>1</sup>	Purslane, common
Bluegrass, annual ( <i>Poa annual</i> ) <sup>1</sup>	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass <sup>2</sup>
Chickweed, common <sup>2</sup>	Junglerice	Shepherdspurse <sup>1</sup>
Chickweed, mouse-ear (from seed)	Knotweed <sup>1</sup>	Signalgrass, broadleaf
Crabgrass (large, smooth)	Kochia	Speedwell, Persian
Crowfootgrass	Lambsquarters, common	Sprangletop
Cupgrass, woolly	Lovegrass	Spurge, prostrate
Foxtails, annual	Panicum, (Texas, fall, browntop)	Witchgrass
Goosegrass	Pigweed	Woodsorrel, yellow (from seed)

<sup>1</sup>To control this weed in non-turf areas, apply this product in late summer, fall, or winter before weeds germinate.

<sup>2</sup>Suppression only.

### Post-Emergent Non-Selective:

#### ANNUAL WEEDS

Barley	Goatgrass, jointed	Rice, red
Barnyardgrass	Goosegrass	Rocket, London
Bluegrass, annual	Groundsel, common	Rye
Bluegrass, bulbous	Henbit	Ryegrass, Italian
Brome, downy	Horseweed / Mare's Tail	Shattercane
Buttercup	Lambsquarters, common	Shepherdspurse
Cheat	Mustard, blue	Sicklepod
Chickweed, common	Mustard, tansy	Signalgrass, broadleaf
Chickweed, mouse-ear (from seed)	Mustard, tumble	Spanishneedles
Cocklebur	Mustard, wild	Sprangletop
Corn	Oats, wild	Spurge, annual
Crabgrass	Panicum, fall	Spurry, umbrella
Dwarf dandelion	Panicum, Texas	Stinkgrass
False flax, smallweed	Pennycress, field	Teaweed
Filaree	Pigweed, redroot	Wheat
Foxtail	Pigweed, smooth	Witchgrass
Foxtail, Carolina	Primrose, cutleaf evening	
Geranium, Carolina	Pusley, Florida	

#### PERENNIAL WEEDS

Alfalfa	Dock, curly	Orchardgrass
Alligatorweed*	Dogbane, hemp	Pampasgrass*
Anise (fennel)	Fescue, tall	Paragrass
Artichoke, Jerusalem	Fescues	Phragmites*
Bahiagrass	Guinea grass	Poison hemlock
Bentgrass	Horsenettle	Quackgrass
Bermudagrass	Horseradish	Redvine*
Bermudagrass, water (knotgrass)	Ice plant	Reed, giant
Bindweed, field	Johnsongrass	Ryegrass, perennial
Bluegrass, Kentucky	Johnsongrass, seedling	Smartweed, swamp
Bluegrass, Texas	Kikuyugrass	Spurge, leafy*
Brackenfern	Knapweed	Starthistle, yellow
Bromegrass, smooth	Lantana	Sweet potato, wild*
Bursage, woollyleaf	Lespedeza	Thistle, artichoke
Canarygrass, reed	Milkweed	Thistle, Canada
Cattail	Morningglory	Timothy
Clover, red	Muhly, wirestem	Torpedograss*
Clover, white	Mullein, common	Trumpetcreeper*
Cogongrass	Napiergrass	Vaseygrass
Dallisgrass	Nightshade, silverleaf	Velvetgrass
Dandelion	Nutsedge; purple, yellow	Wheatgrass, Western

\* Partial Control

## How to Apply/Use

1. Turn sprayer nozzle to open. Adjust sprayer nozzle as desired.
2. Spray to thoroughly cover unwanted weeds and grasses or spray to thoroughly cover areas where you want to prevent weed and grass growth.
3. Do not spray ornamental plants or grasses, as damage will occur.
4. When treating around desirable plants, shield plants from drift with a sheet of cardboard or plastic. If desirable plants are accidentally sprayed, rinse off immediately with water.

### TANK SPRAYER (plastic, aluminum, fiberglass, plastic-lined steel or stainless steel)

- For longer pre-emergent control and post-emergent control of larger weeds use the higher rate.
- Apply 1 gallon of this product to treat 1,220 sq. ft. to 2,400 sq. ft. (1 quart of this product to treat 305 sq. ft. to 600 sq. ft.) (1 pint of this product to treat 155 sq. ft. to 300 sq. ft.)
- Do not apply with a galvanized or unlined steel (except stainless steel) sprayer, or through any irrigation system.

### HOW TO CLEAN SPRAYER

To clean sprayer after use, rinse sprayer and all sprayer parts with soap and water 3 times. Spray rinse water on bare soil or gravel. After cleaning, sprayer may be used for other products.

### WHEN TO REPLANT

All ornamental bedding plants, trees, shrubs, sod, and seed (flowers and grasses) may be planted 12 months after application when applied at the 1 gallon per 1,220 sq. ft. (1 quart per 305 sq. ft.) (1 pint per 155 sq. ft.) use rate or 6 months when applied at the 1 gallon per 2,400 sq. ft. (1 quart per 600 sq. ft.) (1 pint per 300 sq. ft.) use rate. For longer pre-emergent control and post-emergent control of larger weeds use the higher rate.

## STORAGE AND DISPOSAL

### [Residential]

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Keep from freezing. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

## STORAGE AND DISPOSAL

### [Commercial]

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Keep from freezing. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

#### **PESTICIDE DISPOSAL AND CONTAINER DISPOSAL:**

**Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Nonrefillable containers larger than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable containers larger than 5 gallons:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

#### **WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

#### **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RVmmdyy)

ProDeuce is a registered trademark of Nufarm Americas Inc.



**OPTIONAL LANGUAGE:**

<b>Product Facts – 1 Gallon Treats Up To 2,400 Sq. Ft. ([1 Qt.] [32 Fl. Oz.] Treats Up To [600 Sq. Ft.])</b>	
Kills Weeds	Listed weeds & grasses
Prevents Weeds	For up to 3 months
Where to Use	<ul style="list-style-type: none"><li>• Driveways, sidewalks, patios, curbs and gutters</li><li>• Along fences</li><li>• In landscaped areas (except vegetables gardens)</li><li>• For outdoor use only</li></ul>

**OPTIONAL MARKETING LANGUAGE:**

Season Long Control\*

\*Season Long is 3 months

Dual Action

Dual Action Formula

Kills weeds and grasses down to the root

Kills to the root

Ready-To-Use Formula

Starts working in hours

Systemic [control] action

Attacks the root

Fast Acting

Kills tough weeds

Kills weeds and grasses around trees and shrubs

Spot treat driveways, sidewalks and patios

Kills most weeds and grasses

Use where no vegetation desired

Quick and easy to use

Single application kills most weeds

**OPTIONAL STATEMENT:**

The preemergent herbicide prodiamine in this product is yellow in color, which allows the user to see the area that has been sprayed. If sprayed on cement curbs, sidewalks, and driveways, it may temporarily stain if not washed off immediately after application. Caution should be used not to wash the product off target weeds or weed control may be adversely affected. Treated areas should be dry before entering to avoid staining onto non-treated surfaces.