

228-530

08/17/2007

117



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:
228-530

Date of Issuance:
AUG 17 2007

NOTICE OF PESTICIDE:
[X] Registration
[] Reregistration
(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:
NUP 06095

Name and Address of Registrant (include ZIP Code):

NuFarm Americas, Inc.
150 Harvester Dr., Suite 200
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A), provided that you:

- 1. Signal Word: Based on the acute toxicity profile for this product, the signal word should be WARNING - AVISO.
2. Precautionary Statements- Hazards to Humans and Domestic Animals section: Based on the acute toxicity profile of this product and current labeling requirements per the Picloram and Salts and 2,4-D and Salts Reregistration Eligibility Decisions (REDs), the hazards statement must read, "Warning - Aviso. Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves. Prolonged or frequent repeated skin contact may cause allergic reactions in some individuals."

Comments continued on page 2 of this Notice of Registration

Signature of Approving Official:
Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Handwritten signature of Joanne I. Miller

Date:
AUG 17 2007

3. Under the Personal Protective Equipment section, modify the statement beginning, "Applicators and other handlers..." to read, "All mixers, loaders, applicators, flaggers, and other handlers must wear:"
4. Under the Engineering Controls section, add the statement "Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)]."
5. Under Environmental Hazards section:
 - a. Add the statement, "This pesticide is toxic to fish and aquatic invertebrates," at the beginning of the section.
 - b. Add the word, "or rinsate," after the word, "washwaters."
 - c. Add the following sentence after the sentence beginning, "Picloram is known to leach...": "2,4-D has properties and characteristics associated with chemicals detected in groundwater."
 - d. Modify the sentence beginning, "Use of this chemical..." to read, "Use of these chemicals..."
6. Due to the implication on the label that grasses treated with this product may be cut for hay, it falls within the scope of the WPS, and an Agricultural Use Requirements box is needed. This box should be placed under the Directions for Use section, immediately above the Non-Agricultural Use Requirements box, and should contain the following statements:
 - a. Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.
 - b. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.
 - c. PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, is:
 - i. Coveralls,
 - ii. Chemical-resistant gloves made of any waterproof material,
 - iii. Shoes plus socks, and
 - iv. Protective eyewear
7. In the Non-Agricultural Use Requirements box, modify the "Entry Restrictions for Non-WPS Uses" statement to read, "For applications on rangeland and permanent grass pastures, do not enter or allow others to enter treated area until sprays have dried."
8. A Picloram Stewardship Program is required to maintain registration of all end use products containing picloram. The program requires that you include the following Picloram Stewardship Program language on your product's label:

- 3/17
- a. "Every 2 years starting January 1, 2008, the registrant will offer training to applicators which will cover application techniques and product stewardship particular to their use(s) of this product (NUP 06095., Registration No 228-530). Applicators of this product must be able to provide certification of such training on demand to the State, Tribal or Federal enforcement agent."
 9. On page 11 of the draft label, under the first line of the Specific Use Directions column, delete the name, "Remedy," as this is not consistent with the proposed name for the current product.
 10. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar product to submit data.
 11. Add the statement, "EPA Reg. No. 228-530."
 12. Submit one copy of the final printed label before the product is released for shipment.

A stamped copy of the label is enclosed for your records.

Enclosure

4/17

NUP06095

RESTRICTED USE PESTICIDE

May Injure (Phytotoxic) Susceptible, Non-Target Plants. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

ACTIVE INGREDIENTS:

Picloram: 4-amino-3,5,6-trichloropicolinic acid, triisopropanolamine salt 10.2%
2,4-dichlorophenoxyacetic acid, triisopropanolamine salt 39.6%

OTHER INGREDIENTS: 50.2%
TOTAL: 100.0%

Acid equivalents:

Picloram: 4-amino-3,5,6-trichloropicolinic acid 5.7%, 0.54 lb/gal
2,4-dichlorophenoxyacetic acid 21.2%, 2 lb/gal

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with COMMENTS
In EPA Letter Dated:
AUG 17 2007

EPA REG. NO.
EPA EST. NO.

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-530

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527



NET CONTENTS:

5/17

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed or inhaled. Do not get in eyes or on clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, Neoprene Rubber, Polyvinyl Chloride (PVC), or Viton
- Shoes plus socks
- Protective eyewear
- **For containers of over 1 gallon, but less than 5 gallons:** Mixers and loaders who do not use a mechanical system (such as probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to other required PPE.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering Controls

For containers of 5 gallons or more: Do not open pour product from this container. A mechanical system (such as probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing immediately if pesticide gets inside. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to some plants at very low concentrations. Non-target plants may be adversely affected if pesticide is allowed to drift from areas of application. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below

6/17

the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes by cleaning of equipment or disposal of wastes. Do not allow run-off or spray to contaminate wells, irrigation ditches, or any body of water used for irrigation or domestic purposes. Do not make application when circumstances favor movement from treatment site.

Picloram is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, picloram may also have a high potential for runoff into surface water (primarily via dissolution in runoff water). These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Notice: Read the entire label. Use only according to label directions.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product through any We of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: For applications on rangeland and permanent grass pastures, do not allow worker entry into areas until sprays have dried, unless applicator and other handler PPE is worn.

GENERAL INFORMATION

This product is a water-soluble liquid product containing picloram and 2,4-D. Use this product in rangeland and permanent grass pastures to selectively control many annual, biennial, and perennial broadleaf weeds and woody species listed on this label.

Herbicidal effects of this product occur primarily from uptake by plant foliage and translocation throughout the plant, however, secondary herbicidal activity may occur from soil uptake of picloram. Very small amounts can kill or damage broadleaf plants. To prevent damage to crops and other desirable plants, carefully follow all directions and precautions.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

Observe any special use and application restrictions and limitations, including method of application and permissible areas of use as required by state or local regulations. When used in tank mix combination with other products, follow all applicable use directions, precautions, restrictions, and limitations on the labels of each product used.

Application Rate Ranges: Use higher rates in areas with dense weed populations or for longer residual control. For best results, the lower rate should be used only when environmental conditions are favorable for plant growth and when the plants are in the recommended growth stage. Compared to results obtained with the higher rate, a lower rate may be slower to show activity, provide a lower level of control, and may require retreatment.

Maximum Use Rates: Total use of this product must not exceed 4 quarts per acre per annual growing season. Repeat treatments may be applied as necessary, but total use must not exceed 4 quarts per acre per annual growing season.

Grazing Restrictions:

- There are no grazing restrictions for non-lactating dairy animals or other livestock including horses, sheep, goats, and other animals in the treatment area.
- Do not allow lactating dairy animals to graze treated areas within 7 days after application.
- Do not harvest grass cut for hay from treated areas for 30 days after application.
- Meat animals must be withdrawn from treated forage at least 3 days before slaughter.

This product should not be applied in residential areas or near ornamental trees and shrubs. Untreated trees can be affected by root uptake of the herbicide through movement into the top soil or by excretion of the product from the roots of nearby treated trees. Do not apply this product within the area occupied by roots of desirable trees, unless such injury can be tolerated.

7/17

On areas treated with this product, do not rotate to crops intended for food or feed use, other than range or pasture grasses, rye, forage sorghum, sudangrass, wheat, barley or oats not underseeded with a legume. **Do not move treated soil, or use treated soil for growing other plants** until soil residues of picloram are no longer detectable as indicated by an adequately sensitive bioassay or chemical test.

Do not spray pastures if the injury to existing forage legumes cannot be tolerated. This product may injure or kill legume plants. Forage legumes may be less sensitive to the herbicide after the seed has set and plant growth is mature. Seeding of legumes may not be successful if made within one year of application.

Established grasses are tolerant to this product, but newly seeded grasses may be injured until well established as indicated by tillering, development of a secondary root system and vigorous growth (see Planting Grasses Section).

This product may **suppress certain established grasses** such as smooth brome grass, Willman's lovegrass and buffalograss. However, subsequent grass growth should be improved by release from weed competition. Smooth brome grass and Willman's lovegrass grown for seed may be sensitive to this product if applied under adverse growing conditions (moisture stress).

Do not transfer livestock from treated grazing areas to broadleaf crop areas without first allowing 7 days of grazing on untreated grass pasture. Otherwise, urine may contain enough picloram to cause injury to sensitive broadleaf plants.

Do not use grass or hay from treated areas or manure from animals being fed treated forage or hay for composting or mulching of desirable, susceptible broadleaf plants.

Do not use manure from animals grazing treated areas on land used for growing broadleaf crops, ornamentals, orchards or other susceptible, desirable plants. Manure may contain enough picloram to cause injury to susceptible plants.

Do not mix with dry fertilizer.

Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

Do not use on sub-irrigated land.

Do not apply or otherwise permit this product or sprays containing this product to contact crops or other desirable broadleaf plants, including but not limited to alfalfa, beans, cotton, grapes, melons, peas, potatoes, safflower, soybeans, sugar beets, sunflower, tobacco, tomatoes, and other vegetable crops, flowers, fruit plants, ornamentals and shade trees.

Do not make application when circumstances favor movement from treatment site.

Avoid injurious spray drift. Applications should be made to avoid spray drift because very small quantities of the spray that may not be visible may severely injure susceptible crops during both growing and dormant periods. To minimize spray drift:

1. Use nozzle pressures no greater than are required to obtain a proper spray pattern for adequate coverage of target plants.
2. Apply as a coarse spray.
3. Use nozzles designed for herbicide application that do not produce a fine droplet spray.
4. Spray when wind velocity is low. Follow local state regulations. Avoid application under conditions which are conducive to air inversions or conditions of atmospheric temperature inversion.

When making applications near susceptible crops, spray drift may be further lessened by using a drift control system such as Microfoil, Thru-Valve boom (or equivalent) or a drift control agent such as Nalco-Trol (or equivalent). If a drift control additive is used, follow all use recommendations and precautions on the product label.

Ground Equipment: With ground equipment, spray drift may be lessened by keeping the spray boom as low as possible; by keeping the operating spray pressures at the manufacturers recommended minimum pressures for the specific nozzle types used (low pressure nozzles are available from spray equipment manufacturers). Do not apply this product with a mistblower. In hand-gun applications, spray drift may be minimized by selecting the minimum pressure that will provide adequate coverage (without forming a mist); by spraying no higher than brush tops.

Aerial Application: Avoid spray drift at the application site. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

1. The distance of the outer most operating nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory. [This information is advisory in nature and does not supersede mandatory label requirements.]

Aerial Drift Reduction Advisory

Information On Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produced larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

Wind: Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature And Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. The presence of inversions can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Determine Air Movement and Direction Before Making Foliar Applications: Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. It is suggested that a continuous smoke column at or near the spray site or a smoke generator on the spray equipment be used to detect air movements, lapse conditions, or temperature inversions (stable air). If the smoke layers or otherwise indicates a potential for hazardous spray drift, do not spray.

APPLICATION DIRECTIONS**Broadcast Foliar Application (Ground or Aerial)**

Unless otherwise specified, apply in water alone or in an oil-water emulsion in a total spray volume of 10 to 40 gallons per acre using ground equipment or 1 or more gallons per acre by aerial application. If aerially applied, results will be more consistent for spray volumes of 2 or more gallons per acre. Use of the lower total spray volume with ground equipment is recommended primarily where this product is applied simultaneously with liquid fertilizer. Good coverage is essential. For aerial application, swath width should not exceed 1-1/4 times the wingspan of the aircraft.

To provide more complete wetting and coverage of the foliage, a non-ionic surfactant may be used at recommended rates. The use of a drift control additive is recommended for drift reduction and improved deposition.

9/17

SECTION I: Control of Broadleaf Weeds and Woody Plants in Rangeland and Permanent Grass Pastures in the Southwest, Southeast, and Mid-Atlantic States

1-2 Pints/Acre or 3-4 Pints/Acre: Apply at the rate indicated by stage of growth to control the following woody plants or broadleaf weeds:	
Weed Species	Specific Use Directions
annual broomweed, bitter sneezeweed, bitterweed, buffalo bur, bull thistle, bursage (bur ragweed), camphor weed, cocklebur, common ragweed, croton, horseweed, lambsquarters, pigweed, prickly lettuce, smartweed, sunflower, tasajillo, wild carrot	<p>Early Season: Apply at a rate of 1-2 pints/acre in early to mid spring when weeds are less than 3 inches tall. Rates in the lower end of the rate range are effective only when weeds are less than 2 inches tall and conditions are favorable for plant growth.</p> <p>Mid to Late Season: Apply at a rate of 3-4 pints/acre in late spring to early summer when weeds are 3 inches tall to early flowering.</p>

2-4 Pints/Acre: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
aster, heath	Apply prior to bud stage when actively growing.
aster, spiny (Mexican devilweed)	Apply prior to bud stage when actively growing.
bee plant, Rocky Mountain	Apply prior to bud stage when actively growing.
bindweed, hedge	Apply prior to bud stage when actively growing.
blackberry	Tank-mix 2 pints per acre of this product with 1 pint per acre of Remedy* herbicide plus surfactant. Apply in late May to early June during or after bloom (not before) when the foliage is dark green. Do not treat blackberries in the same year after mowing, shredding, or burning. Even one year after removal of top growth, blackberry stands will be more difficult to control than undisturbed stands and will require retreatment.
buckwheat, climbing false	Apply prior to seed development when actively growing.
buckwheat, wild	Apply prior to seed development when actively growing.
bullnettle, western	Apply in spring when plants begin to flower.
bundleflower, Illinois	Apply prior to bud stage when actively growing.
burdock, common	Apply prior to bud stage when actively growing.
buttercup	Apply in early spring prior to bud stage.
chickweed, mouseear	Apply prior to bud stage when actively growing.
chicory	Apply from rosette stage to early bud stage when actively growing.
coneflower, upright prairie	Apply when plants are 6 inches tall, but before flowering.
common goldenweed, Drummond's goldenweed (<i>Isocoma spp.</i>)	Apply in spring (April-June) when favorable growing conditions result in substantial canopy development. Thorough and uniform coverage is essential. Use higher spray volumes (20-25 gpa for ground and 4-5 gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (see Mixing instructions).
curly dock	<p>Early Season: Apply 2 pints per acre prior to bolting stage of growth.</p> <p>Mid-to-Late Season: Apply at a rate of 3 to 4 pints/acre from bolting to bud stage.</p>

10/17

2-4 Pints/Acre: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds: <i>(continued)</i>	
Weed or Brush Species	Specific Use Directions
devil's-claw	Apply prior to flowering when actively growing.
dogfennel (cypressweed)	Apply when plants are from 6 to 24 inches tall, but before flowering. Increase rate within the rate range as season progresses and plants become larger.
erigonum, annual	Apply prior to bud stage when actively growing.
fleabane, rough	Apply prior to bud stage when actively growing.
gray goldaster narrowleaf goldaster	Apply in the spring during the bud stage (pre-bloom) using an oil-water emulsion spray. Thorough coverage is essential.
goldenrod, Missouri	Apply prior to bud stage when actively growing.
goldenweed, common, goldenweed, Drummond's (<i>Socoma spp.</i>)	Apply in spring (April-June) when favorable growing conditions result in substantial canopy development. Thorough and uniform coverage is essential. Use higher spray volumes (20-25 gpa for ground and 4-5 gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (see Mixing instructions).
hemlock, poison	Apply from rosette stage in spring or fall up to 36" tall.
hemlock, water (common)	Apply from rosette stage in spring or fall up to bud stage.
horsenettle, Carolina	Apply 2 pints per acre when plants are 4-6 inches tall. At 2 pints per acre retreatment may be necessary for acceptable control. Apply 3 to 4 pints per acre when flowering or for longer residual control of later emerging plants and greater stand reduction the following year.
horehound	Apply during active growth.
jimsonweed	Apply prior to bud stage when actively growing.
morningglory, ivyleaf	Apply prior to bud stage when actively growing.
mugwort	Apply prior to bud stage when actively growing.
nightshade, silverleaf	Apply 2 pints per acre when plants are 4-6 inches tall. Apply 3 to 4 pints per acre when flowering or for longer residual control of later emerging plants and greater stand reduction the following year. Retreatment is necessary for total control.
pennycress, field	Apply when plants are to 6 inches tall, but before flowering.
plantain, buckhorn	Apply prior to bud stage when actively growing.
pricklypoppy, annual	Apply prior to bud stage when actively growing.
puncturevine	Apply prior to bud stage when actively growing.
ragweed, common, giant, lanceleaf and western	Use lower rates in rate range when weeds are no more than 2 inches tall and conditions are favorable for plant growth. Use higher rates when weeds are from 3 inches tall to early flowering.
sagebrush, sand	Apply when new terminal growth reaches 6-12" and before average daytime temperature reaches 95°F. Use low rate only in early season.
snow-on-the-mountain	Apply prior to bud stage when actively growing.
sowthistle, spiny (prickly)	Apply prior to bud stage when actively growing.
stickweed	Apply 2 to 3 pints/acre prebloom.
thistles, biennial: including bull, musk, plumeless or scotch	Apply 2 pints/acre at rosette stage. Apply 3 to 4 pints/acre in mid to late season from bolting to bud stage.
vervain, blue vervain, hoary	Apply when plants are 6 inches tall to early flowering. Increase rate within the rate range as season progresses and weeds mature.

11/17

2-4 Pints/Acre: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds: <i>(continued)</i>	
Weed or Brush Species	Specific Use Directions
vetch, hairy	Apply prior to bud stage when actively growing.
wingstem	Apply 2-3 pints/acre prebloom.
yankeeweed	Apply when plants are 8 to 10 inches tall.

3-4 Pints/Acre: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
marshelder (sumpweed)	Apply in early season when weeds are less than 4 inches tall. Older plants require higher rates. <i>Thorough uniform coverage is essential. Use higher spray volumes (20-25 gpa for ground and 5 or more gpa for aerial equipment.</i>
mesquite and oak sprouts (suppression of regrowth):	Delay applications of this product for weed control until the foliage of regrowth brush in the treatment area is fully expanded and turned from light to dark green.
milkweed	Apply 4 pints/acre to actively growing milkweeds less than 4 inches tall. Add a surfactant at the manufacturer's recommended rate to improve wetting of foliage.
mullein, common	Apply 4 pints/acre during the rosette stage in spring or fall prior to bolting. Add a surfactant at the manufacturer's recommended rate to improve wetting of foliage.
poisonous plants such as: groundsel (Senecio spp.), garbancillo, (Wooton loco) and Woolly loco	Apply in fall or winter when moisture conditions are favorable. Because locoweeds are difficult to wet, use of a surfactant (0.25% to 0.5% v/v) or oil-water emulsion is recommended (see Mixing Instructions). Herbicide treatment may increase palatability of poisonous plants. Treated areas should not be grazed until the toxic plants are no longer palatable.
thistle, wavyleaf	Apply from rosette to late bolt stage.
tropical soda apple	Apply when plants are beginning to flower.

1 Gallon/Acre: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
cactus, pricklypear or cholla	Make ground broadcast application in the spring or early summer to control a broad spectrum of broadleaf weeds in addition to pricklypear.
Chinese tallowtree	Apply in spring or fall when conditions are favorable for plant growth. Thorough and uniform spray coverage is required. Use higher spray volumes (20-25 gpa for ground and 5 or more gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (see Mixing Instructions).
Macartney rose multiflora rose	Apply in spring or fall when conditions are favorable for plant growth. Thorough and uniform spray coverage is essential. Use higher spray volumes (20-25 gpa for ground and 5 or more gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (see Mixing Instructions). Avoid application within 9-12 months after mowing or when plants have a high percentage of new growth. Poor control will result if plants are less than 3 ft tall.
locust (honey and black) wild plum	Apply in spring when leaves are fully expanded and mature. Use of a surfactant (0.25 to 0.5% v/v) is recommended.

Section II: Control of Broadleaf Weeds and Woody Plants in Rangeland and Permanent Grass Pastures in the North and Northwestern U.S. including Colorado, Idaho, Iowa, Kansas, Minnesota, Missouri, Montana, Nebraska, North Dakota, Oregon, South Dakota, Utah, Washington and Wyoming

For best results in terms of forage response, desirable forage grasses should be present in the area to be treated in sufficient density to provide competition to lessen weed re-establishment following treatment. Additionally, good grazing management practices are recommended, particularly in the year following treatment, to allow forage grass density to increase.

12/17

Application Rates: Use higher rates in areas with dense weed populations or for longer residual control. For best results, the lower rate should be used only when environmental conditions are favorable for plant growth and when the plants are in the recommended growth stage. Compared to results obtained with the higher rate, a lower rate may be slower to show activity, provide a lower level of control, and may require retreatment.

2-4 Pints/Acre: Apply at the indicated stage of growth to control the following broadleaf plant species. Increase rate within rate range as growing season progresses:	
Weed or Brush Species	Specific Use Directions
absinth wormwood annual broomweed	Apply when actively growing in spring or early summer.
biennial thistles, such as bull, musk, plumeless or scotch	Apply 2 pt/acre at rosette stage. Apply 3 to 4 pt/acre to bolted thistle, but apply before early bud stage.
broom snakeweed	Apply after full leaf development to early bloom stage when plants are actively growing.
curly dock	Apply 2 pt/acre early season prior to bolting. Apply 3 to 4 pt/acre in mid to late season from bolting to early flower.
curlycup gumweed	Apply when new growth and seedlings have fully emerged before bloom stage.
fringed sagebrush	Apply a minimum of 3 pt/acre after seed stalk elongation and early flowering (mid - late June) and throughout the summer under good growing conditions.
goldenrod	Apply prior to bud stage during active growth.
hemp (marijuana) hemlock, poison	Apply from rosette stage in spring or fall up to 36" tall.
hemlock, water (common)	Apply from rosette stage in spring or fall up to bud stage.
ironweed, western	Apply 2 to 3 pt/acre prior to bud stage during active growth. A surfactant is recommended.
locoweeds, such as silky crazyweed (white point loco) and lambert crazyweed	Apply from early bud to early bloom stage. Herbicide application may increase palatability of these poisonous plants. Therefore, treated areas should not be grazed until after the toxic plants have dried up. Higher rate range should be considered to provide greater reduction of poisonous plants.
phlox, hoods	Apply during active growth.
plains pricklypear	Apply when the majority of plants are in the flower stage. The lower rate will provide a partial stand reduction. More complete control may be obtained with the higher rate. Treatment response is very slow and may continue for 2 years or longer.
ragweed, common, giant, fanceleaf and western	Use the lower rate in early season when weeds are no more than 2 inches tall. Use the higher rate when weeds range from 3 inches tall to early flowering, when conditions are favorable for plant growth.
thistles, biennial: including bull, musk, plumeless or scotch	Apply 2 pt/acre at rosette stage. Apply 3 to 4 pt/acre in mid to late season from bolting to bud stage.
vervain, blue and hoary	Apply when plants are 6 inches tall to early flowering. Increase rate within the rate range as season progresses and plants mature.
wormwood, Louisiana and absinth	Apply during active growth prior to woody stem development.
yarrow	Apply 2 pt/acre prior to bud stage. A surfactant is recommended.

13/17

4 Pints/Acre: Apply at the indicated stage of growth to control the following broadleaf weed species:	
Weed or Brush Species	Specific Use Directions
dense clubmoss	Apply in early summer with a surfactant at 0.25% v/v.
geyer larkspur	Apply from rosette to flower bud formation.
hairy goldenaster	Apply at bloom stage during active growth.
houndstongue	Apply to rosettes in late fall or early summer.
larkspur, plains	Apply prior to bud stage when actively growing.
licorice, wild	Apply at bloom stage, but before bur formation.
loco, woolly	Apply from bolting to early bloom. Herbicide application may temporarily increase palatability of this poisonous plant. Therefore, treated areas should not be grazed until toxic plants have dried up.
milkweed, common	Apply at bud stage when actively growing.
mullein, common	Apply during rosette stage in spring or fall prior to bolting. Add a surfactant at the manufacturer's recommended rate to improve wetting of foliage.
oxeye daisy	Apply 3-4 pt/acre when all plants have emerged to late flowering.
pussytoes	Apply prior to bud stage when actively growing. Use a surfactant at the manufacturer's recommended rate to improve wetting of foliage.

1 Gallon/Acre: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
Macartney rose multiflora rose	Apply in spring or fall when conditions are favorable for plant growth. Thorough and uniform spray coverage is essential. Use higher spray volumes (20-25 gpa for ground and 5 or more gpa for aerial equipment). Use of a non-ionic surfactant or oil-water emulsion is recommended (see Mixing Instructions). Avoid application within 9-12 months after mowing or when plants have a high percentage of new growth. Poor control will result if plants are less than 3 ft tall.
locust (honey and black) wild plum	Apply in spring when leaves are fully expanded and mature. Use of a surfactant (0.25 to 0.5% v/v) is recommended.

High-Volume Foliar Applications

Spray to thoroughly wet foliage and stems. The use of an approved agricultural surfactant is recommended. Do not use more than 1 gallon of this product (0.54 lb of picloram) per acre. To minimize spray drift, use lowest possible pressure and coarse spray to achieve good coverage. Keep sprays no higher than brush tops. Use of an approved drift control agent is recommended to reduce the potential for spray drift.

14/17

1 Gallon/100 Gallons of Spray: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
blackberry, elm, granjeno, locust, maple, oaks, sweetgum, sumac	Tank mix recommended rate of this product with 1 to 2 qt/100 gallons of Remedy and apply in late spring to early summer when leaves are fully expanded and mature. Use of a surfactant (0.25 to 0.5% v/v) is recommended. Spray to thoroughly wet foliage. For best results on blackberry, treat during or after bloom.
annual broomweed, bitterweed, bitter sneezeweed, bullnettle, bursage (bur ragweed), bull thistle, buffalo bur, camphorweed, cocklebur, common ragweed, croton, gray goldaster, lanceleaf ragweed, marshelder (sumpweed), musk thistle, narrowleaf goldaster, prickly lettuce, smartweed, sunflower, wild carrot, silverleaf nightshade, tasajillo, upright prairie cone flower, western hosenettle, western ragweed, yankeeweed	Apply when target weeds are 2 to 3 inches tall until early flowering.
flameleaf sumac honeylocust,	Apply in spring when leaves are fully expanded and mature. Use of a surfactant (0.25 to 0.5% v/v) is recommended. Spray thoroughly to wet foliage.
tropical soda apple	Apply when plant begins to flower.

1-2 Gallons/100 Gallons of Spray: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds:	
Brush Species	Specific Use Directions
Macartney rose multiflora rose	Apply in spring or fall when conditions are favorable for plant growth. High volume application is recommended for control of large undisturbed clumps or small regrowth.

2 Gallons/100 Gallons of Spray: Apply at the indicated stage of growth to control the following woody plants or broadleaf weeds:	
Weed or Brush Species	Specific Use Directions
Chinese tallow tree	Apply in spring or fall when conditions are favorable for plant growth.
cactus, pricklypear or cholla	Applications may be made throughout the year. Spray to wet all pads to runoff. Use of a surfactant (0.25-0.5% v/v) is recommended. Water soluble dye may be added to the spray mixture to mark treated plants.
common goldenweed, Drummond's goldenweed	Apply in spring (April-June) when favorable growing conditions result in substantial canopy development.
poisonous plants such as: groundsel (Senecio spp.), garbancillo (Wooton loco), and Woolly loco	Apply in fall or winter when moisture conditions are favorable. Herbicide treatment may increase palatability of poisonous plants. Treated areas should not be grazed until the toxic plants have dried up and lost their palatability.

TREATMENT AFTER PLANTING GRASSES, INCLUDING CONSERVATION RESERVE PROGRAM (CRP) ACRES

Weed Control Prior to Seeding Planting Grasses

This product may be applied to control weeds prior to planting **cool season grasses**.

Apply this product at 4 pints per acre or less depending on the target species. This product may be tank-mixed with Glyphomax Plus (glyphosate) to control grasses prior to seeding.

- To optimize weed control, minimal disturbance of the treatment area with the seeding operation is suggested. The site should be left undisturbed for a minimum of 21 days prior to seedbed preparation or seeding. To optimize weed control and reduce the potential for injury of seeded grasses, increase the interval between application of this product and planting grass seed.
- Do not plant smooth brome grass for 60 days after treatment.

15/17

Perennial Grasses

Applications of this product to perennial grasses should be made only after perennial grasses are well established as indicated by vigorous growth and a well-developed secondary root system.

Sprigged Bermudagrass: This product at 1.5 pints per acre or less can be used on sprigged bermudagrass once the runners (stolons) have reached 6 inches in length and growing conditions are favorable.

Overseeding: This product at rates of 1.5 pints per acre or less can be applied to permanent pastures that have been over seeded with small grains (such as barley, forage sorghum, oats, rye, ryegrass, sudangrass or wheat) grown for pasture or hay only. Young seedling small grains or grasses are sensitive to this product. This product should not be applied until overseeded grasses are well established and at tillering stage of growth or later.

Precautions:

- Applications of this product to established warm season grasses such as bermudagrass during initial greenup in early spring could delay or suppress emergence of new growth. If temporary suppression of new growth cannot be tolerated, application of this product should be made prior to greenup or after vigorous vegetative growth has resumed.
- Do not use this product if legumes are a desired cover during CRP.
- Conditions unfavorable to plant growth, such as drought, will increase potential for injury to grasses at all stages of growth.
- Crop Rotation: Do not rotate to grain sorghum (milo) if greater than 4 pints per acre of this product has been applied. For rates below 4 pints per acre, do not plant grain sorghum for 8 months after application. This product is not intended for use on land planted to sweet sorghum. To avoid potential crop injury, planting of small grains should be delayed a minimum of 60 days of soil temperatures above 40°F following application, except in Idaho, North Dakota, Nebraska, Montana, Oregon, South Dakota, Washington and Wyoming, where the minimum interval should be 90 days.
- After CRP, do not plant broadleaf crops in treated acres until an adequately sensitive bioassay (described below) shows that no detectable picloram is present in the soil.

Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application. The test area should sample field conditions such as soil texture, soil pH, drainage, and any other variable that could affect the seed bed of the new crop. The field bioassay can be initiated at any time between harvest of the treated crop and the planting of the rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the test rotational crop; plant only a labeled crop such as pasture grasses, small grains (barley, oats, rye or wheat), or, after a rotational interval of 8 months, grain sorghum.

Mixing Instructions

Ground or Aerial Application - For Use With Water Alone

Start with about half the required amount of water in the spray tank. With agitation operating, add the required amount of this product. If a surfactant is needed, it should be added as the remainder of the required water is added to complete the spray mix. When using a drift control additive, carefully follow the manufacturer's directions. Complete dispersion and uniform mixing is essential to proper performance of drift control additives. This can be aided by thorough circulation through a mixing pump with moderate to high shearing action.

Use With Oil/Water Emulsions

Ground Application: Add oil to the total spray mix at a rate of 5 to 10% of the total mix, up to a maximum of 1 gallon of oil per acre, using agricultural spray emulsifiers and mixing procedures given below.

Aerial Application: Use oil and water in the spray mixture in a 1:5 ratio (1 part oil to 5 parts water), up to a maximum of 1 gallon of oil per acre using mixing procedures given below.

Mixing Instructions for Oil/Water Emulsions (Batch Mixing)

With continuous, vigorous agitation:

1. Add to the spray tank half the amount of water to be used.
2. Add the amount of this product required for the total volume of spray being mixed.
3. Premix the required amount of oil with an emulsifier such as Sponto 712 or Triton X-100, using the manufacturer's recommended rate of emulsifier per gallon of oil. Add the oil-emulsifier premix to the spray tank.
4. Finally, add the remaining amount of water required to bring the spray batch to the desired total volume.
5. Maintain agitation in the spray tank during application.

Mixing with Liquid Fertilizer for Broadleaf Weed Control in Rangeland and Permanent Grass Pastures

This product may be tank mixed with liquid fertilizers and used in foliar application for weed control and fertilization of rangelands and permanent grass pastures. Avoid using liquid fertilizers in applications to brush as efficacy may be reduced. Use liquid fertilizers at rates recommended by supplier or local Extension Service Specialist.

Compatibility with Liquid Fertilizer: Prior to large scale batch mixing, conduct a "jar test" for spray mixture compatibility by mixing each component in the required order and proportion in a clear glass jar. Close the jar and agitate the mixture until evenly dispersed. Use of a compatibility agent is indicated if components of the mixture do not disperse readily or do not remain dispersed after mixing. Use of a compatibility aid such as Unite or Complex is recommended to help obtain and maintain a uniform spray solution during mixing and application. **Compatibility is best with straight liquid nitrogen fertilizer solutions. Mixing with N-P-K fertilizer**

16/17

solutions or suspensions is more difficult and should not be attempted without first conducting a successful jar test. Agitation in the spray tank must be vigorous to compare with jar test agitation.

Suggested Mixing and Application Procedure

With continuous vigorous agitation:

1. Add half the amount of liquid fertilizer to the spray tank.
2. Add compatibility aid such as Unite or Compex at 1 quart per 100 gallons of total spray mix.
3. First add the amount of this product needed for the total spray mixture. Mixing with N-P-K fertilizer solutions may be improved by premixing this product with water (1 part this product to 25-30 parts water) before adding to the spray tank.
4. Add the remaining liquid fertilizer to produce the needed total spray volume.
5. Apply as soon as mixing is complete, maintaining continuous, vigorous agitation throughout mixing and application without interruption.

Application during very cold (near freezing) weather is not advisable. The likelihood of mixing or compatibility problems with liquid fertilizer increases under cold conditions.

Do not store the spray mixture.

Note: Do not use spray equipment for application of other products to land planted, or to be planted, to susceptible crops or desirable sensitive plants, **unless** it has been determined that all phytotoxic herbicide residue has been removed by thorough cleaning of the equipment. See "Cleaning Instructions for Sprayer Equipment" General Use Precautions section of this label.

Cleaning Instructions for Spray Equipment

To avoid injury to desirable plants, equipment used to apply this product should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use. Flush the entire system at least three times with water, and dispose of rinse water in non-cropland area away from water supplies.
2. During the second rinse, add 1 quart of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 min.). Let the solution stand for several hours, preferable overnight.
3. Flush the solution out the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Nozzles and screens should be removed separately.

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OKLAHOMA SUPPRESSION OF SEEDLING AND REGROWTH MESQUITE IN CRP, RANGELAND AND PERMANENT PASTURES

Directions for Use

This product may be applied as a broadcast foliar spray at 1.5 to 3.0 pints per acre for suppression of seedling and regrowth mesquite plants. Apply this product when soil moisture and temperatures are favorable for plant growth and after the foliage from new seasonal growth on mesquite plants has fully expanded and changed from a light green to dark green color. Use highest rate in rate range where the density or height of woody plants is greater or when growth conditions are less than optimal.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY

17/17

SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(NEW))