

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



Office of Pesticide Programs

NuFarm Americas, Inc.
150 Harvester Dr.
Burr Ridge, IL 60527

Subject: Amendment
EPA Reg. No.: 228-531
Quincept Herbicide

MAR 11 9 2008

Dear Mr. Head,

The Agency has received your application for an amendment, dated December 19, 2007. The label described above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided the following modifications are made:

1. Remove box for the text beginning, "Follow manufacturers instructions for cleaning/maintaining PPE....," and place immediately following PPE requirements, ahead of "Engineering Controls."
2. Add a User Safety Recommendations box containing the following language:
 - a. "Users should wash hand before eating, drinking, chewing gum, using tobacco, or using the toilet."
 - b. "Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing."
 - c. "Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

This labeling supersedes all previous accepted labeling for this product. Please submit one (1) copy of the final printed label before the product is released for shipment. A stamped copy of the accepted label is enclosed for your records. If you have any questions, please contact Kathryn Montague (703-305-1243 or montague.kathryn@epa.gov).

Sincerely,

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2 8 9

Quincept™ Herbicide

ACCEPTED
with COMMENTS
In EPA Letter Dated:

MAR 19 2008
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

FOR SELECTIVE BROADLEAF AND CERTAIN GRASSY WEED CONTROL

CONTAINS 2,4-D, QUINCLORAC AND DICAMBA

228-531

CONTROLS: DANDELION, CHICKWEED, BLACK MEDIC, KNOTWEED, PLANTAIN, OXALIS, CLOVER, COCKLEBUR, THISTLE, LARGE AND SMOOTH CRABGRASS AND MANY OTHER SPECIES OF BROADLEAF AND CERTAIN GRASSY WEEDS; SOME OF WHICH ARE LISTED ON THIS LABEL.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	13.24%
Quinclorac (3,7-Dichloro-8-quinolinecarboxylic Acid)**	8.25%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	1.38%

OTHER INGREDIENTS:

TOTAL: 77.13%
100.0%

By Isomer Specific AOAC Method, Equivalent to:

* 2,4-Dichlorophenoxyacetic Acid	11.34%, 1.00 lbs./gal.
** 3,7-Dichloro-8-quinolinecarboxylic Acid	8.25%, 0.75 lbs./gal.
*** 3,6-Dichloro-o-Anisic Acid	1.38%, 0.125 lbs./gal.

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-531
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS

00228.00531.20071218.Pending
NUP-12D02

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or on clothing. Wear protective eyewear. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks, and
- Chemical-resistant gloves

See engineering controls for additional requirements and exceptions.

Engineering Controls Statements: If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. If this container is five gallons or more in capacity, do not pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

USER SAFETY RECOMMENDATIONS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material (such as nitrile, butyl, neoprene, and/or barrier laminate), chemical-resistant footwear plus socks, protective eyewear and chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter or allow others to enter treatment areas until sprays have dried.

PRODUCT INFORMATION

Quincept combines Quinclorac, 2,4-D and Dicamba to provide broad-spectrum control of grassy and broadleaf weeds.

- Post-emergent control of a comprehensive list of both grassy and broadleaf weeds
- Excellent clean-up product for areas that did not receive a pre-emergent grass herbicide application in the spring
- Controls newly germinated 1-2 leaf crabgrass, to 1-tiller crabgrass and when crabgrass has matured to 5 tillers or greater
- Quinclorac contributes grassy weed control and is absorbed by foliage and roots and translocated throughout the plant. Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death

USE SITES

Quincept may be applied postemergence to residential and nonresidential turfgrasses. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms.

Highly Tolerant Species	Moderately Tolerant Species	Susceptible (Do Not Use on these grass species)
Bluegrass, Annual (<i>Poa annua</i>) Bluegrass, Kentucky Fescue, Tall Ryegrass, Annual Ryegrass, Perennial	Bentgrass, Creeping ^{1,2} Bermudagrass, Common ¹ Bermudagrass (Hybrids) ¹ Bluegrass, Rough (<i>Poa trivialis</i>) Fescue, Chewing's Fescue, Fine ² Fescue, Hard Fescue, Red Zoysiagrass	Bahiagrass Bentgrass, Colonial Bentgrass, Seaside Buffalograss Carpetgrass Centipedegrass Dichondra Paspalum, Seashore St. Augustinegrass

1) Yellowing that may occur on these species can be reduced by the addition of chelated iron or sprayable soluble nitrogen
 2) Only apply to fine fescue when it is part of a blend
 3) Application to established creeping Bentgrass must be applied in 2-3 split applications at 1.1 to 1.9 ounces per 1,000 sq. ft.
 Do not exceed 1.5 pounds a.i. Quinclorac per acre per year. Sequential application(s) should be timed 21 days apart.
 Broadleaf weed control may be reduced due to sequential applications.
 Do NOT use on golf course greens, tees and collars
 Do NOT use on lawns with desirable clovers or legumes or on ornamentals

Warm season grasses such as Common Bermuda and Zoysia may see temporary discoloration. Use reduced rates if grass is stressed from heat or drought. Do not apply during growth stages from dormancy to green-up and from green-up to dormancy.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed the maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

TIMING

Apply Quincept to actively growing weeds as a postemergence broadcast or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds. Under certain conditions, application of Quincept made to annual grasses at 2-4 tiller may not provide complete control.

NEWLY SEEDED AREAS:

The application of Quincept to grass seedlings is not recommended until after the third mowing.

NEWLY SODDED, SPRIGGED, OR PLUGGED AREAS:

The application of Quincept to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Delay applications for 4 weeks after seeding and emergence of Kentucky bluegrass, perennial ryegrass and fine fescues.

SEEDING:

Delay applications for 4 weeks after seeding and emergence of turf species listed on this label.

MOWING:

Do not mow for two days before or two days after application. Clippings for the first three mowings should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

IRRIGATION AND RAINFALL

If soil moisture is not sufficient prior to application, irrigation may improve weed control. For best results, DO NOT water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is required.

Do not apply this product by air or through any type of irrigation equipment.

MIXING INSTRUCTIONS

Begin with a clean spray tank. Fill the spray tank with one-half the required amount of clean water. Slowly add Quincept while agitating, then complete filling the tank with water. Maintain continuous agitation until spraying is complete. If left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Adding adjuvants may cause slight leaf burn, but turf vigor is not reduced. Delaying applications when relative humidity and temperatures are high may help to avoid potential for leaf burn and turfgrass damage. Low mowing heights may also increase the possibility of turf injury. The addition of chelated iron or sprayable solution nitrogen fertilizer will reduce slight yellowing.

For best results, the addition of methylated seed oil is recommended when it meets all of the following criteria:

- be nonphytotoxic
- contain only EPA-exempt ingredients
- provide good mixing quality in the jar test
- be successful in local experience

Do not include additives when tank-mixing with emulsifiable concentrate (EC) products as this may cause phytotoxicity. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

Quincept may be tank mixed with EPA-registered preemergent herbicides (if compatible) for extended residual control.

The following compatibility test should always be performed prior to full-scale tank mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

Sprayer Cleaning: Clean application equipment thoroughly before and after application to prevent cross-contamination. Use a strong detergent or approved spray tank cleaner and rinse thoroughly.

SPRAYING INSTRUCTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Do not apply at wind speeds greater than 15 mph.

Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Avoid broadcast applications when air temperature exceeds 90 degrees. When using small, spot treatment applications in temperature over 90 degrees, turf injury may occur.

SPRAY VOLUMES:

Higher water volumes are generally required to control grassy weeds. Use 20 to 300 gallons per acre (.5 to 6.9 gallons per 1,000 per square feet) and a spray pressure of 20 to 40 psi. Higher water volumes should be used for control of dense weed populations to ensure weed foliage is completely covered.

LOWER VOLUME EQUIPMENT:

Use at least 0.5 gallons of water per 1,000 square feet. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE:

For all grasses (1) do not overlap spray patterns; (2) use reduced rates, if grass is stressed from heat, drought, etc.; and (3) follow CDA equipment spray instructions.

Make no more than 2 applications per year.

POST EMERGENT BROADLEAF WEED CONTROL

Quincept will control or suppress the following list of broadleaf weeds. For best results, apply this product when weeds are actively growing and in early stages of growth. More mature weeds will be more difficult to control and may require a second application. Mature, drought stressed weeds will be more difficult to control so adequate soil moisture is preferred. Adverse or extreme environmental conditions such as poor soil conditions, high temperatures, drought and cultural conditions may affect the performance of this product.

Do not broadcast apply this product above 90°F. Spot treatments above 90°F may result in some turf injury.

BROADLEAF WEEDS CONTROLLED

Alder	Canada thistle	Florida betony	Milk vetch	Redroot pigweed
Annual yellow sweet clover	Carolina geranium	Florida pusley	Milkweed bloodflower	Red sorrel
Artichoke	Carpetweed	Frenchweed	Mugwort	Redstem filaree
Austrian fieldcress	Catchweed bedstraw	Galinsoga	Morningglory	Rough cinquefoil
Bedstraw	Catsear	Garlic mustard	Mousear chickweed	Rough fleabane
Beggartick	Catnip	Goathead	Musk thistle	Roundleafed marigold
Biden	Chickweed	Goatsbeard	Mustard	Rush
Bindweed	Chicory	Goldenrod	Narrowleaf plantain	Russian pigweed
Bird vetch	Cinquefoil	Ground ivy	Narrowleaf vetch	Russian thistle
Bitterweed	Common chickweed	Gumweed	Nettle	St. Johnswort
Bitter wintercress	Common mullein	Hairy bittercress	Orange hawkweed	Scarlet pimpernel
Black-eyed Susan	Common sowthistle	Hairy fleabane	Oxalis	Scotch thistle
Black medic	Corn chamomile	Hawkweed	Oxeye daisy	Sheep sorrel
Black mustard	Creeping Jenny	Heatall	Parsley-piert	Shepherdspurse
Black-seed plantain	Crimson clover	Heartleaf drymary	Parsnip	Slender plantain
Blessed thistle	Croton	Hedge bindweed	Pearlwort	Smallflower galinsoga
Blue lettuce	Cudweed	Hedge mustard	Pennycress	Smartweed
Blue vervain	Curly dock	Hemp	Pennywort	Smooth dock
Box elder	Curly indigo	Henbit	Peppergrass	Smooth pigweed
Bracted plantain	Dandelion	Hoary cress	Pepperweed	Sneezeweed
Brassbuttons	Dead nettle	Horsetail	Pigweed	Southern wild rose
Bristly oxtongue	Dock	Indiana mallow	Pineywoods bedstraw	Sowthistle
Broadleaf dock	Dollar Weed	Ironweed	Plains coreopsis	Spanishneedle
Broadleaf plantain	Dogbane	Jewelweed	Plantain	Spatterdock
Broomweed	Dogfennel	Jimsonweed	Poison hemlock	Speedwell
Buckhorn	Elderberry	Kochia	Poison ivy	Spiny amaranth
Buckhorn plantain	English daisy	Knawel	Poison oak	Spiny cocklebur
Bulbous buttercup	Fall dandelion	Knotweed	Pokeweed	Spotted catsear
Bull nettle	False dandelion	Lambsquarter	Poorjoe	Spotted knapweed
Bull thistle	False flax	Lespedeza	Povertyweed	Spotted spurge
Burdock	False sunflower	Locoweed	Prostrate spurge	Spurge
Burning nettle	Fiddleneck	Lupine	Prostrate vervain	Spurweed
Bur ragweed	Field bindweed	Mallow	Puncture vine	Stinging nettle
Burweed	Field pansy	Marshelder	Purslane	Stinkweed
Buttercup	Flea bane (daisy)	Matchweed	Ragweed	Stitchwort
	Flixweed	Mexicanweed	Red clover	Strawberry clover

Sumac	Velvet leaf	Western salsify	Wild onion	Woolly morning-glory
Sunflower	Venice mallow	White clover	Wild parsnip	Woolly plantain
Sweet clover	Veronica	White mustard	Wild radish	Wormseed
Tall nettle	Vervain	Wild aster	Wild rape	Yarrow
Thistle	Vetch	Wild buckwheat	Wild strawberry	Yellow rocket
Tick trefoil	Violet, Wild	Wild carrot	Wild sweet potato	Yellowflower
Toadflax	Virginia buttonweed	Wild four-o'clock	Wild vetch	pepperweed
Trailing crownvetch	Virginia creeper	Wild garlic	Willow	and other broadleaf
Tumble mustard	Virginia pepperweed	Wild geranium	Witchweed	weeds.
Tumble pigweed	Wavyleaf bullthistle	Wild lettuce	Woodsorrel	
Tumbleweed	Western clematis	Wild marigold	Woolly croton	

POST EMERGENT CONTROL OF GRASSY WEEDS

This product can provide control and suppression of certain grassy weeds. For best results, apply this product when weeds are actively growing and in early stages of growth. More mature grasses will be more difficult to control and may require a second application. Mature, drought stressed grassy weeds will be more difficult to control so adequate soil moisture is preferred. Adverse or extreme environmental conditions such as poor soil conditions, high temperatures, drought and cultural conditions may effect the performance of this product. Do not broadcast apply this product above 90°F. Spot treatments above 90°F may result in some turf injury.

GRASSY WEEDS CONTROLLED IN ORNAMENTAL LAWNS AND TURF

COMMON NAME	COMMENTS
Barnyardgrass Crabgrass, Large Crabgrass, Smooth Foxtail, Giant Foxtail, Green Foxtail, Yellow Signalgrass, Broadleaf	Under certain conditions annual grasses at the 2 to 4 tiller stage may not be completely controlled and a sequential application may be needed at 14 to 21 days. Do not exceed two applications per year.

USE RATES

Quincept may be applied postemergence to residential and nonresidential turfgrasses according to the following rate chart. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms. Not for use on sod farms in Arizona.

APPLICATION GUIDELINES			
APPLICATION	RATE	SPRAY VOLUME	MAXIMUM APPLICATIONS
Single	7 to 8 pints/A or 112 to 128 ounces/A	20 to 300 gal/A or .5 to 6.9 gal/1,000 sq. ft.	2 per year
Spot Treatment	2.6 to 2.9 fluid ounces per 1,000 sq. ft.	1 gal/1,000 sq. ft.	

Application Guidelines:

- Ensure that spray volumes are adequate to completely cover weeds, especially when dense weed infestations make it difficult to completely cover the foliage of target weeds.
- Early applications of this product will not control weeds germinating later in the season so a second application may be necessary.
- If turf injury is observed after the first application, delay the second application until turf has recovered.
- Dual Applications: A single application should control less mature, light weed populations; however, a second application may be required to control heavy, mature broadleaf weed infestations. Under certain conditions, annual grasses in the 2 to 4 tiller stages may require a sequential application. If a second application is required, allow 14-21 days between applications.
- For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV121807A)

Quincept is a trademark of Nufarm Americas Inc.

Advertising Claims:

Proven performance

"Three-In-One" or "3-in-1"

Grassy and broadleaf weed control

Delivers a powerful punch

Crabgrass control

"Complete Formula"

Controls over 200 broadleaf weeds

Controls crabgrass, foxtail, Barnyardgrass and Signalgrass

Designed to pick up...when your pre-emerge lets go

Perfect for late season weed escapes