


228-536

12/07/2007

1/15

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p>	EPA Reg. Number: 228-536	Date of Issuance: DEC 7 2007
	Term of Issuance: Unconditional with label comments	
Name of Pesticide Product: NUP06013 HERBICIDE		
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)		
Name and Address of Registrant (include ZIP Code): Mr. George Meindl Nufarm Americas, Inc. 150 Harvester Drive, Suite 200 Burr Ridge, IL 60527		
<p><small>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</small></p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data. 2. Make the following changes to your labeling: <ol style="list-style-type: none"> a. Change the registration number to "228-536" b. Make all of the changes detailed in the attached document "Summary of Comments on Polaris specimen.qxd" 		
		12-7-07

2/15

Page 2
EPA Reg. No. 228-536

3. Submit final labeling for this product **within 30 days** of the date of this letter.
4. Submit studies of one year storage stability (Guideline 830.6317) and corrosion characteristics (Guideline 830.6320). The observations must be made at 0, 3, 6, 9, and 12 month intervals. The results should be submitted to the Agency in an electronic file and a paper copy.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

If you have any questions about this letter, please contact Tobi Colvin-Snyder at 703-305-7801.



Jim Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505C)

Summary of Comments on Polaris specimen.qxd

Page: 2

Author: tsnyder
Subject: Cross-Out
Date: 11/26/2007 3:17:37 PM -05'00'

⌘

Author: tsnyder
Subject: Replacement Text
Date: 11/26/2007 3:17:00 PM -05'00'

⌘ Causes moderate eye irritation.

Author: tsnyder
Subject: Cross-Out
Date: 11/26/2007 3:17:51 PM -05'00'

⌘

Author: tsnyder
Subject: Comment on Text
Date: 11/26/2007 3:24:18 PM -05'00'

⌘ Move this statement to above "If on skin...", since eye irritation is tox category III, and the other endpoints are IVs.

Page: 5

Author: tsnyder
Subject: Replacement Text
Date: 11/26/2007 4:12:24 PM -05'00'

T_A specified

Author: tsnyder
Subject: Replacement Text
Date: 11/26/2007 4:13:16 PM -05'00'

T_A specified

Author: tsnyder
Subject: Inserted Text
Date: 11/27/2007 9:53:13 AM -05'00'

T_A Tank mix products may be used provided that the label of the tank mix product does not prohibit such mixing.

Author: tsnyder
Subject: Inserted Text
Date: 11/27/2007 9:54:34 AM -05'00'

T_A instructions for all products used.

Page: 7

Author: tsnyder
Subject: Cross-Out
Date: 12/7/2007 5:06:13 PM -05'00'

T Not registered for this use.

Author: tsnyder
Subject: Comment on Text
Date: 11/27/2007 10:15:12 AM -05'00'

T While not mandatory, I suggest changing this text to "NUP 06013" for greater clarity.

6/15

NUP06013

FOR CONTROL OF VEGETATION ON FORESTRY SITES

ACTIVE INGREDIENT:

Isopropylamine salt of Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)* 27.6%

OTHER INGREDIENTS: 72.4%

TOTAL 100.0%

*Equivalent to 22.6% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid or 2 pounds acid per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527



NET CONTENTS GALS.

000228-00000.20070228.
NUP06013

ACCEPTED
with COMMENTS
in EPA Letter Dated

DEC 7 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-536

7/15

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

~~Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.~~

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- chemical-resistant gloves, such as barrier laminate, butyl rubber or polyethylene
- shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS	
Users should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored, and applied only in stainless steel, fiberglass, plastic, and plastic-lined steel containers. Do not mix, store, or apply this product or spray solutions of this product in unlined steel (except stainless steel) containers or spray tanks.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This herbicide is phytotoxic at extremely low concentrations. Non-target plants may be adversely affected from drift.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. The requirements in this box apply to use on trees being grown for sale or other commercial use, or for commercial seed production, or for production of timber or wood products, or for research purposes.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical-resistant gloves, such as barrier laminate, butyl rubber or polyethylene
- shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Noncrop weed control is not within the scope of the Worker Protection Standard. See the GENERAL INFORMATION section of this label for a description of noncrop sites.

Do not enter treated areas without protective clothing until sprays have dried.

This product may be used only in accordance with recommendations and restrictions in this label. Keep containers closed to avoid spills and contamination.

IMPORTANT

Do not use on food or feed crops. Do not apply to the inside of ditches used to transport irrigation water. Keep from contact with fertilizers, insecticides, fungicides, and seeds to prevent unintentional exposure of desirable vegetation to this product. Do not apply or drain or flush equipment on or near sensitive desirable plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on Christmas trees. Thoroughly clean application equipment after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

GENERAL INFORMATION

This product is an aqueous formulation that is readily mixable with water, diesel oil, or recommended seed oils and penetrating oils. For foliar applications, this product may be mixed with water as the spray carrier or an emulsion carrier may be prepared by mixing this product into water and then adding a suitable seed oil at 12 to 50%, by volume. This product is to be mixed with water or a penetrating oil and applied as a spray to cut stumps. This product should be mixed with a penetrating oil for application to the basal area of brush and trees. Adequate agitation should be maintained with all emulsion mixtures of this product to prevent phase separation. Prior to actual tank mixing with other products, herbicides and carrier oils, compatibility testing in small containers is recommended!

This product is recommended for vegetation control in forestry sites. Roadsides contiguous with the treated area may be included.

This product is recommended for control of vegetation in forestry site preparation, in directed applications for conifer release and for mid-rotation release using understory broadcast applications.

This product is also recommended for the control of undesirable vegetation along non-irrigation ditch banks and for the establishment and maintenance of wildlife openings, except in the state of California. See use directions for CUT STUMP TREATMENTS, TREE INJECTION TREATMENTS, FRILL OR GIRDLE TREATMENTS, THINLINE BASAL AND STEM APPLICATIONS, LOW VOLUME BASAL BARK TREATMENTS and LOW VOLUME FOLIAR APPLICATIONS.

This product may be applied on forestry sites that contain areas of temporary surface water caused by the collection of water between planting beds, in equipment ruts, or in other depressions created by forest management activities, except in the states of California and New York. It is permissible to treat drainage ditches, intermittent drainage, intermittently flooded low lying sites, seasonally dry flood plains, and transitional areas between upland and lowland sites when no water is present, except in the states of California and New York. Only the edge of drainage ditches can be treated for drainage ditches that contain water. It is also permissible to treat marshes, swamps, and bogs after water has receded, as well as seasonally dry flood deltas, except in the states of California and New York. Do not make applications to natural or man-made bodies of water such as lakes, reservoirs, ponds, streams, rivers and canals.

SYMPTOMOLOGY:

This product is readily absorbed through foliage, bark and roots and is translocated rapidly throughout the plant, with accumulation in meristematic regions. Treated plants stop growing soon after herbicide application. Chlorosis first appears in the youngest leaf tissue. In perennials, the herbicide is translocated into the roots, thus preventing resprouting. Chlorosis and tissue necrosis may not be apparent in some species for several weeks after application. Woody plants, brush, and trees may not display the full extent of herbicide control until several months following application.

9/15

MANAGING OFF-TARGET MOVEMENT

The following information is provided as general guidance for managing off-target movement. Specific use recommendations for this product may differ depending on the application technique used and the vegetation management objective.

Spray Drift: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Spray drift from applying this product may result in damage to sensitive plants adjacent to the treatment area. Only apply this product when the potential for drift to these and other adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal. Do not apply when the following conditions exist that increase the likelihood of spray drift from intended targets: high or gusty winds, high temperatures, low humidity, temperature inversions.

To minimize spray drift, the applicator should be familiar with and take into account the following drift reduction advisory information. Additional information may be available from state enforcement agencies or the Cooperative Extension on the application of this product.

The best drift management strategy and most effective way to reduce drift potential are to apply large droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity and Temperature Inversions).

CONTROLLING DROPLET SIZE

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift. Do not use nozzles producing a mist droplet spray.

Application Height: Making applications at the lowest possible height (aircraft, ground driven spray boom) that is safe and practical reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the application equipment (e.g. aircraft, ground) upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind: Drift potential is lowest between wind speeds of 3 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud, which can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind Erosion: Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

Aerial Application Methods and Equipment: Use 2 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

Managing spray drift from aerial applications: Applicators must follow these requirements to avoid off-target drift movement: 1) boom length - the distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor, 2) nozzle orientation - nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees, and 3) application height - without compromising aircraft safety, applications should be made at a height of 10 feet or less above the crop canopy or tallest plants. Applicators must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Ground Application (Broadcast): Use 5 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

10/15

CONIFER SITE PREPARATION TREATMENTS

This product may be used to control labeled grasses, broadleaf weeds, vines and brambles, and woody brush and trees on forest sites in advance of regeneration for the following conifer crop species:

Crop Species	Rate (oz/A)
Loblolly Pine (<i>Pinus taeda</i>)	48 – 80
Loblolly X Pitch Hybrid	48 – 80
Longleaf Pine (<i>Pinus palustris</i>)	48 – 80
Shortleaf Pine (<i>Pinus echinata</i>)	48 – 80
Virginia Pine (<i>Pinus virginiana</i>)	48 – 80
Slash Pine (<i>Pinus elliottii</i>)	40 – 64
Douglas-Fir (<i>Pseudotsuga menziesii</i>)	24 – 48
Western Hemlock (<i>Tsuga heterophylla</i>)	24 – 48
Coastal Redwood (<i>Sequoia sempervirens</i>)	24 – 48
California Red Fir (<i>Abies magnifica</i>)	24 – 40
California White Fir (<i>Abies concolor</i>)	24 – 40
Jack Pine (<i>Pinus banksiana</i>)	24 – 32
Lodgepole Pine (<i>Pinus contorta</i>)	24 – 32
Pitch Pine (<i>Pinus rigida</i>)	24 – 32
Ponderosa Pine (<i>Pinus ponderosa</i>)	24 – 32
Sugar Pine (<i>Pinus lambertiana</i>)	24 – 32
White Pine (<i>Pinus strobus</i>)	24 – 32
Black Spruce (<i>Picea mariana</i>)	24 – 32
Red Spruce (<i>Picea rubens</i>)	24 – 32
White Spruce (<i>Picea glauca</i>)	24 – 32

Use the recommended rate of this product per acre applied as a broadcast foliar spray for long-term control of labeled woody plants and residual control of herbaceous weeds. Within 4 to 6 weeks of treatment, grasses and other herbaceous weeds will be controlled and may provide fuel to facilitate a site preparation burn, if desired, to control conifers or other species tolerant to the herbicide.

For tracts to be planted with loblolly, loblolly x pitch hybrid, longleaf pine, shortleaf pine, Virginia pine and slash pine, this product may be applied at a rate of 64 oz per acre on areas that have little to no resprouting vegetation because of recent management activities such as harvesting, mechanical shearing, burning piling or bedding. Applications must be made after September 1.

MIXING AND APPLICATION INSTRUCTIONS FOR SITE PREPARATION

Apply the recommended rate of this product per acre in 5 to 20 gallons total spray carrier for helicopter applications or 5 to 40 gallons total spray carrier for mechanical or backpack ground spray applications. Enhanced brownout for burning and improved control of brush and grasses may be obtained by application of this product in 12 to 50% oil:water (volume:volume) emulsion carrier. Methylated or ethylated seed oils containing at least 50% esterified seed oil by volume are recommended. Mix this product into the water portion of the carrier thoroughly, then add the oil and mix thoroughly again to obtain a uniform emulsion. Use the higher label rates of this product and higher spray volumes when controlling particularly dense or multi-layered canopies of hardwood stands, or difficult to control species. Make applications during the growing season; beginning in the spring after full leaf expansion of the target weed or brush has occurred and complete applications before leaf drop in the fall.

Tank mixes may be necessary for chemical control of conifers and other species tolerant to this product in certain cases. Observe all precautions and restrictions on the product labels. Always follow the most restrictive label. Combinations with other products labeled for forest site preparation may kill certain plants such as legume's and blackberry which are desirable for wildlife habitat.

Do not plant seedlings of black spruce (*Picea mariana*) or white spruce (*Picea glauca*) on sites that have been site prepared with a broadcast application of this product or into the treated zone of spot or banded site preparation applications for three months following treatment or injury may occur.

HELICOPTER SPRAY EQUIPMENT

All precautions should be taken to minimize or eliminate spray drift. Applications should not be made under gusty conditions. The use of controlled droplet booms and nozzle configurations is recommended.

IMPORTANT: Do not make applications by fixed wing aircraft. Maintain adequate buffer zones. Thoroughly clean application and mixing equipment, including landing gear, immediately after use. Prolonged exposure of this product to uncoated steel (except stainless steel) surfaces may result in corrosion and failure of the exposed part.

HARDWOOD SITE PREPARATION TREATMENTS

For site preparation prior to planting hardwood species in the southeast and gulf coast states (Virginia to Texas), use this product at a rate of 48 oz per acre and spray before the end of July. Application in an emulsion carrier with a minimum of 12% oil is recommended. Do not plant hardwood seedlings before January of the year following site preparation or injury may occur.

11/15

DIRECTED FOLIAR APPLICATIONS FOR CONIFER RELEASE

This product may be applied as a directed spray using water or oil emulsion carrier for control and suppression of labeled brush and weed species. Directed spray applications may be made using low carrier volumes (generally 10 gallons total spray per acre or less) in labeled conifer stands of all ages by targeting the unwanted vegetation and avoiding direct application to the conifer. Ensure that the maximum labeled rates per acre listed for the conifer species are not exceeded.

Use directed foliar applications of this product for release of the following conifers from hardwood competition:

Crop Species	Rate (fl oz/Acre)
Loblolly Pine (<i>Pinus taeda</i>)	24 – 40
Loblolly X Pitch Hybrid	24 – 40
Virginia Pine (<i>Pinus virginiana</i>)	24 – 40
Longleaf Pine (<i>Pinus palustris</i>)	24 – 32
Pitch Pine (<i>Pinus rigida</i>)	24 – 32
Shortleaf Pine (<i>Pinus echinata</i>)	24 – 32
Slash Pine (<i>Pinus elliottii</i>)	24 – 32
White Pine (<i>Pinus strobus</i>)	16 – 32
Lodgepole Pine (<i>Pinus contorta</i>)	16 – 24
Douglas-Fir (<i>Pseudotsuga menziesii</i>)	16 – 24
Jack Pine (<i>Pinus banksiana</i>)	12 – 24
Black Spruce (<i>Picea mariana</i>)	12 – 24
Red Spruce (<i>Picea rubens</i>)	12 – 24
White Spruce (<i>Picea glauca</i>)	12 – 24

For applications directed to the foliage of undesirable brush mix 2 to 10% this product in water. For brush species with thick leaf cuticles or difficult to control species use oil emulsion carrier containing 12 to 50%, by volume, recommended oil diluent. Apply the spray solution or emulsion to at least two-thirds of each hardwood crown using backpack sprayers or hand held equipment. Do not spray to the point of runoff and avoid spraying the conifers for best results. For low volume foliar applications to control big leaf maple a 5% by volume solution or emulsion of this product is recommended.

Some minor conifer growth inhibition may be observed when release treatments are made during periods of active conifer growth. To minimize potential conifer height growth inhibition, release treatments may be made late in the growing season after formation of final conifer resting buds. To prevent possibility of conifer injury, do not apply this product when conifers are under stress from drought, diseases, animal or winter injury, or other stresses reducing conifer vigor.

Injury may occur to non-target or desirable hardwoods if they extend from the same root system as treated stems, or their root systems are grafted to those of the treated tree, or if their roots extend into the soil near treated trees.

BAG AND BROADCAST APPLICATIONS FOR CONIFER RELEASE

In Douglas-fir and Ponderosa pine stands, broadcast applications of this product up to 32 oz per acre are permissible when the trees are covered by bags prior to the application. The bags must prevent the spray mix from contacting the conifer foliage. For improved control of brush species, particularly evergreens, add a suitable seed oil at 5 to 12 % by volume. On sites with coarse textured soils (e.g. decomposed granite, pumice, sandy or rocky sites) or low levels of soil organic matter (generally 5% or less) significant conifer growth inhibition and mortality is possible. Do not use this treatment on these types of sites if conifer growth inhibition and mortality cannot be tolerated.

LATE ROTATION VEGETATION CONTROL IN WESTERN CONIFERS

In California, the Pacific Northwest and Inland Northwest, broadcast aerial applications of this product up to 48 oz per acre are permissible in conifer stands that are targeted for harvesting the year following treatment. Use a minimum spray volume of 15 gallons per acre. For improved control of brush species, particularly evergreens, add a suitable seed oil at 5 to 12 % by volume. Significant conifer injury or mortality must be expected. Do not use this treatment if conifer injury or mortality cannot be tolerated.

UNDERSTORY BROADCAST APPLICATIONS FOR MID-ROTATION RELEASE

This product may be applied as a broadcast application below the conifer canopy to control understory brush and suppress trees for labeled species. Ground spray machinery or hand held equipment may be used to broadcast this product in water or oil emulsion carrier below the crop tree canopy in a manner as to minimize spray contact by the live crown of crop trees.

Ensure that maximum labeled rates per acre listed for crop species below are not exceeded.

Crop Species	Maximum Rate (fl oz/Acre)
Loblolly Pine (<i>Pinus taeda</i>)	64
Loblolly X Pitch Hybrid	64
Virginia Pine (<i>Pinus virginiana</i>)	64
Longleaf Pine (<i>Pinus palustris</i>)	32
Pitch Pine (<i>Pinus rigida</i>)	32
Shortleaf Pine (<i>Pinus echinata</i>)	32
Slash Pine (<i>Pinus elliottii</i>)	32

12/15

CUT STUMP TREATMENTS

Mix 8.0-16.0 fluid ounces of this product in one gallon of water*, diesel oil, or a penetrating oil.

This product may be tank mixed with Tahoe® 3A, Tahoe® 4E, picloram, Patriot®-Razor/Razor® Pro, or Foresters® Non-Selective to control labeled species. Spray or brush this product's solution onto the cambium area of the freshly cut stump surface. Insure that this product's solution thoroughly wets the cambium area (the wood next to the bark) of the stump. The use of a surfactant or penetrating agent may improve uptake through partially callused cambiums. Applications can be made anytime during the year except during periods of heavy sap flow in the spring. Do not over apply causing puddling.

*Note: Use water as a diluent only when temperatures are sufficient to prevent freezing or add antifreeze (ethylene glycol) according to label directions to prevent freezing.

TREE INJECTION TREATMENTS

No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

Mix 8.0-12.0 fluid ounces of this product in one gallon of water*. Using standard injection equipment, apply 1 ml. of this product's solution at each injection site around the tree with no more than 1 inch intervals between cut edges. Insure that the injector completely penetrates the bark at each site.

*Note: Use water as a diluent only when temperatures are sufficient to prevent freezing or add antifreeze (ethylene glycol) according to label directions to prevent freezing.

FRILL OR GIRDLE TREATMENTS

Mix 8.0-12.0 fluid ounces of this product in one gallon of water*, diesel oil, or a penetrating oil. Using a hatchet, machete, or similar tool, make cuts through the bark and completely around the tree with no more than 2 inch intervals between cut edges. Spray or brush this product's solution into each cut until thoroughly wet.

*Note: Use water as a diluent only when temperatures are sufficient to prevent freezing or add antifreeze (ethylene glycol) according to label directions to prevent freezing.

THINLINE BASAL AND STEM APPLICATIONS

This product may be applied as a thinline basal or arcing application to the stems of susceptible species such as big leaf maple (Acer macrophyllum), willow (Salix spp.) and Eucalyptus (Eucalyptus spp.) with a stem ground line diameter of 3 inches or less. Mix 24 to 48 ounces of this product in one gallon of diesel oil or penetrating oil. Maintain uniform mixtures with frequent agitation. Direct a thin line of the spray solution to the stems beginning a few feet from the ground and descending toward the base of the tree making a zig-zag motion. Do not over apply causing puddling.

LOW VOLUME BASAL BARK TREATMENTS

Mix 8.0-12.0 fluid ounces of this product in one gallon of diesel oil or a penetrating oil. To control mixed brush species with up to 4 inch stem diameter at breast height, spray to wet the lower 12-18 inches of the stem with this product's oil mixture (include the root collar area). Do not over apply causing dripping or puddling. Maintain uniform mixtures with frequent agitation.

LOW VOLUME FOLIAR APPLICATIONS

This product may be applied as a low volume foliar application. Mix 3-5% of this product in water and adjuvant or in a penetrating oil. For small brush spray down on the crown to cover approximately 70% of the plant foliage. For larger brush insure coverage on as much of the crown as possible and spray at least two sides of the plant. May be tank mixed with other labeled herbicides. Use a tank mix of 3 to 5% of this product plus 15 to 20% Tahoe 4E in basal oil to control black locust, honey locust, hackberry, elms and other species listed on manufacturer's labels. Use the higher rate of this product (5%) in areas containing sassafras, oak, hickory, cherry, and maples or in the southern 2/3's of the U.S. A tank mix of 3% of this product + Tahoe 4E is effective in the Northeastern U.S.

SPRAY SOLUTION MIXING GUIDE FOR LOW VOLUME FOLIAR APPLICATIONS

AMOUNT OF SPRAY SOLUTION BEING PREPARED	DESIRED CONCENTRATION (fluid volume)			
	This Product		Tahoe® 4E	
	3%	5%	15%	20%
1 gallon	3.8 oz	6.4 oz	19.2 oz	25.6 oz
3 gallons	11.5 oz	19.2 oz	57.6 oz	76.8 oz
4 gallons	15.4 oz	25.6 oz	76.8 oz	102.4 oz
5 gallons	19.2 oz	32.0 oz	96.0 oz	1.0 gallon
50 gallons	1.5 gallons	2.5 gallons	7.5 gallons	10.0 gallons
100 gallons	3.0 gallons	5.0 gallons	15.0 gallons	20.0 gallons

INVERT EMULSIONS

This product can be applied as an invert emulsion carrier. The carrier is a thick invert water-in-oil spray emulsion designed to minimize spray drift and spray run off, resulting in more herbicide on the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Consult the invert chemical label for proper mixing directions. Do not exceed 3 quarts/Acre of this product.

WEEDS CONTROLLED

This product will provide postemergence control and some residual control of the following target vegetation species. Degree of control is both species and rate dependent.

GRASSES

The species of annual and perennial grasses controlled by this product include the following:

- | | | |
|---------------------------------------------------------|-------------------------------------------------|---------------------------------------------------|
| Annual bluegrass (<i>Poa annua</i>) | Feathertop (<i>Pennisetum villosum</i>) | Quackgrass (<i>Agropyron repens</i>) |
| Bahiagrass (<i>Paspalum notatum</i>) | Fescue (<i>Festuca spp.</i>) | Reed canary grass (<i>Phalaris arundinacea</i>) |
| Barnyardgrass (<i>Echinochloa crus-galli</i>) | Foxtail (<i>Setaria spp.</i>) | Saltgrass (<i>Distichlis stricta</i>) |
| Beardgrass (<i>Andropogon spp.</i>) | Giant reed (<i>Arundo donax</i>) | Sand dropseed (<i>Sporobolus cryptandrus</i>) |
| Bermudagrass (<i>Cynodon dactylon</i>) | Goosegrass (<i>Eleusine indica</i>) | Sandbur (<i>Cenchrus spp.</i>) |
| Big bluestem (<i>Andropogon gerardii</i>) | Guineagrass (<i>Panicum maximum</i>) | Smooth brome (<i>Bromus inermis</i>) |
| Broadleaf signalgrass (<i>Brachiaria platyphylla</i>) | Italian ryegrass (<i>Lolium multiflorum</i>) | Sprangletop (<i>Leptochloa spp.</i>) |
| Canada bluegrass (<i>Poa compressa</i>) | Itchgrass (<i>Rottboellia exaltata</i>) | Timothy (<i>Phleum pratense</i>) |
| Cattail (<i>Typha spp.</i>) | Johnsongrass (<i>Sorghum halepense</i>) | Torpedograss (<i>Panicum repens</i>) |
| Cheat (<i>Bromus secalinus</i>) | Junglerice (<i>Echinochloa colonum</i>) | Vaseygrass (<i>Paspalum urvillei</i>) |
| Cogongrass (<i>Imperata cylindrica</i>) ¹ | Kentucky bluegrass (<i>Poa pratensis</i>) | Wild barley (<i>Hordeum spp.</i>) |
| Crabgrass (<i>Digitaria spp.</i>) | Lovegrass (<i>Eragrostis spp.</i>) | Wild oats (<i>Avena fatua</i>) |
| Crowfootgrass (<i>Dactyloctenium aegyptium</i>) | Orchardgrass (<i>Dactylis glomerata</i>) | Wirestem muhly (<i>Muhlenbergia frondosa</i>) |
| Dallisgrass (<i>Paspalum dilatatum</i>) | <i>Panicum spp.</i> | Witchgrass (<i>Panicum capillare</i>) |
| Downy brome (<i>Bromus tectorum</i>) | Paragrass (<i>Brachiaria mutica</i>) | Woolly cupgrass (<i>Eriochloa villosa</i>) |
| Fall panicum (<i>Panicum dichotomiflorum</i>) | Phragmites (<i>Phragmites australis</i>) | |
| | Prairie cordgrass (<i>Spartina pectinata</i>) | |
| | Prairie threeawn (<i>Aristida oligantha</i>) | |

¹Use minimum of 48 oz per acre.

BROADLEAF WEEDS

The species of annual and perennial broadleaf weeds controlled by this product include the following:

- | | | |
|----------------------------------------------------|----------------------------------------------------------|---------------------------------------------------------|
| Arrowwood (<i>Pluchea sericea</i>) | <i>nauseosus</i>) | Rush skeletonweed (<i>Chondrilla juncea</i>) |
| Broom snakeweed (<i>Gutierrezia sarothrae</i>) | Henbit (<i>Lamium alexicaule</i>) | Russian knapweed (<i>Centaurea repens</i>) |
| Bull Thistle (<i>Cirsium vulgare</i>) | Hoary vervain (<i>Verbena stricta</i>) | Russian thistle (<i>Salsola kali</i>) |
| Burclover (<i>Medicago spp.</i>) | Horseweed (<i>Conyza canadensis</i>) | Saltbush (<i>Atriplex spp.</i>) |
| Burdock (<i>Arctium spp.</i>) | Indian mustard (<i>Brassica juncea</i>) | Shepherd's purse (<i>Capsella bursa-pastoris</i>) |
| Camphorweed (<i>Heterotheca subaxillaris</i>) | Japanese bamboo/knotweed (<i>Polygonum cuspidatum</i>) | Silverleaf nightshade (<i>Solanum elaeagnifolium</i>) |
| Carolina geranium (<i>Geranium carolinianum</i>) | Knotweed, prostrate (<i>Polygonum aviculare</i>) | Smartweed (<i>Polygonum spp.</i>) |
| Carpetweed (<i>Mullugo verticillata</i>) | Kochia (<i>Kochia scoparia</i>) | Sorrell (<i>Rumex spp.</i>) |
| Chickweed, mouseear (<i>Cerastium vulgatum</i>) | Lambsquarters (<i>Chenopodium album</i>) | Sowthistle (<i>Sonchus spp.</i>) |
| Clover (<i>Trifolium spp.</i>) | Little mallow (<i>Malva parviflora</i>) | Spurge, annual (<i>Euphorbia spp.</i>) |
| Cocklebur (<i>Xanthium strumarium</i>) | Milkweed (<i>Asclepias spp.</i>) | Stinging nettle (<i>Urtica dioica</i>) |
| Common chickweed (<i>Stellaria media</i>) | Miners lettuce (<i>Montia perfoliata</i>) | Sunflower (<i>Helianthus spp.</i>) |
| Common ragweed (<i>Ambrosia artemisiifolia</i>) | Mullein (<i>Verbascum spp.</i>) | Sweet clover (<i>Melilotus spp.</i>) |
| Cudweed (<i>Gnaphalium spp.</i>) | Nettleleaf goosefoot (<i>Chenopodium murale</i>) | Tansymustard (<i>Descurainia pinnata</i>) |
| Dandelion (<i>Taraxacum officinale</i>) | Oxeye daisy (<i>Chrysanthemum leucanthemum</i>) | Texas thistle (<i>Cirsium texanum</i>) |
| Desert camelthorn (<i>Alhagi pseudalhagi</i>) | Pepperweed (<i>Lepidium spp.</i>) | Velvetleaf (<i>Abutilon theophrasti</i>) |
| Diffuse knapweed (<i>Centaurea diffusa</i>) | Pigweed (<i>Amaranthus spp.</i>) | Western ragweed (<i>Ambrosia psilostachya</i>) |
| Dock (<i>Rumex spp.</i>) | Plantain (<i>Plantago spp.</i>) | Wild carrot (<i>Daucus carota</i>) |
| Dogfennel (<i>Eupatorium capillifolium</i>) | Pokeweed (<i>Phytolacca americana</i>) | Wild lettuce (<i>Lactuca spp.</i>) |
| Fiddleneck (<i>Amsinckia intermedia</i>) | Primrose (<i>Oenothera kunthiana</i>) | Wild parsnip (<i>Pastinaca sativa</i>) |
| Filaree (<i>Erodium spp.</i>) | Puncturevine (<i>Tribulus terrestris</i>) | Wild turnip (<i>Brassica campestris</i>) |
| Fleabane (<i>Erigeron spp.</i>) | Purple loosestrife (<i>Lythrum salicaria</i>) | Woollyleaf bursage (<i>Ambrosia grayi</i>) |
| Giant ragweed (<i>Ambrosia trifida</i>) | Purslane (<i>Portulaca spp.</i>) | Yellow starthistle (<i>Centaurea solstitialis</i>) |
| Goldenrod (<i>Solidago spp.</i>) | Pusley, Florida (<i>Richardia scabra</i>) | Yellow woodsorrel (<i>Oxalis stricta</i>) |
| Gray rabbitbrush (<i>Chrysothamnus</i> | Rocket, London (<i>Sisymbrium irio</i>) | |

VINES AND BRAMBLES

The species of vines and brambles controlled by this product include the following:

- | | | |
|---------------------------------------------------|---------------------------------------------------------|------------------------------------------------------|
| Field bindweed (<i>Convolvulus arvensis</i>) | Trumpet creeper (<i>Campsis radicans</i>) | Wild rose (<i>Rosa spp.</i>) ¹ |
| Hedge bindweed (<i>Calystegia sepium</i>) | Virginia creeper (<i>Parthenocissus quinquefolia</i>) | Including Multiflora rose (<i>Rosa multiflora</i>) |
| Honeysuckle (<i>Lonicera spp.</i>) ¹ | Wild buckwheat (<i>Polygonum convolvulus</i>) | Macartney rose (<i>Rosa bracteata</i>) |
| Morningglory (<i>Ipomoea spp.</i>) | Wild grape (<i>Vitis spp.</i>) | |
| Poison ivy (<i>Rhus radicans</i>) | | |
| Redvine (<i>Brunnichia cirrhosa</i>) | | |

¹Use higher labeled rates.

WOODY BRUSH AND TREES

The species of woody brush and trees controlled by this product include the following:

- | | | |
|-----------------------------------------------------------------|--------------------------------------------------------------------|-----------------------------------------------------------------|
| Alder (<i>Alnus spp.</i>) | Eucalyptus (<i>Eucalyptus spp.</i>) | Privet (<i>Ligustrum vulgare</i>) |
| American beech (<i>Fagus grandifolia</i>) | Hawthorn (<i>Crataegus spp.</i>) | Red alder (<i>Alnus rubra</i>) |
| Ash (<i>Fraxinus spp.</i>) ¹ | Hazel (<i>Corylus cornuta</i>) ⁵ | Red maple (<i>Acer rubrum</i>) ¹ |
| Aspen (<i>Populus spp.</i>) | Hickory (<i>Carya spp.</i>) ¹ | Saltcedar (<i>Tamarix pentandra</i>) |
| Australian pine (<i>Casuarina equisetifolia</i>) ⁵ | Holly (<i>Ilex spp.</i>) ^{1,4} | Sassafras (<i>Sassafras albidum</i>) |
| Autumn olive (<i>Elaeagnus umbellata</i>) | Including Gallberry (<i>Ilex glabra</i>) ^{4,5} | Scotch broom (<i>Cytisus scoparius</i>) ⁵ |
| Bald cypress (<i>Taxodium distichum</i>) ⁴ | Tall gallberry (<i>Ilex coriacea</i>) ⁴ | Sourwood (<i>Oxydendrum arboreum</i>) ² |
| Bigleaf maple (<i>Acer macrophyllum</i>) ¹ | Yaupon (<i>Ilex vomitoria</i>) ⁴ | Sumac (<i>Rhus spp.</i>) |
| Birch (<i>Betula spp.</i>) | Honeylocust (<i>Gleditsia triacanthos</i>) ⁵ | Sweetbay magnolia (<i>Magnolia virginiana</i>) ^{4,5} |
| Black locust (<i>Robinia pseudoacacia</i>) ⁵ | Huckleberry (<i>Gaylussacia spp.</i>) | Sweetgum (<i>Liquidambar styraciflua</i>) |
| Black oak (<i>Quercus kelloggii</i>) | <i>Lyonia spp.</i> | Sycamore (<i>Platanus occidentalis</i>) |
| Blackgum (<i>Nyssa sylvatica</i>) ² | Including Fetterbush (<i>Lyonia lucida</i>) | Tanoak (<i>Lithocarpus densiflorus</i>) ^{1,4,5} |
| Boxelder (<i>Acer negundo</i>) | Staggerbush (<i>Lyonia mariana</i>) | TITI (<i>Cyrtilla racemiflora</i>) ^{1,4} |
| Brazilian peppertree (<i>Schinus terebinthifolius</i>) | Madrone (<i>Arbutus menziesii</i>) | Tree of heaven (<i>Ailanthus altissima</i>) ⁵ |
| Ceanothis (<i>Ceanothis spp.</i>) | Manzanita, greenleaf (<i>Arctostaphylos patula</i>) ⁴ | <i>Vaccinium spp.</i> |
| Cherry (<i>Prunus spp.</i>) ^{1,2} | Maple (<i>Acer spp.</i>) | Including Blueberry (<i>Vaccinium spp.</i>) |
| Chinaberry (<i>Melia azedarach</i>) | Melaleuca (<i>Melaleuca quinquenervia</i>) | Sparkleberry (<i>Vaccinium arboreum</i>) |
| Chinese tallow-tree (<i>Sapium sebiferum</i>) | Mulberry (<i>Morus spp.</i>) ^{1,3} | Waxmyrtle (<i>Myrica californica</i>) ^{4,5} |
| Chinquapin (<i>Castanopsis chrysophylla</i>) ⁴ | Oak (<i>Quercus spp.</i>) ^{1,3} | (<i>Myrica cerifera</i>) ^{4,5} |
| Cottonwood (<i>Populus spp.</i>) | Persimmon (<i>Diospyros virginiana</i>) ² | Willow (<i>Salix spp.</i>) |
| Cypress (<i>Taxodium spp.</i>) | Poison oak (<i>Rhus diversiloba</i>) | Yellow-poplar (<i>Liriodendron tulipifera</i>) ¹ |
| Dogwood (<i>Cornus spp.</i>) ¹ | Popcorn-tree (<i>Sapium sebiferum</i>) | |
| Elderberry (<i>Sambucus spp.</i>) ⁵ | Poplar (<i>Populus spp.</i>) ² | |
| Elm (<i>Ulmus</i>) ⁵ | | |

¹Use higher labeled rates.

²Best control with applications prior to formation of fall leaf color.

³The degree of control may be species dependent.

⁴Oil emulsion carrier is recommended.

⁵Tank mix with Tahoe 4E as a basal or cut stump treatment

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not store below 10°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: FOR QUART, 1.0 GALLON, 2.5 GALLON, 15 GALLON AND 30 GALLON: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR FIELD KEG, MINI BULK AND BULK: When this container is empty, replace the cap and seal all openings that have been opened during use, and return the container to the point of purchase, or to a designated location. This container must only be refilled with the pesticide product. Do not reuse the container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. Check for leaks after refilling and before transport. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling. Disposal of container must be in compliance with state and local regulations.

15/15

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV022807)

Polaris, Patriot, Razor, Tahoe and Foresters' are registered trademarks of Nufarm Americas Inc.