

228-568

04/08/2008

1/11



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (H7505C)
 401 "M" St., S.W.
 Washington, D.C. 20460

EPA Reg. Number: 228-568
 Date of Issuance: APR 8 2008

Term of Issuance:
 Unconditional with label comments

Name of Pesticide Product:
Brazen SF Herbicide

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. George Meindl
 Nufarm Americas, Inc.
 150 Harvester Drive, Suite 200
 Burr Ridge, IL 60527


Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. **Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.**
2. **Make the following changes to your labeling:**
 - a. **Change the registration number to "228-568"**
 - b. **Make all of the changes detailed in the attached document "Summary of Comments on NUP060125 EPA draft 2 062807".**


 Jim Tompkins, PM 24
 Branch, (Registration Division 7505P)


4-8-08
 Date

- 3. Submit final labeling for this product **within 30 days** of the date of this letter.
- 4. Submit studies of one year storage stability (Guideline 830.6317) and corrosion characteristics (Guideline 830.6320). The observations must be made at 0, 3, 6, 9, and 12 month intervals. The results should be submitted to the Agency in an electronic file and a paper copy.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

If you have any questions about this letter, please contact Tobi Colvin-Snyder at 703-305-7801.



Jim Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505C)

BRAZEN™ SF

HERBICIDE

FOR CONTROL OF ANNUAL AND PERENNIAL BROADLEAF WEEDS IN ESTABLISHED TURFGRASS, INCLUDING, BUT NOT LIMITED TO, SOD FARMS.

ACTIVE INGREDIENT:

Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt*33.0%
Clopyralid: 3,6-dichloro-2-pyridinylcarboxylic acid, triethylamine salt**12.1%

OTHER INGREDIENTS: 54.9%

TOTAL: 100.0%

*Acid Equivalent Triclopyr: 23.7%, 2.25 lbs./gal.

**Acid Equivalent Clopyralid: 7.9%, 0.75 lbs./gal.

IN CALIFORNIA, OREGON AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY.

KEEP OUT OF REACH OF CHILDREN
WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-LAI
EPA EST. NO.

MANUFACTURED BY
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



NET CONTENTS: GALS.

MADE IN USA

000228-00000.20071017.NEW
NUP-06125

ACCEPTED
with COMMENTS
In EPA Letter Dated

APR 8 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-568

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING - AVISO**

Harmful if swallowed. Causes substantial but temporary eye injury. Do not get in eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selections chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- long-sleeved shirt and long pants
- shoes and socks, plus
- protective eyewear (goggles, face shield or safety glasses)



Clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate should be discarded. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protections Standard (WPS) for agriculture pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS	
Users should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and other invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when cleaning and/or disposing of equipment wash waters or rinsate.

Triclopyr has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

S/11

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product for ground application only and is to be used in accordance with its labeling and with the Worker Protection Standard (40 CFR part 170). This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE is required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material; shoes plus socks; protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Do not enter or allow people (or pets) to enter treatment area until sprays have dried.

GENERAL INFORMATION

This product controls a wide variety of broadleaf weeds in established cool and warm season turfgrasses, including, but not limited to, turfgrass in sod farms with noted exceptions.

Apply this product to healthy, well established turfgrass with a well-anchored root system. This product may discolor and/or stunt turfgrass that is not well established or is stressed or weakened due to factors which damage or weaken turf, such as unfavorable climactic conditions, temperature extremes, drought and nematodes.

This product is recommended for use on the following turfgrass species:

ESTABLISHED COOL SEASON TURFGRASS	
COMMON NAME	SCIENTIFIC NAME
bentgrass [†]	<i>Agrostis species</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra var. communtata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceac</i>
ryegrass, perennial	<i>Lolium perenne</i>
[†] On bentgrass, do not apply more than 1 pint of product per acre (0.37 fl. oz. or 2.5 tsp. per 1000 sq.ft.) unless turfgrass injury can be tolerated. To minimize turfgrass injury, additional applications should be made at least four weeks apart. Avoid swath overlaps.	

ESTABLISHED WARM SEASON TURFGRASS [†]	
COMMON NAME	SCIENTIFIC NAME
bahiagrass [†]	<i>Agrostis species</i>
Bermudagrass ^{††}	<i>Poa pratensis</i>
buffalograss	<i>Festuca rubra var. communtata</i>
centipedegrass	<i>Festuca rubra</i>
fescue, tall (growing in warm season areas)	<i>Festuca ovina</i>
zoysiagrass	<i>Zoysia juponica</i> , <i>Zoysia tenuifolia</i>
[†] Do not treat warm season turfgrass with this product when mowing height is less than 1/2 inch. Do not apply more than 1 pint of product per acre (0.37 fl. oz. or 2.5 tsp. per 1000 sq.ft.) unless turfgrass injury can be tolerated. To minimize turfgrass injury, additional applications should be made at least four weeks apart. Avoid swath overlaps. ^{††} Do not apply this product to Bermudagrass on sod farms.	

APPLICATION RESTRICTIONS

- The sale and use of this product in Nassau and Suffolk counties in the state of New York is strictly prohibited.
- In the states of California, Oregon and Washington, turfgrass and lawn uses are restricted to golf courses only.
- The use of this product in the spring when warm season turfgrass is breaking dormancy may significantly delay green up time for turfgrass.
- For ground application only.
- Do not use on residential turf. Turfgrass and lawn uses are restricted to non-residential sites.
- Do not use this product on golf course putting greens or tee areas.
- Do not apply to Bermudagrass on sod farms.
- Do not apply this product in a manner that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area being treated during application.
- This product can affect susceptible broadleaf plants directly through foliage and indirectly by root uptake from treated soil. As a result, do not apply this product directly to, or allow spray drift to come in contact with desirable broadleaf crops and ornamental plants or soil where sensitive crops will be planted the same season, such as: alfalfa, beans, flowers, grapes, lentils, peas, potatoes, safflower, soybeans, sunflowers and tomatoes.
- Do not apply to exposed roots of shallow rooted trees and shrubs.
- Do not allow sprays of this product to contact exposed suckers and/or roots of trees and shrubs or injury may occur.
- Do not reseed for three weeks after application.
- Do not collect grass clippings applied with this product for mulch or compost nor send applied grass clippings to a compost facility. Applicator must give notice to landowners/property managers to not use grass clippings for composting.
- Do not apply on ditches used to transport irrigation water.
- Do not apply this product through any type of irrigation system.
- Do not contaminate irrigation ditches or water used for irrigation or domestic purposes.
- Do not apply where runoff or irrigation water may flow onto susceptible crops resulting in injury to crops.

TREATMENT OF TURFGRASS SPECIES NOT LISTED ON LABEL

Users who wish to use this product on a turfgrass species that is not listed on this label may determine the suitability for such use by treating a small area at a recommended rate. Before application treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine whether or not the treatment is safe to the targeted turfgrass species.

The user assumes full responsibility for plant damage or other liability resulting from using this product on turfgrass species which are not recommended on this label.

APPLICATION DIRECTIONS

Add one-half the desired amount of clean water to spray tank. Add product to spray tank and complete addition of water with agitation running. Mix thoroughly and continue agitation while spraying.

Apply treatment spray using equipment that will insure uniform coverage (see specific application directions below). Sprays should be applied when weeds are actively growing. Application under drought conditions may provide less than desirable results. Broadleaf weed species germinate at different times. Only emerged weeds present at time of application are controlled.

Apply 1 to 2 pints of product per acre to control broadleaf weeds. A maximum of 0.19 lb. ae of clopyralid / 10.56 lbs. ae of triclopyr per acre (2 pints of product per acre) per application is recommended. To minimize turfgrass injury, repeat applications, if necessary, should be made not less than 4 weeks apart. Newly seeded turf should be mowed 2 or 3 times before treating. Do not water for 6 hours after application.

Additional Turfgrass Application Restrictions:

- Do not use more than 0.38 lb. ae of clopyralid / 1.125 lb. ae of triclopyr per acre (4 pints of product per acre) per year of treatment.
- In Florida and New York, the maximum use rate is 0.25 lb. ae of clopyralid / 0.75 lb. ae of triclopyr per acre (2-2/3 pints of product per acre) per growing season.

Avoid overlapping of the spray pattern which could result in higher than recommended application rates. Rates above those recommended on this label could result in turf injury.

SPRAY DRIFT MANAGEMENT

Apply this product in such a manner to avoid contact to nearby susceptible crops or other desirable plants. Applications should be made only when hazards from spray drift are at a minimum. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants including ornamental trees and shrubs. Do not spray when winds will carry spray mist toward susceptible crops or ornamental plants.

GROUND TURFGRASS APPLICATION

Spray drift can be reduced by:

- keeping the spray boom as low as possible
- applying no less than 20 gallons of spray per acre (except under Low Volume Application)
- keeping the operating spray pressures at the manufacturer's minimum recommended pressures for the specific nozzle type used
- spraying when the wind velocity is low (follow state regulations)

Avoid application under completely calm conditions which may be conducive to air inversion.

In hand-gun applications, select the minimum pressure required to obtain adequate plant coverage without forming a mist.

Do not apply with a mist blower.

STANDARD BROADCAST TURFGRASS APPLICATION

Apply 1 to 2 pints of this product in enough water to deliver 20 to 200 gallons of total spray mix per acre (0.5 to 5.0 gallons spray per 1000 sq. ft.). Higher application volumes may be used when this product is tank mixed with fertilizers.

LOW VOLUME TURFGRASS APPLICATION

Apply 1 to 2 pints of this product in enough water to deliver from 5 to 20 gallons of total spray mixture per acre (0.125 to 0.5 gallon spray per 1000 sq. ft.).

Use low pressures and application equipment capable of delivering a uniform droplet size that can wet the weed leaf surface.

To improve spray coverage, the addition of a non-ionic surfactant at a rate of 0.25 to 0.5 pint per acre is suggested.

Use the higher rates of surfactant for lower rates of product and lower spray volumes.

The use of ULV applications is not recommended.

SPOT TREATMENT OF ORNAMENTAL TURFGRASS USING PORTABLE SPRAYERS

Mix 0.5 fl. oz. of this product in enough water to make 1 gallon of spray and apply at any time broadleaf weeds are susceptible by wetting foliage of undesirable plants to point of runoff. This is enough spray to treat approximately 1000 sq. ft. of turf.

WEEDS CONTROLLED AND USE RATE RECOMMENDATIONS

Use the higher rates when hard to control species are prevalent, when applications are made in late summer on mature weeds, and during periods of drought stress.

WEEDS	SUGGESTED USE RATE		
	pint / acre	fl.oz. / 1000 sq.ft.	tsp / 1000 sq.ft.
black medic hop clover red clover white clover	1.0 [†]	.037 (11 ml)	2.5
American burnweed common chickweed common cocklebur common vetch creeping beggarweed dwarf beggarweed false dandelion hawkweed henbit matchweed mouse ear chickweed round leaf mallow sheep sorrel spotted catsear spurweed	1.5	0.55 (16 ml)	3.5
broadleaf plantain burdock coffeeweed common dandelion common ragweed lambsquarters narrowleaf plantain (buckhorn) shepherd's purse Virginia pepperweed	1.5 - 2.0	0.55 - 0.74 (16 - 22 ml)	3.5 - 4.5
Canada thistle common yellow woodsorrel ^{†††} creeping woodsorrel ^{†††} curly dock English lawn daisy ^{††} goldenrod lespedeza musk thistle poison ivy smartweed wild buckwheat wild violet ^{†††}	2.0	0.74 (22 ml)	4.5
[†] For faster activity under good growing conditions. 1.5 pints per acre is recommended. Repeat applications for extended weed control are recommended. ^{††} Six to eight weeks required for control. ^{†††} Repeat treatment may be necessary.			

CROP ROTATION PRECAUTIONS

Residues of this product in treated plant tissues, including the treated crop or weeds, which have not completely decayed may affect succeeding susceptible crops.

FIELD BIOASSAY INSTRUCTIONS

In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample field conditions such as soil texture, soil pH, drainage, and any other variable that could affect the seed bed of the new crop. The field bioassay can be initiated at any time between harvest of the treated crop and the planting of the intended rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the test rotational crop; plant only a labeled crop or crop listed in the table below for which the rotational interval has clearly been met.

Crop Rotation Intervals for ALL STATES EXCEPT Idaho, Nevada, Oregon, Utah and Washington

Rotation Crops (1)	Rotation Interval † (Soils > 2% organic matter AND Rainfall > 15 inches DURING 12 months following application.)	Rotation Interval † (Soils < 2% organic matter AND Rainfall < 15 inches DURING 12 months following application.)
barley, field corn, grasses, oats, wheat	30 days	30 days
canola (rapeseed), flax, sugar beets	5 months	5 months
alfalfa, asparagus, cole crops, dry beans, grain sorghum, mint, onions, popcorn, safflower, soybeans, strawberries, sunflowers, sweet corn	10.5 months	10.5 months
lentils, peas, potatoes (incl. potatoes grown for seed) and broadleaf crops grown for seed (excluding <i>Brassica</i> species)	18 months (2, 3)	18 months (2, 3)
(1) A field bioassay is recommended prior to planting any broadleaf crops that are not listed. Do not rotate to unlisted crops prior to 10.5 months following application. (2) An 18-month crop rotation is recommended due to potential crop injury. Note: For these crops, a minimum 10.5-month rotation interval must be observed to avoid illegal residues in the harvested crop. (3) The potential for injury may be reduced by burning, removal, or incorporation of treated crop residues followed by a minimum of two supplemental fall irrigations.		
† Note: The above intervals are based upon average annual precipitation, regardless of irrigation practices. Observance of recommended crop rotation intervals should result in adequate safety to rotational crops. However, this product is dissipated in the soil by microbial activity and the rate of microbial activity is dependent upon several interrelating factors including soil moisture, temperature and organic matter. Therefore, accurate prediction of rotational crop safety is not possible. In areas of low organic matter (<2%) and less than 15 inches average annual precipitation, potential for crop injury may be reduced by burning or removing plant residues, supplemental fall irrigation and deep moldboard plowing prior to planting the sensitive crop.		

Crop Rotation Intervals for Idaho, Nevada, Oregon, Utah and Washington ONLY

Rotation Crops (1)	Rotation Interval † (Soils > 2% organic matter AND Rainfall > 15 inches DURING 12 months following application.)	Rotation Interval † (Soils < 2% organic matter AND Rainfall < 15 inches DURING 12 months following application.)
barley, field corn, grasses, oats, wheat	30 days	30 days
canola (rapeseed), flax, sugar beets	5 months	5 months
asparagus, <i>Brassica</i> species grown for seed, cole crops, grain sorghum, mint, onions, popcorn, strawberries, sweet corn	12 months	12 months
alfalfa, dry beans, soybeans, sunflowers	12 months (2)	18 months (2)
lentils, peas, potatoes (incl. potatoes grown for seed) and broadleaf crops grown for seed (excluding <i>Brassica</i> species)	18 months (2)	18 months (2, 3)

- (1) A field bioassay is recommended prior to planting any broadleaf crops that are not listed. Do not rotate to unlisted crops prior to 12 months following application.
- (2) An 18-month crop rotation is recommended due to potential crop injury.
Note: For these crops, a minimum 12-month rotation interval must be observed to avoid illegal residues in the harvested crop.
- (3) Crop injury and/or yield loss may occur up to 4 years after application. A field bioassay is also recommended prior to planting these sensitive crops. See instructions above.

† ~~Note: The above intervals are based upon average annual precipitation, regardless of irrigation practices. Observance of recommended crop rotation intervals should result in adequate safety to rotational crops. However, this product is dissipated in the soil by microbial activity and the rate of microbial activity is dependent upon several interrelating factors including soil moisture, temperature and organic matter. Therefore, accurate prediction of rotational crop safety is not possible. In areas of low organic matter (<2%) and less than 15 inches average annual precipitation, potential for crop injury may be reduced by burning or removing plant residue, supplemental fall irrigation and deep moldboard plowing prior to planting the sensitive crop.~~

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Plastic Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Metal Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINER: Close all openings which have been opened during use and replace all caps. For returns of empty container, contact Nufarm Americas, Inc. Customer Service Department at 1-800-345-3330 to obtain proper handling instructions.

GENERAL: Consult Federal, State or local disposal authorities for approved alternative procedures.

11/11

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(NEW)

Brazen is a trademark of Nufarm Americas Inc.