

228-572

9/22/2008

1/38

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460



SEP 22 2008

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Mr. Matthew Granahan  
Nufarm Americas, Inc.  
150 Harvester Drive, Suite 200  
Burr Ridge, IL 60527

Dear Mr. Granahan:

Subject: Amendment of label to revise crop grouping 10 for white sapote to clarify the  
brassica vegetable group  
Nuprid 2SC Soil/Foliar Insecticide  
EPA Registration No. 228-572  
Your Submissions Dated 6/17/08

The labeling referred to above submitted in connection with the Federal Insecticide,  
Fungicide and Rodenticide, as amended is acceptable.

A stamped copy of the labeling is enclosed for your records. Please submit one final  
printed copy of the labeling before releasing the product for shipment. If you have any questions  
regarding this label, please contact Autumn Metzger at (703) 305-5314.

Sincerely,

A handwritten signature in black ink, appearing to read "Venus Eagle", written over a horizontal line.

Venus Eagle  
Product Manager (1)  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

2/38

GROUP 4A INSECTICIDE

# NUPRID® 2SC

## SOIL/FOLIAR INSECTICIDE

FOR USES IN PEST MANAGEMENT AND SUPPRESSION OF INSECT  
VECTORED DISEASES AND MAINTENANCE OF PLANT HEALTH.

**ACTIVE INGREDIENT:**

Imidacloprid, 1[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine. . . . . 21.4%

**OTHER INGREDIENTS:** . . . . . 78.6%

**TOTAL:** . . . . . 100.0%

Contains 2 pounds of imidacloprid per gallon.

SHAKE WELL BEFORE USING

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION – PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-572  
EPA EST. NO.

MANUFACTURED FOR  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



NET CONTENTS      GALS.

000228-00572.20080617.EPA.N  
NUP-07387

**ACCEPTED**

SEP 22 2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
Registered under  
EPA Reg. No. 228-572

3/38

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION - PRECAUCION**

Harmful if absorbed through skin. Harmful if inhaled. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as Natural Rubber, Selection Category A).

<b>FIRST AID</b>	
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move the person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
No specific antidote is available. Treat the patient symptomatically.	

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and Other Handlers Must Wear:**

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS STATEMENTS**

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.
- Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.**

4/38

## SPRAY DRIFT MANAGEMENT

The responsibility of avoiding spray drift is with the applicator. The applicator should consider weather related factors and the interaction of application equipment when making application decisions.

### Mixing and Loading Requirements

The use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading area and potential surface to groundwater conduits such as field sumps, uncased well heads, sink-holes, or field drains.

### Aerial Applications

The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

### Importance of Droplet Size

The droplet size is an important factor and can influence drift. Typically smaller droplet sizes, such as less than 150 to 200 microns, have a greater tendency to drift compared to larger droplets. Applications typically should be made to deliver the largest droplet range that provides adequate control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection.

### Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions.

### Restrictions During Temperature Inversions

Do not make ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions typically restrict vertical air mixing, which then could cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions typically are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

### Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward-pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

### No-Spray Zone Requirements for Soil Applications

Do not apply within 25 feet of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

### Runoff Management

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When using this product on erodible soils, Best Management Practices for minimizing runoff should be employed.

### Endangered Species Notice

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

### Resistance Management

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

This product contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

The active ingredient in this product belongs to the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to this product. To maintain susceptibility to this class of chemistry in insect species with high resistance development potential, it is recommended that for each crop season: 1) only a single, soil application of this product be made; 2) foliar applications of products from this same class not be made following a long residual, soil application of this product, or other neonicotinoid products.

Foliar applications of this product or other Group 4A products from the neonicotinoid chemical class should not be used on crops previously treated with a long-residual, soil-applied product from the neonicotinoid chemical class.

Examples of other Group 4A, neonicotinoid products used as foliar treatments include: Actara, Assail, Calypso, Centric, Clutch, Couraze, Gallant, Impulse, Intruder, Leverage, Passada, Provado, Trimax, Trimax Pro and Venom.

5/38

Examples of other Group 4A, neonicotinoid products used as soil treatments include: *Admire, Admire Pro, Advise, Alias, Couraze, Cruiser, Gaucho, Macho, Macho Max, Platinum, Venom and Widow.*

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

### APPLICATION INSTRUCTIONS

#### For Foliar Applications

Apply as a directed or broadcast foliar spray. Thorough coverage of foliage is necessary without runoff for optimum insecticidal efficacy. Use adequate spray volumes, properly calibrated application equipment and spray adjuvant if necessary to obtain thorough coverage. Failure to provide adequate coverage and retention of this product on leaves and fruit may result in loss of insect control or delay in onset of activity. This product may be applied with properly calibrated ground or aerial application equipment. Minimum recommended spray volumes, unless otherwise specified on crop specific application instructions sections, are 10 gallons/Acre by ground applications and 5 gallons/Acre through aerial equipment. This product may also be applied by overhead chemigation (see additional CHEMIGATION DIRECTIONS FOR USE section below) if allowed in crop specific application instruction section.

#### For Soil Applications

Applications of this product should apply active ingredient into the seed or root-zone of crop. Failure to place this product into root-zone may result in loss of control or delay in onset of activity. This product may be applied with ground or chemigation application. Do not apply with aerial application equipment. Broadcast, foliar applications are only recommended to seedling flats or trays, or where product is intended to be washed from foliage to soil prior to drying on foliage.

Optimum activity results from applications to the root-zone of plants to be protected. The earlier this product is available to a developing plant, the earlier the protection begins. This product is continuously taken into the roots over a long period of time and the systemic nature of this product allows movement from roots through the xylem tissue to all vegetative parts of the plant. This results in extended residual activity, the control of insects and the prevention and/or reduction of virus transmission or symptom expression, and plant health benefits. The rate applied affects the length of the plant protection period. Higher rates are recommended when infestations occur later in crop development, or where pest pressure is continuous. This product will generally not control insects infesting flowers, blooms or fruit. Additional crop protection may be required for insects feeding in or on these plant parts and for insects not listed in the crop-specific, pests controlled sections of this label.

Suppression, or less than complete control of certain diseases and insect pests including reduced feeding, may also result from applications of this product. Complete control of these pests/diseases may require supplemental control measures.

Use of this product on crops grown for production of true seed intended for private or commercial planting is generally not recommended but may be allowed under State specific supplemental labeling. As with any insecticide, care should be taken to minimize exposure of this product to honey bees and other pollinators. Use of this product on crops requiring bee pollination should be avoided during bloom and a minimum of 10 days prior to bloom. Additional information on this product uses for these crops and other questions may be obtained from the Cooperative Extension Service, PCAs, or local Nufarm representative.

Pre-mix with water or other appropriate diluent prior to application. Keep this product and water suspension agitated to avoid settling.

Do not apply more than 0.5 lbs active ingredient per acre, per crop season, regardless of formulation or method of application, unless specified within a crop-specific, Application Instructions section for a given crop.

**Mixing Instructions**

To prepare the application mixture, add a portion of the required amount of water to the tank and with agitation then add this product. Complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application. This product may also be used with other pesticides and/or fertilizer solutions. **Please see Compatibility Note below.** When tank mixtures of this product and other pesticides are involved, prepare the tank mixture as recommended above and follow suggested Mixing Order below.

**Mixing Order**

When pesticide mixtures are needed, add wettable powders first, then this product and other flowable (suspension concentrate) products second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add an additional component until the previous is thoroughly mixed. If a fertilizer solution is added, a fertilizer/pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

**Compatibility Note**

Test compatibility of the intended mixture before adding this product to the spray or mix tank. Add proportionate amounts of each ingredient in the appropriate order, to a pint or quart jar, cap, shake for 5 minutes, and let set for 5 minutes. Poor mixing or formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used. For further information, contact your local Nufarm representative.

**CHEMIGATION - DIRECTIONS FOR USE**

Refer to DIRECTIONS FOR USE section before proceeding with chemigation application.

**Types of Irrigation Systems****For Soil Application**

Chemigation applications of this product may only be made to crops through chemigation systems as specified in crop-specific application sections and only through low-pressure systems unless specifically recommended for a given crop. Do not apply this product through any other type of irrigation system.

**For Foliar Application**

Chemigation applications of this product may be made to crops through overhead sprinkler chemigation systems if specified in crop-specific instruction sections. Do not apply this product through any other type of irrigation system.

**Water Volume**

Chemigation applications of this product should be made as concentrated as possible. Retention of this product on target site of insect infestation is necessary for optimum activity. Chemigation of this product in water volumes exceeding 0.10 inch/Acre is not recommended.

**Uniform Water Distribution and System Calibration**

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

**Chemigation Monitoring**

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Drift**

Do not apply when wind speed favors drift beyond the area intended for treatment.

**Required System Safety Devices**

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**Using Water from Public Water Systems**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally-closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must

contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

ROTATIONAL CROPS*	
Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.	
<b>IMMEDIATE PLANT-BACK:</b> All crops on this label plus the following crops not on this label: barley, canola, corn (field, pop & sweet), rapeseed, sorghum, sugar beet and wheat.	
<b>30-DAY PLANT-BACK:</b> Cereals (including buckwheat, millet, oats, rice, rye, and triticale), soybeans and safflower	
<b>12-MONTH PLANT-BACK:</b> All Other Crops	
<b>10-MONTH PLANT-BACK:</b> Onion and bulb vegetables	
* Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.	

## FIELD CROPS APPLICATION RATES

### COTTON - SOIL

Pests Controlled	Rate fluid ounces/1,000 row-feet	Rate fluid ounces/Acre
Cotton aphid Plant bugs Thrips Whiteflies	1.3	17.0 to 21.1 (Depending on row-spacing)
<b>Notes and Restrictions</b> Maximum amount allowed per crop season: 21.1 fluid ounces/Acre (0.33 lb AI/Acre)		
<b>Applications</b> Apply specified dosage in one of the following methods: 1. In-furrow spray during planting directed on or below seed; 2. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; 3. Chemigation into root zone through low-pressure drip or trickle irrigation.		

Regardless of formulation or method of application, apply no more than 0.5 lb active ingredient per acre per season, including seed treatment as Gaucho®, soil and foliar uses. Do not apply more than a total of 6 applications of the active ingredient per season. Do not graze treated fields after any application of this product. Please see Resistance Management section of this label.

8/38

# **COTTON - FOLIAR**

Pests Controlled		Rate fluid ounces/Acre
Aphids Cotton Fleahoppers Plant bugs (excludes <i>Lygus hesperus</i> ) Bandedwinged whitefly Green stink bug Southern green stink bug Bollworm/Budworm (ovicidal effect)		2.0 to 4.0
Pests Suppressed		
Lygus bugs ( <i>Lygus hesperus</i> ) Whiteflies (other than bandedwinged whitefly)		3.0 to 4.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 7 days Maximum amount allowed per season: 20.0 fluid ounces/Acre (0.31 lb. AI/A) Do not graze treated fields after any application of this product.		
<b>Applications</b> This product may be applied through properly calibrated ground, aerial or chemigation application equipment.		
Tank Mix Recommendations		
Pests Controlled (In addition to pests listed above)	This Product Rate fluid ounces/Acre	Bidrin® 8* Rate fluid ounces/Acre
<b>For early season control of:</b> Thrips	2.0 to 3.0	1.6 to 3.2
<b>For mid to late season control of:</b> Plant bugs Stink bugs (including Brown stink bug) Grasshoppers Saltmarsh caterpillar Cotton leafperforator	2.0 to 3.0	4.0 to 8.0
<b>Notes and Restrictions (In addition to Notes and Restrictions listed above)</b> *Refer to the Bidrin® 8 product label for specific use recommendations; observe all restrictions and precautions that appear on the label.		



# POTATO - SOIL

Pests Controlled	Rate fluid ounces/1,000 row-feet	Rate fluid ounces/Acre
Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid	0.9 to 1.3	13.0 to 20.0
<b>Pests / Diseases Suppressed</b>		
Symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis (PLRV) Wireworms (with in-furrow spray at planting)	0.9 to 1.3	13.0 to 20.0
<b>Notes and Restrictions:</b> Maximum amount allowed per crop season: <b>20.0 fluid ounces/Acre</b> (0.31 lb AI/Acre) <b>Applications:</b> Apply specified dosage in one of the following methods: 1. In-furrow spray during planting directed on seed pieces or seed potatoes; 2. Subsurface side-dress on both sides of the row covered with 3 or more inches of soil; 3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil; 4. Narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting. For effective pest control or suppression, applications of this product must be placed below soil surface and in contact with seed piece or within root zone. For potatoes grown on highly permeable soils with shallow water table, at-plant applications of this product may be made in a 2 to 4 inch band (width of planter shoe opening) and completely covered.		

## POTATO \* - Seed Piece Treatment

Pests Controlled	Rate fluid ounces/100 lbs seed	Rate fluid ounces/Acre
Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid Wireworms (seed piece protection)	0.4 to 0.8	8.0 to 16.0
<b>Pests / Diseases Suppressed</b>		
Symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis (PLRV)	0.8	16.0
<b>Notes and Restrictions</b> Maximum amount allowed per crop season: <b>20.0 fluid ounces/Acre</b> (0.31 lb AI/Acre) Do not use treated seed pieces for food, feed, or fodder. Do not apply any subsequent application of this product (in-furrow), Admire®, Gaucho®, Leverage® or Provado® following a seed-piece treatment of this product. <b>Applications</b> Apply specified dosage as a diluted spray onto seed pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part this product. Agitate or stir spray solution as needed. Fungicidal or inert absorbent dusts may be applied after this product's application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed pieces as soon as possible after treating avoiding prolonged exposure of seed pieces treated with this product to sunlight and in accordance with the recommendation of your local Extension specialist. Consult your local Nufarm representative or crop protection product dealer for information relevant to your area. * Based on a seeding rate of 2000 lbs/acre.		

POTATO - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Colorado potato beetle Flea beetles Fleahoppers Psyllids	3.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 7 days Maximum amount allowed per season: 12.8 fluid ounces/Acre (0.2 lb. AI/A)	

TOBACCO - SOIL

Pests Controlled	Rate fluid ounces/1,000 plants (as seedling tray drench)	Rate fluid ounces/1,000 plants (in-furrow or transplant-water)
Aphids Flea beetles	1.0	1.4
Mole crickets Whiteflies Wireworms	1.4 to 2.8	1.8 to 2.8
<b>Pests / Diseases Suppressed</b>		
Cutworms Symptoms of: Tomato spotted wilt virus (TSWV)	1.4 to 2.8	1.8 to 2.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per crop season: 32.0 fluid ounces/Acre (0.50 lb AI/Acre) <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Uniform, broadcast foliar spray to seedlings in trays (tray drench) not more than 7 days prior to transplanting, followed immediately by overhead irrigation to wash this product from foliage into potting media. Failure to wash this product from foliage may result in a reduction in pest control. Transplants should be handled carefully during setting to avoid dislodging treated potting media from roots. 2. In-furrow spray or transplant-water drench during setting. 3. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. <b>Important Note:</b> Proper tray drench applications of this product have been shown to be the most efficacious method of application. However, the specified rate of this product may be applied as a combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of this product into the plant and a delay in control.		

TOBACCO - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids	1.6 to 3.2
Flea beetles Japanese beetle	3.2
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 7 days Maximum amount allowed per season: 18.0 fluid ounces/Acre (0.28 lb. AI/A)	

## VEGETABLE and SMALL FRUIT CROPS APPLICATION INSTRUCTIONS

### For Foliar Applications

Apply specified rate per acre as a broadcast or directed foliar spray as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimal control. A spray adjuvant may be used to improve coverage. This product may not knock down established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. This product may be tank mixed with other insecticides for knockdown of pests or for improved control of other pests. Crops contained with certain Crop Groups recognized by USEPA are subject of change. Please refer to USEPA website ([www.epa.gov](http://www.epa.gov)) for latest Crop Groups.

### CUCURBIT VEGETABLES - SOIL <sup>1/</sup>

**Crops of Crop Group 9 including:** Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

Field application rates. See details on next page for additional planthouse instructions.	
Pests Controlled	Rate fluid ounces/Acre
Aphids Cucumber beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	16.0 to 24.0
Pests / Diseases Suppressed	
Bacterial wilt (as vectored by various cucumber beetles) Leaf silvering resulting from whitefly feeding	16.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 21 days Maximum amount allowed per application: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre)	
<b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.	
Planthouse Application Instructions	
Pests Controlled	Rate fluid ounces/1,000 plants
Aphids Whiteflies	0.1
<b>Notes and Restrictions</b> Maximum amount applied in the planthouse: <b>0.1 fluid ounce</b> (0.00156 lb AI)/1,000 plants. Maximum number of applications in planthouse: 1	
<b>Applications</b> Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners: 1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash this product from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash this product from foliage may result in reduced pest control;	
(continued)	

<b>Planthouse Application Instructions - (Continued)</b>
<b>Applications - continued</b>
2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.
The application made in the planthouse is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants should be handled carefully during setting to avoid dislodging treated potting media from roots.
<b>Important Note:</b> Not all varieties of cucurbit vegetables have been tested for tolerance to this product applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

**GREENHOUSE VEGETABLES - SOIL <sup>1/</sup>**

**Cucumber and Tomato ONLY** (Mature plants in production greenhouses)

<b>Pests Controlled</b>	<b>Rate fluid ounces/1,000 plants</b>
Aphids Whiteflies	1.4
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>0 days</b> Maximum number of applications per crop season: 1 Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Do not apply to immature plants since phytotoxicity may occur. Applications should be made when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials ( <i>Orius</i> sp.) can occur when this product is applied. Many varieties of vegetables have been tested for tolerance to this product and show good safety. However, certain varieties may show more sensitivity to this product. Therefore, treat a few plants before treating the whole greenhouse.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

# FRUITING VEGETABLES - SOIL <sup>1/</sup>

Crops of Crop Group 8 plus Okra including: Eggplant, Ground cherry, Okra, Pepper (including bell, chilli, cooking, pimento and sweet) Tomato, Pepinos, Tomatillo

Field application instructions. See details that follow for additional planthouse instructions.	
Pests Controlled	Rate fluid ounces/Acre
Aphids Colorado potato beetle Flea beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	Okra and Pepper 16.0 to 32.0  Other Crops 16.0 to 24.0
Pests / Diseases Suppressed	
Symptoms of: Tomato mottle virus Tomato spotted wilt virus Tomato yellow leaf curl virus	Okra and Pepper 16.0 to 32.0 Other Crops 16.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 21 days  Maximum amount allowed on pepper and okra crops per application: <b>32.0 fluid ounces/Acre</b> (0.5 lb AI/Acre) Maximum amount allowed on other fruiting vegetable crops per application: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre)  <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.	
Planthouse Application Instructions	
Pests Controlled	Rate fluid ounces/1,000 plants
Aphids Whiteflies	0.1
<b>Notes and Restrictions</b> Maximum amount applied in the planthouse: <b>0.1 fluid ounce</b> (0.00156 lb AI)/1,000 plants. Maximum number of applications in planthouse: 1  <b>Applications</b> Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners: 1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash this product from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash this product from foliage may result in reduced pest control; 2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.  The application made in the planthouse is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants should be handled carefully during setting to avoid dislodging treated potting media from roots.  <b>Important Note:</b> Not all varieties of fruiting vegetables have been tested for tolerance to this product applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

14/38

# FRUITING VEGETABLES - FOLIAR <sup>1/</sup>

Crops of Crop Group 8 plus Okra Including: Eggplant, Ground cherry, Okra, Pepper (including bell, chill, cooking, pimento and sweet), Tomato, Pepinos, Tomatillo

Pests Controlled	Rate fluid ounces/Acre
Aphids Colorado potato beetle Leafhoppers Whiteflies	3.0 to 5.0
Pepper weevil	5.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 0 days Minimum interval between applications: 5 days Maximum amount allowed per season: 15.4 fluid ounces/Acre (0.24 lb. AI/A) <b>Applications</b> For pepper weevil, apply specified dosage of this product by ground equipment only, timing applications prior to a damaging pest population becoming established. Good coverage of foliage and fruit is necessary for optimal control. Applications of this product must be incorporated into a full-season program, where alternations of effective products from multiple classes of chemistry and different modes of actions are utilized in a blocked or windowed approach. For additional information, please contact your Nufarm representative, Extension Specialist or crop advisor. Higher rate should be used when targeting adult whiteflies.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

## GLOBE ARTICHOKE - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers	3.2 to 8.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 14 days Maximum amount allowed per season: 32.0 fluid ounces/Acre (0.50 lb. AI/A)	

**HERBS - SOIL**

**Crops of Crop Subgroup 19A including:** Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Burnet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea Beetles Leafhoppers Whiteflies	16.0 to 24.0
<b>Pests Suppressed</b>	
Thrips (foliage-feeding thrips only)	16.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per season: 24.0 fluid ounces/Acre (0.38 lb AI/Acre) <b>Applications</b> Apply specified dosage in one of the following methods: 1. In-furrow spray during planting directed on or below seed; 2. In-furrow spray or transplant-water drench during setting or transplanting; 3. Shanked-into or below eventual seed-line; 4. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. <b>Notes</b> Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Nufarm strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use.	

**HERBS - FOLIAR**

**Crops of Crop Subgroup 19A including:** Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Burnet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea Beetles Leafhoppers Whiteflies	2.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum amount allowed per crop season: 8.4 fluid ounces/Acre (0.13 lb. AI/Acre) <b>Applications</b> This product may be applied through properly calibrated ground and aerial application equipment. Thorough coverage with direct contact of the spray material to the target pests is required for optimum control. The addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage and control. <b>Notes</b> Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Nufarm strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use.	

# **HEAD and STEM BRASSICA VEGETABLES and LEAF BRASSICA GREENS, plus TURNIP TOPS - SOIL**

**Crop of Crop Group 5 including:** Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (*gai lon*) broccoli, Chinese (*bok choy*) cabbage, Chinese (*napa*) cabbage, Chinese mustard (*gai choy*) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, plus Turnip tops (leaves)

## **LEAFY VEGETABLES - SOIL <sup>1/</sup>**

**Crops of Crop Subgroup 4A plus Watercress including:** Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Rocket), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Raddicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only, applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

Pests Controlled	Rate fluid ounces/Acre (on 36 inch rows)
Aphids Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	10.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 21 days Maximum amount allowed per application: 24.0 fluid ounces/Acre (0.38 lb AI/Acre) <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

## **HEAD and STEM BRASSICA VEGETABLES and LEAF BRASSICA GREENS - FOLIAR**

**Crops of Crop Group 5 including:** Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (*gai lon*) broccoli, Chinese (*bok choy*) cabbage, Chinese (*napa*) cabbage, Chinese mustard (*gai choy*) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach and Rape greens



**LEAFY GREEN VEGETABLES - FOLIAR <sup>1/</sup>**

**Crops of Crop Subgroup 4A plus Watercress including:** Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (rocket), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach), Watercress (commercial production only; applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	3.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>5 days</b> Maximum amount allowed per season: <b>15.4 fluid ounces/Acre</b> (0.24 lb. AI/A) <b>Applications</b> For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following application. Applications must be made to fully leafed-up canopies only.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

**LEAFY PETIOLE VEGETABLES - SOIL <sup>1/</sup>**

**Crops of Crop Subgroup 4B including:** Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, finocchio), Rhubarb, Swiss chard

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	10.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>45 days</b> Maximum amount allowed per application: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre) <b>Applications</b> Apply specified dosage of this product in one of the following methods: <ol style="list-style-type: none"> <li>1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;</li> <li>2. In-furrow spray directed on or below seed;</li> <li>3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application;</li> <li>4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;</li> <li>5. Post-seeding drench, transplant-water drench, or hill drench;</li> <li>6. Subsurface side-dress on both sides of each row. This product must be incorporated into root zone.</li> </ol>	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

## LEGUME VEGETABLES - SOIL <sup>1/</sup>

**Crops of Crop Group 6 ( Except soybean, dry ) including:** Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean, Bean (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean), Pea (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea), Other Beans and Peas [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	16.0 to 24.0
<b>Pests / Diseases Suppressed</b>	
Symptoms of: Bean common mosaic virus (BCMV) Bean golden mosaic virus (BGMV) Beet curly top/hybrigeminivirus (BCTV)	16.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>21 days</b> Maximum amount allowed per crop season: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre) <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray at planting directed on or below seed; 3. In a narrow (2" or less) surface band over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours following application; 4. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; 5. As a post-seeding drench, transplant drench, or hill drench.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

## LEGUME VEGETABLES - FOLIAR <sup>1/</sup>

**Crops of Crop Group 6 ( Except soybean, dry ) including:** Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean, Bean (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean), Pea (*Pisum* spp. includes dwarf pea, edible pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea), Other Beans and Peas [Broad bean (fava), chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), lentil, Pigeon pea, soybean (immature seed), Sword bean]

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Whiteflies	2.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>7 days</b> Maximum amount allowed per season: <b>8.4 fluid ounces/Acre</b> (0.13 lb. AI/A)	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

**ROOT VEGETABLES - SOIL <sup>1/</sup>**

Crops of Crop Subgroup 1B except Sugarbeet including: Beet (garden)<sup>2/</sup>, Burdock (edible)<sup>2/</sup>, Carrot<sup>2/</sup>, Celeriac<sup>2/</sup>, Chervil (turnip-rooted)<sup>2/</sup>, Chicory<sup>2/</sup>, Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip<sup>2/</sup>, Radish<sup>2/</sup>, Oriental radish (daikon)<sup>2/</sup>, Rutabaga<sup>2/</sup>, Salsify (oyster plant), Salsify (black)<sup>2/</sup>, Salsify (Spanish), Skirret and Turnip<sup>2/</sup>.

Pests Controlled	Rate fluid ounces/1,000 row-feet	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	0.7 to 1.7	10.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>21 days</b> Maximum amount allowed per crop season: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre) Maximum number of applications per crop season: <b>1</b> <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray (rate specified per 1,000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting; 3. In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting. <b>Important Note:</b> The rate applied affects the length of control. Use higher rates where infestations occur later in crop development, or where pest pressure is continuous. Rates of this product less than 0.7 fluid ounce/1,000 row-feet will not provide adequate residual pest control. Crops treated with this product grown on very high organic matter soils (muck) may also require additional pest management control.		

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

<sup>2/</sup> Tops or greens from these crops may be utilized for food or feed.

**TUBEROUS and CORM VEGETABLE - SOIL <sup>1/</sup>**

Crops of Crop Subgroup 1C including: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter & sweet)<sup>2/</sup>, Chayote (root), Chufa, Dasheen (taro)<sup>2/</sup>, Ginger, Leren, Sweetpotato, Tanier (cocoyam)<sup>2/</sup>, Turmeric, Yam bean (jicama, manioc pea), Yam (true)<sup>2/</sup>

For applications on potato see Field Crops section for Potato - Soil

Pests Controlled	Rate fluid ounces/1,000 row-feet	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	0.7 to 1.7	10.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI) from planting application: <b>3 days</b> (leaves); <b>125 days</b> (corms) Maximum amount allowed per crop season: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre) Maximum number of applications per crop season: <b>1</b> <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. In-furrow spray (rate specified per 1,000 row-feet) over planting material (hulis) or shanked-in 1 to 2 inches below hulis depth at planting; 2. Side-dress not more than 0.6 fluid ounce/1,000 row-feet no later than 45 days after-planting. Observe same PHI as above. <b>Important Note:</b> The rate applied affects the length of control. Use higher rates where infestations occur later in crop development, or where pest pressure is continuous. Rates of this product less than 0.7 fluid ounce/1,000 row-feet may not provide adequate residual pest control. Crops treated with this product grown on very high organic matter soils (muck) may also require additional pest management control.		

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

<sup>2/</sup> Tops or greens from these crops may be utilized for food or feed.

**ROOT, TUBEROUS and CORM VEGETABLES - FOLIAR <sup>1/</sup>**

**Crops of Crop Group 1 ( Except for sugarbeet ) including:** Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Beet (garden)<sup>2/</sup>, Brudock (edible)<sup>2/</sup>, Canna (edible, Queensland arrowroot), Carrot<sup>2/</sup>, Cassava (bitter & sweet)<sup>2/</sup>, Celery<sup>2/</sup>, Chayote (root), Chervil (turnip-rooted)<sup>2/</sup>, Chicory<sup>2/</sup>, Chufa, Dasheen (taro)<sup>2/</sup>, Ginger, Ginseng, Horseradish, Leren, Parsley (turnip-rooted), Parsnip<sup>2/</sup>, Radish<sup>2/</sup>, Oriental radish (dalkon)<sup>2/</sup>, Rutabaga<sup>2/</sup>, Salsify (black)<sup>2/</sup>, Salsify (oyster plant), Salsify (Spanish), Skirret, Sweet potato, Tanier (cocoyam)<sup>2/</sup>, Turmeric, Turnip<sup>2/</sup>, Yam bean (Jicama, manioc pea), Yam (true)<sup>2/</sup>

For applications on potato see Field Crops section for Potato - Foliar

Pests Controlled	Rate fluid ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	2.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum amount allowed per season: 2.8 fluid ounces/Acre (0.044 lb AI/A) on radish; 8.4 fluid ounces/Acre (0.13 lb. AI/A) on other crops Maximum applications of this product per crop season: 1 on radish; 3 on other crops.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

<sup>2/</sup> Tops or greens from these crops may be utilized for food or feed.

**STRAWBERRY - SOIL <sup>1/</sup>**

Annual and Perennial Crops	
Pests Controlled	Rate fluid ounces/Acre
Aphids Whiteflies	24.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per crop season: 32.0 fluid ounces/Acre (0.50 lb AI/Acre) <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening; 2. As a plant material or plant hole treatment just prior to, or during transplanting. 3. As a band spray over the row in a minimum of 20 gallons of water per acre, followed immediately by overhead irrigation to incorporate product into root zone. Do not use plastic or other mulch that limits movement of this product into root zone. The rate applied affects the length of control. Use higher rates where infestations may occur later in crop development or where pest pressure is continuous.	

21/38

Post-harvest Use on Perennial Crops	
Pests Controlled	Rate fluid ounces/Acre
White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	16.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per crop season: 24.0 fluid ounces/Acre (0.38 lb AI/Acre) <b>Applications</b> Apply a single application <b>post harvest to coincide with renovation of strawberry fields</b> and during active egg-laying period of beetles. Apply specified dosage of this product in one of the following methods: 1. As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre; 2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed; 3. As a chemigation application with 600 to 1,000 gallons of water followed by 0.10 to 0.25 inch irrigation. <b>Important Note:</b> All soil-surface applications must be followed by 0.25 inch of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate this product into egg-deposition zone may result in decreased activity of beetle grubs.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

#### STRAWBERRY - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Spittlebugs Whiteflies	3.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum amount allowed per season: 9.1 fluid ounces/Acre (0.14 lb. AI/A) Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.	

#### SUGARBEET - SOIL <sup>1/</sup> (For Use Only in CA)

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Whiteflies Flea beetles	6.0 to 12.0
Pests / Diseases Suppressed	
Symptoms of: Western yellows / Beet curly top hybrigeminivirus (BCTV)	6.0 to 12.0
<b>Notes and Restrictions</b> Maximum amount allowed per crop season: 12.0 fluid ounces/Acre (0.18 lb AI/Acre) Do not apply immediately prior to bud opening or during bloom or when bees are actively foraging. <b>Applications</b> Apply specified dosage of this product in the following method: Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting. The low rate may be applied to aid establishment of stands in whitelly areas, or for early season control of the other pests listed.	

<sup>1/</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

22/38

NUPRID™ 2SC SOIL/FOLIAR INSECTICIDE CONVERSION CHART FOR LINEAR APPLICATION								
RATE fluid ounces/Acre	RATE fluid ounces/1,000 row-feet Based on average row spacing (In inches):							
	10	15	20	25	30	35	40	45
10	0.19	0.29	0.38	0.48	0.57	0.67	0.76	0.86
12	0.23	0.34	0.46	0.57	0.69	0.80	0.92	1.03
14	0.27	0.40	0.54	0.67	0.80	0.94	1.07	1.21
16	0.31	0.46	0.61	0.77	0.92	1.07	1.22	1.38
18	0.34	0.52	0.69	0.86	1.03	1.21	1.38	1.55
20	0.38	0.57	0.76	0.96	1.15	1.34	1.53	1.72
22	0.42	0.63	0.84	1.05	1.26	1.47	1.68	1.89
24	0.46	0.69	0.92	1.15	1.38	1.61	1.84	2.07
26	0.50	0.75	0.99	1.24	1.49	1.74	1.99	2.24
28	0.54	0.80	1.07	1.34	1.61	1.87	2.14	2.41
30	0.57	0.86	1.15	1.43	1.72	2.01	2.29	2.58
32	0.61	0.92	1.22	1.52	1.84	2.14	2.45	2.75
<b>Important Note:</b> Rate of this product applied affects the length of control and, to a considerable extent, the degree of control or effect. Row-spacing X rate combinations in shaded blocks may not provide adequate residual pest control and are not recommended for long-term, residual control. Use higher labelled rates where infestations may occur later in crop development or where pest pressure is continuous. Nufarm offers no warranty for use of this product at rates below 0.7 fluid ounce/1,000 row-feet.								

23/38

## TREE, BUSH and VINE CROPS

### Application Instructions

#### Application Instructions - For Foliar Applications Only

Apply specified rate per acre as a broadcast or directed foliar spray as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimal control. A spray adjuvant may be used to improve coverage. This product may not knock down established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. This product may be tank mixed with other insecticides for knockdown of pests or for improved control of other pests. Aerial application of this product may result in slower activity and reduced control relative to results from ground application. For trees and vine crops, application rates are based on full size, mature trees or vines.

#### BANANA and PLANTAIN - SOIL

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers	16.0 to 32.0
<b>Pests / Diseases Suppressed</b>	
Scales	16.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 0 day Maximum amount allowed per crop season: 32.0 fluid ounces/Acre (0.5 lb AI/Acre)	
<b>Applications</b> Apply specified dosage of this product in the following method: Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	

#### BANANA and PLANTAIN - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Thrips	6.4
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 0 day Minimum interval between applications: 14 days Maximum amount allowed per crop season: 32.0 fluid ounces/Acre (0.5 lb AI/A)	
<b>Applications</b> Apply specified dosage of this product as a broadcast or directed spray to infested area insuring thorough coverage. This product may be applied through properly calibrated ground or aerial application equipment. Aerial application of this product may result in slower activity and reduced control relative to results from ground application.  Addition of an organosilicone adjuvant at a rate not to exceed 2.0 fluid ounces/100 gallons finished spray solution may improve coverage and pest control.	

## BUSHBERRY - SOIL

Crops of Crop Subgroup 13B including: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Lingonberry, Salal

Pests Controlled	Rate fluid ounces/Acre
Japanese beetle (adults, feeding on foliage) White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	16.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. 18-inch band on each side of the row followed with 0.25 inch of irrigation immediately after application. For optimal grub control, apply this product to control 1st or 2nd instar larvae. Application may be made post-bloom up to 7 days prior to harvest, or post-harvest until October 1st. For optimum control of Japanese beetle larvae, make applications from June 1 to July 15. Do not apply during bloom. Application to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field will control resident grub populations. Applications directed to the root zone will help protect berry plant roots from grub feeding. Apply this product to moist soil. If necessary, apply one hour of irrigation water immediately before application. To ensure maximum efficacy, 0.5 to 1 inch of irrigation water or rainfall should be applied or received within 24 hours of application of this product to facilitate movement into the soil and into the root zone.	

## BUSHBERRY - FOLIAR

Crops of Crop Subgroup 13B including: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Lingonberry, Salal

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers/Sharpshooters	2.4 to 3.2
Blueberry maggot Japanese beetles (adults) Thrips (foliage-feeding thrips only)	4.8 to 6.4
<b>Notes and Restrictions:</b> Pre-Harvest Interval (PHI): <b>3 days</b> Minimum interval between applications: <b>7 days</b> Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.5 lb. AI/A) Maximum number of applications of this product per crop season: <b>5</b> Maximum application volume (water): 20.0 GPA - ground; 5.0 GPA - aerial Do not apply pre-bloom or during bloom or when bees are actively foraging.	



# **CANEBERRY - SOIL ( For Use Only in CA )**

**Crops of Crop Subgroup 13A including:** Blackberry (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these). Raspberry (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*).

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Whiteflies	16.0 to 32.0
Rednecked cane borer	24.0 to 32.0
<b>Pests / Diseases Suppressed</b>	
Thrips (foliage-feeding thrips only)	16.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Maximum amount allowed per season: 32.0 fluid ounces/Acre (0.5 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Soil Application</b> Apply specified dosage in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. 2. Basal, soil drench in a minimum of 500 gallons solution per acre.	

# **CITRUS - SOIL ( Containerized )**

**Crops of Crop Group 10, plus White sapote including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote (*Casimiroa* spp), and other cultivars and/or hybrids of these.

Pests Controlled	Rate mL/ft <sup>3</sup> container media
Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Whiteflies	0.75
Citrus root weevil (larval complex)	1.25 to 2.50
<b>Pests / Diseases Suppressed</b>	
Citrus thrips (foliage-feeding thrips only)	2.50
<b>Application</b> Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of this product per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, treatment should be made at planting prior to insect infestation. Retreat if necessary. For control of larvae of the citrus root weevil complex, application should be made prior to neonate larvae entering potting media. Utilize higher dosage for heavy infestations.	

26/38

# **CITRUS - SOIL ( Field )**

**Crops of Crop Group 10, plus White sapote including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote (*Casimiroa* spp), and other cultivars and/or hybrids of these.

Pests Controlled	Rate fluid ounces/Acre
Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Termites ( FL only ) Whiteflies	16.0 to 32.0
Pests / Diseases Suppressed	
Citrus nematode Symptoms of: Citrus tristeza virus (CTV) through vector control Citrus yellows Thrips (foliage-feeding thrips only)	32.0
<p><b>Notes and Restrictions</b>  Pre-Harvest Interval (PHI): 0 day  Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre)</p> <p><b>Applications</b>  Apply specified dosage of this product in one of the following methods:</p> <ol style="list-style-type: none"> <li>1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil should be lightly pre-wetted to break soil surface tension prior to applications of this product. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move this product into root zone. Allow 24 hours before initiating subsequent irrigations;</li> <li>2. Soil surface band spray on both sides of the tree. Bands should overlap at the tree base to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root zone. This method is suitable for very coarse soils with 0.75% organic matter or less;</li> <li>3. Drench to base of tree not exceeding one quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only recommended for trees up to 8 feet tall;</li> <li>4. For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk;</li> <li>5. For suppression of citrus nematode, apply specific dosage through low pressure chemigation or soil surface spray only, ensuring complete coverage of the root system and utilizing application directions stated above for the respective application method. Repeated and regular use of this product over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.</li> </ol>	

## CITRUS - FOLIAR

Crops of Crop Group 10, plus White sapote including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, White sapote (*Casimiroa spp.*), and other cultivars and/or hybrids of these listed.

Pests Controlled	Rate fluid ounces/100 gallons	Rate fluid ounces/Acre
Aphids Asian citrus psyllid Black fly Leafhoppers/Sharpshooters Leafminers Mealy bugs Scales Whiteflies	2.8 to 4.0 (for dilute applications)	8.0 to 16.0 (depending on tree size, target pest and infestation pressure)
<b>Pests Suppressed</b>		
Thrips (foliage-feeding thrips only)	2.8 to 4.0	8.0 to 16.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 0 days Minimum interval between applications: 10 days Maximum amount allowed per crop season: 32.0 fluid ounces/Acre (0.5 lb. AI/A) Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.		
<b>Applications</b> Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. This product may be applied through properly calibrated ground or aerial equipment. Aerial application of this product may result in slower activity and reduced control to results from ground application. Scales - time applications to the crawler stage. Treat each generation. Where concentrated applications are appropriate, increase the spray solution concentration to apply an equivalent rate per acre to that applied in the diluted application. The 20.0 fluid ounce/Acre rate is based on full sized trees. This rate may be reduced proportionally for smaller trees.		

## COFFEE - SOIL

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Leafminer	16.0 to 32.0
<b>Pests Suppressed</b>	
Scales	16.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Maximum amount allowed per season: 32.0 fluid ounces/Acre (0.5 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<b>Applications</b> Apply specified dosage in one of the following methods: <ol style="list-style-type: none"> <li>1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.</li> <li>2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation;</li> <li>3. Basal, soil drench in sufficient water to insure incorporation into the root zone followed by irrigation.</li> </ol>	

28/38

# COFFEE - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Whiteflies	6.4
Pests Suppressed	
Scales	6.4
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 7 days Maximum amount allowed per crop season: <b>32.0 fluid ounces/Acre</b> (0.5 lb. AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<b>Applications</b> Apply specified dosage of this product as a broadcast or directed spray to infested area insuring thorough coverage. This product may be applied through properly calibrated ground or aerial application equipment. Aerial application of this product may result in slower activity and reduced control relative to results from ground application.	

# CRANBERRY - SOIL

Pests Controlled	Rate fluid ounces/Acre
Rootgrubs (Scarabaeidae) Rootworms (Chrysomelidae)	16.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 30 days Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<b>Applications</b> Apply this product to moist soil. Apply specified dosage of this product in one of the following methods: 1. As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre; 2. As a chemigation application with 600 to 1,000 gallons water. Immediately upon application, this product must be incorporated into root zone by 0.1 to 0.3 inch water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.	
<b>Rootgrubs and Rootworms</b> Best control may be achieved when application is made post-bloom immediately after bees are removed. Applications should target early instar larvae. This product has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of this product and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.	

29/38

## GRAPE - SOIL

Including: American bunch grape, Muscadine grape and Vinifera grape

Pests Controlled	Rate fluid ounces/Acre
European fruit lecanium Leafhoppers/Sharpshooters Mealybugs <i>Phylloxera</i> * spp	16.0 to 32.0
Pests / Diseases Suppressed	
Grapeleaf skeletonizer Nematodes Pierce's disease	24.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>30 days</b> Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre) <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation; 3. Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation. 4. For suppression of nematodes, apply 14 fluid ounces in a single application or two 7-fluid ounce applications on a 30 to 45-day interval. Treatments should be made only by 1) chemigation into root zone through above ground low pressure drip, trickle, micro sprinkler or equivalent equipment or 2) French plow technique, followed immediately by sufficient irrigation to move the product into the entire root zone of the plant. Repeated and regular use of this product over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response. For optimum results, make application between bud-break and the pea-berry stage. A total of 14 fluid ounces/acre is recommended under the following conditions: 1. Where vigorous vine growth is expected 2. In warmer growing areas 3. Where mealybug and European fruit lecanium populations are expected to be heavy 4. Where vine populations exceed 600 per acre, or; 5. For suppression of nematodes * Repeated and regular use of this product over several, consecutive growing seasons controls existing <i>Phylloxera</i> infestations over time or prevents <i>Phylloxera</i> from becoming established.	

## GRAPE - FOLIAR

American bunch grape, Muscadine grape and Viniferous grape

Pests Controlled	Rate fluid ounces/Acre
Leafhoppers/Sharpshooters Mealybugs	2.4 to 3.2
Grapeleaf skeletonizer	3.0 to 3.2
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>0 days</b> Minimum interval between applications: <b>14 days</b> Maximum amount allowed per season: <b>6.4 fluid ounces/Acre</b> (0.1 lb. AI/A) <b>Applications</b> Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. This product may be applied through properly calibrated ground or aerial equipment. For application on grapes, ground application is recommended.	

30/38

### HOP - SOIL

Pests Controlled	Rate fluid ounces/Acre
Aphids	19.2
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>60 days</b> Maximum amount allowed per season: <b>19.2 fluid ounces/Acre</b> (0.30 lb AI/Acre) <b>Applications</b> Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation; 3. Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation. Use higher dosages where extended residual control is desired or for treating larger vines or vines with dense foliage volume.	

### HOP - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids	6.4
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>28 days</b> Minimum interval between applications: <b>21 days</b> Maximum amount allowed per season: <b>19.2 fluid ounces/Acre</b> (0.3 lb. AI/A)	

### POME FRUIT - SOIL

Crops of Crop Group 11 including: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Pests Controlled	Rate fluid ounces/Acre
Aphids (including woolly apple aphid) Leathoppers	16.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>21 days</b> Maximum amount allowed per season: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Applications</b> Apply specified dosage of this product in the following method: Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	

# **POME FRUIT - FOLIAR**

Crops of Crop Group 11 Including: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Pests Controlled	Rate fluid ounces/100 gallons	Rate fluid ounces/Acre
Leathoppers	0.8 to 1.6	3.2 to 6.4
Aphids (except woolly apple aphid) Apple maggot Leafminers San Jose scale	1.6	6.4
FOR PEARS ONLY Mealybugs Pear psylla	4.0	16.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum of this product allowed per season: 32.0 fluid ounces/Acre (0.50 lb. AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Applications</b> Applications targeting apple maggot should be combined with manufacturer's recommended rate of a sticker.		

# **POMEGRANATE - SOIL**

Pests Controlled	Rate fluid ounces/Acre
Aphids Leathoppers/Sharpshooters Whiteflies	16.0 to 32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 0 day Maximum amount allowed per season: 32.0 fluid ounces/Acre (0.5 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Applications</b> Apply specified dosage of this product in the following method: Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	

# **POMEGRANATE - FOLIAR**

Pests Controlled	Rate fluid ounces/Acre
Aphids Leathoppers/Sharpshooters Whiteflies	6.4
<b>Pests Suppressed</b>	
Scales	6.4
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 7 days Maximum amount allowed per crop season: 19.2 fluid ounces/Acre (0.3 lb. AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

## STONE FRUIT - SOIL

**Crops of Crop Group 12 including:** Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

In-field, Soil Application	
Pests Controlled	Rate fluid ounces/Acre
Aphids (including woolly apple aphid) Leafhoppers	16.0 to 24.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>21 days</b> Maximum amount allowed per season: <b>24.0 fluid ounces/Acre</b> (0.38 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<b>Applications</b> Apply specified dosage of this product in the following method: Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	
Pre-plant, Root Dip Application	
Pests Controlled	Rate fluid ounces/10 gallons root-dip solution
Black peach aphid (infesting roots)	2.0
Mix this product at a rate of 2.0 fluid ounces per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in this product's solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment.	

## STONE FRUIT - FOLIAR

**Crops of Crop Group 12 including:** Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

Pests Controlled	Rate fluid ounces/100 gallons	Rate fluid ounces/Acre
Aphids Green June beetle Japanese beetle Leafhoppers/Sharpshooters Plant bugs Rose chafer San Jose scale	1.6	3.2 to 6.4
Cherry fruit fly	1.6	4.8 to 6.4
Pests Suppressed		
Plum curculio Stink bugs	1.6	6.4
<b>Notes and Restrictions for Apricot, Nectarine, Peach</b> Pre-Harvest Interval (PHI): <b>0 days</b> Minimum interval between applications: <b>7 days</b> Maximum amount allowed per season: <b>19.2 fluid ounces/Acre</b> (0.30 lb. AI/A) Minimum application volume (water): 50 GPA - ground application; 25 GPA - aerial application. Do not apply pre-bloom or during bloom or when bees are actively foraging.		
<b>Notes and Restrictions for Cherries, Plums, Plumcot, Prune</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>10 days</b> Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb. AI/A) Minimum application volume (water): 50 GPA - ground application; 25 GPA - aerial application. Do not apply pre-bloom or during bloom or when bees are actively foraging.		



33  
38

# **TREE NUTS - SOIL**

**Crops of Crop Group 14 including:** Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers/Sharpshooters Mealybugs Spittlebugs Termites Whiteflies	16.0 to 32.0
Pests / Diseases Suppressed	
Pecan scab (from reduction in honeydew deposition)	24.0 to 32.0
Thrips (foliage-feeding thrips only)	32.0
<p><b>Notes and Restrictions</b>            Pre-Harvest Interval (PHI): 7 days            Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre)            Do not apply pre-bloom or during bloom or when bees are actively foraging.</p> <p><b>Applications</b>            Apply specified dosage prior to or at onset of pest infestation in one of the following methods:</p> <ol style="list-style-type: none"> <li>1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. Pre-wet soil prior to applications of this product and allow soil to dry following application and prior to subsequent irrigation;</li> <li>2. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site;</li> <li>3. Shank or subsurface side-dress, injected to a depth just above or just within the root zone and between the trunk and drip line of the tree canopy. Product should be applied in a minimum of 10 gallons per acre using multiple shanks on both sides of trees. Ensure product placement is below sod or orchard floor debris. Irrigation covering entire treated area should follow within 48 hours to promote uptake by root system.</li> <li>4. For control of termites, apply specified dosage to slightly moist soil as a high-volume drench to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. Utilize sufficient carrier volume to penetrate the soil to a depth of 18 to 24 inches to obtain optimum control. Allow soil to dry following treatment and prior to applying any irrigation.</li> </ol> <p><b>Remarks</b>            Use the higher rates when applied by shank or subsurface side-dress, used on larger trees, soils with high clay content, for high plant populations, and/or where extended control is desired. Under some conditions, control may not occur for 14 or more days or until two (2) irrigations have been made. Applications made later in the season may result in reduced efficacy.</p>	

34/38

### TREE NUTS- FOLIAR

**Crops of Crop Group 14 including:** Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)

Pests Controlled	Rate fluid ounces/Acre
Aphids (except Black pecan aphid) Leafhoppers/Sharpshooters <i>Phylloxera</i> spp. (leaf infestations) Spittlebugs Whiteflies	2.8 to 5.6
Black pecan aphid Mealybugs San Jose scale	6.4
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 6 days Maximum amount allowed per crop season: <b>23.0 fluid ounces/Acre</b> (0.36 lb. AI/A) Minimum application volume (water): 50 GPA - ground application, 25 GPA - aerial application Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Applications</b> Applications for control of San Jose scale should be timed according to crawler stage, treating each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.	

### TROPICAL FRUIT - SOIL

**Including:** Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Jaboticaba, Guava, Llama, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursap, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

Pests Controlled	Rate fluid ounces/Acre
Aphids Avocado lacebug Leafhoppers Whiteflies	24.0 to 32.0
<b>Pests / Diseases Suppressed</b>	
Scales Thrips (foliage-feeding thrips only)	32.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 6 days Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Applications</b> Apply specified dosage of this product in the following method: Chemigation through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	

### TROPICAL FRUIT - FOLIAR

Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Liana, Jaboticaba, Guava, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers/Sharpshooters Mealybugs Thrips (foliage-feeding thrips only) Whiteflies	6.4
Pests Suppressed	
Scales	6.4
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum amount allowed per season: 32.0 fluid ounces/Acre (0.50 lb. AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

### OTHER CROPS - FOLIAR

#### Application Instructions

Apply specified rate per acre as a broadcast or directed foliar spray as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimal control. A spray adjuvant may be used to improve coverage. This product may not knock down established and heavy insect populations. Two applications may be required to achieve control. Scout fields and retreat if needed. This product may be tank mixed with other insecticides for knock down of pests or for improved control of other pests.

### CHRISTMAS TREE - SOIL

Pests Controlled	Rate fluid ounces/Acre
White grub complex (damage from grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and oriental beetle)	16.0 to 32.0
<b>Notes and Restrictions</b> Maximum amount allowed per season: 32.0 fluid ounces/Acre (0.5 lb. AI/Acre)	
<b>Applications</b> Soil incorporation and movement of this product to the root zone is required for activity. This product can be incorporated most readily when applied to moist soil. Apply specified dosage in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. 18-inch band on each side of the row (small trees) to full broadcast application (large trees) followed by rainfall or 0.25 to 1 inch of irrigation within 12 hours after application. For optimal grub control apply this product during adult flight activity, or up to mid-July, when first instar larvae are present.	

### CHRISTMAS TREE - FOLIAR

Pests Controlled	Rate fluid ounces/Acre
Aphids Adelgids Sawflies	3.2 to 6.4
<b>Notes and Restrictions</b> Minimum interval between applications: <b>7 days</b> Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb. AI/A) <b>Applications</b> Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. This product may be applied through properly calibrated ground or aerial equipment. Aerial application of this product may result in slower activity and reduced control relative to results from ground application. Gall-forming adelgids - time applications to coincide with full bud-swell or first bud-break of earliest bud-breaking trees. Once galls form spraying will be ineffective.	

### POPLAR/COTTONWOOD - SOIL

(Includes members of the genus *Populus* grown for pulp or timber)

Pests Controlled	Rate fluid ounces/Acre
Aphids Cottonwood leaf beetle	16.0 to 32.0
<b>Pests / Diseases Suppressed</b>	
<i>Phylloxera popularia</i>	16.0 to 32.0
<b>Notes and Restrictions</b> Maximum amount allowed at-plant per crop season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging. <b>Applications</b> Apply specified dosage of this product in the following method: 1. Chemigation through low-pressure drip irrigation. 2. For narrow row, cutting orchards/nurseries used for plant propagation, shank into root zone followed by adequate irrigation to promote uptake. Adequate irrigation depends on soil moisture level at application. Under dry conditions 0.25 inch/acre is recommended. For Cottonwood leaf beetle, protection against damage will occur when application is made early-season, when beetles first begin feeding. Larger trees may require earlier treatment as a result of slower uptake. For <i>Phylloxera</i> , apply early in the year, from break of dormancy through May.	

Cutting/Whip Application Instructions. See details above for Field Application Instructions.

Pests Controlled	Cutting Whip Soaking Solution fluid ounces of this product needed per 100 gallons
Cottonwood leaf beetle	13.3 to 26.6 (unhydrated cuttings/whips) 26.6 to 40.1 (partially hydrated cuttings/whips)
<b>Pests / Diseases Suppressed</b>	
Aphids	13.3 to 26.6 (unhydrated cuttings/whips)
Phylloxera popularia	26.6 to 40.1 (partially hydrated cuttings/whips)
<b>Notes and Restrictions</b> Maximum amount allowed at-plant per crop season: <b>32.0 fluid ounces/Acre</b> (0.50 lb AI/Acre) <b>Applications</b> Moisture content of cuttings/whips prior to application, the solution concentration and the length of soaking interval interact to affect the amount of product absorbed into plant material. For a constant soaking interval of 24 hours, drier cuttings/whips absorb a higher quantity of solution and require a lower concentration. Conversely, more hydrated cuttings/whips absorb less solution and require a higher concentration. Soaking of cuttings/whips should occur in a covered container in absence of UV light. Not all <i>Populus</i> sp. clones/varieties/hybrids have been tested for crop safety. Without specific knowledge about a particular <i>Populus</i> sp. clone/variety/hybrid, a small number of cuttings/whips of each should be treated and evaluated prior to commercial use. <u>Apply this product in one of the following cuttings/whips soaking methods:</u> For freshly cut (hydrated) cuttings/whips, soak plant material in specified solution concentration for 24 hours prior to cold storage. After removal from cold storage, plant as needed. For previously hydrated cuttings/whips removed from cold storage, allow plant material to reach room temperature and soak in specified solution concentration for 24 hours prior to planting. Proper care should be taken in disposal of any residual soaking solution. Solution may be applied to existing trees or other registered crops as long as all product label precautions and restrictions are observed.	

#### POPLAR/COTTONWOOD - FOLIAR

(Includes members of the genus *Populus* grown for pulp or timber)

Pests Controlled	Rate fluid ounces/Acre
Aphids Leaf beetles	3.2 to 6.4
<b>Notes and Restrictions</b> Minimum interval between applications: <b>10 days</b> Maximum amount allowed per season: <b>32.0 fluid ounces/Acre</b> (0.50 lb. AI/A) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

# **WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

# **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RV061708N)

NUPRID is a registered trademark of Nufarm Americas Inc.

Admire, Admire Pro, Calypso, Gaucho, Provado are registered trademarks of Bayer.

Leverage, Trimax and Trimax Pro are trademarks of Bayer.

Actara, Centric, Cruiser and Platinum are registered trademarks of the Syngenta Group Company.

Assail and Intruder are registered trademarks of Nippon Soda Company, LTD.

Advise and Gallant are trademarks of Agrilience, LLC.

Alias and Passada are trademarks of Makhteshim Agan of North America, Inc.

Clutch is a registered trademark of Arysta LifeScience.

Couraze is a trademark of Cheminova, Inc.

Cruiser is a registered trademark of the Syngenta Group Company.

Impulse, Macho and Macho Max are trademarks of Albaugh, Inc.

Venom is a trademark of Valent USA Corporation.

Widow is a trademark of Loveland Products, Inc.