

228-575

11/05/2008

1/10



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505C)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

228-575

Date of Issuance:

NOV 5 2008

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Atera LC Granular Insecticide

Name and Address of Registrant (include ZIP Code):

Nufarm Americas, Inc.  
150 Harvester Drive Suite 200  
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
  - a. Revise the EPA Registration Number to read, "EPA Reg. No. 228-575".
3. The data requirements for storage stability (830-6317) and corrosion characteristics (830-6320) have not been satisfied, and must be submitted within eighteen months of the date of this notice.

Signature of Approving Official:

Venus Eagle, Product Manager (01)  
Insecticide-Rodenticide Branch, Registration Division (7505P)

Date:

NOV 5 2008

2/10

4. The data requirements for Oxidation/Reduction (830-6314) and Explodabililty (830.6316) have not been satisfied, and must be submitted within eighteen months of the date of this notice.
5. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact Kable Bo Davis at (703) 306-0415.

Venus Eagle  
Product Manager (01)  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

Enclosure- Stamped Labeling

3/10

# Atera™ LC Granular

## INSECTICIDE

FOR CONTACT AND SYSTEMIC INSECT PEST CONTROL IN LAWNS AND LANDSCAPE ORNAMENTALS AROUND RESIDENTIAL, INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, AND ATHLETIC FIELDS.

**ACTIVE INGREDIENTS BY WEIGHT:**

|                                 |               |
|---------------------------------|---------------|
| Imidacloprid .....              | 0.31%         |
| Bifenthrin .....                | 0.16%         |
| <b>OTHER INGREDIENTS:</b> ..... | <u>99.53%</u> |
| <b>TOTAL</b> .....              | 100.00%       |

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
**NOV 5 2008**

*Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
228-575*

EPA REG. NO. 228-LTL  
EPA EST. NO.

Manufactured For  
**NUFARM AMERICAS INC.**  
150 Harvester Drive  
Burr Ridge, IL 60527



**NET CONTENTS**

00228-00000.20081105.EPA New  
NUP-07322

4  
19

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

| <b>USER SAFETY RECOMMENDATIONS</b>  |  |
|---|--|
| <b>Users Should:</b>  |  |
| <ul style="list-style-type: none"> <li>• Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.</li> <li>• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul> |  |

| <b>FIRST AID</b>  |   |
|-------------------|---|
| <b>IF IN EYES</b> | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul> |

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**NOTE TO PHYSICIAN**

This pesticide contains a pyrethroid. If significant quantities have been swallowed, promptly eliminate the ingested quantities from the stomach and intestines. Treatment is symptomatic and supportive. Alcohol, milk, cream, and other digestible fats and oils should NOT be administered as they may increase absorption.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is hallow, may result in groundwater contamination.

This product is toxic to bees exposed to treatment and for more than 5 days following treatment. Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Carefully read and understand the Directions for Use and restrictions before applying this product.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**RESTRICTIONS**

Apply using ground equipment only; do not apply by air.

Do not apply on the following: nurseries, commercial greenhouses, and grass grown for seed, golf courses and sod farms.

Do not allow people or pets to enter the treated area until the dust has settled.

Do not apply more than 132 pounds of this product (0.4 pounds of imidacloprid, 0.2 pounds of bifenthrin) per acre per application on residential use sites (i.e. around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, preschools or day care facilities).

This product made be applied at up to 160 pounds of product (0.5 pounds imidacloprid, 0.26 pounds of bifenthrin) per application on non-residential use sites (i.e. around institutional, public, commercial or industrial buildings, parks, recreational areas or athletic fields).

Total amount applied cannot exceed a total of 160 pounds (0.5 pounds if imidacloprid, 0.26 pounds of bifenthrin) per acre per year.

5/10

## INFORMATION

This product is a broad spectrum, long lasting contact and systemic insecticide for control of insects infesting lawns and ornamentals in landscapes and surrounding areas. This product can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, cemeteries, arboretums, and non-production greenhouses.

### RESISTANCE MANAGEMENT

The active ingredients in this product are imidacloprid, a Group 4A insecticide, and bifenthrin a Group 3 insecticide. Insect biotypes with acquired resistance to Group 4A and Group 3 insecticides may eventually dominate the insect population if Group 4A or Group 3 insecticides are used alone or repeatedly in the same field or in successive years as the primary method of control for targeted species. Previous extensive use of Group 4 or Group 3 insecticides in a given area may result in resistance to selected target insects. This may result in partial or total loss of control of those species by this product.

To delay insecticide resistance in insects that are at risk of developing resistance to insecticides consider:

- Avoid the consecutive use of other Group 3 or Group 4A insecticides on the same insect species.
- Use tank-mixtures or pre-mixes of insecticides from a different target site of action group as long as both products are registered for the same use.
- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your State Agricultural Experiment Station, State Extension Specialists, or other lawn experts and/or Nufarm Americas Inc. for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Nufarm Americas Inc.

### APPLICATION EQUIPMENT

Apply this product uniformly over the area being treated with normally used granular application equipment. Use either drop-type or rotary-type spreaders to apply this product.

Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately. In order to insure continuing delivery of the correct amount of this product further adjustments to the settings should be made periodically during application.

### APPLICATION INSTRUCTIONS

#### LAWNS, LAWN PERIMETERS AND LANDSCAPED AREAS

This product can be used for broad spectrum insect control in lawns, lawn perimeters, and landscaped areas. This product can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields and cemeteries.

This product has sufficient contact and residual activity to provide high levels of control. The need for an application can be based on historical monitoring of the presence of or anticipated occurrence of individual or multiple insect pest species, the site, previous records or experiences, current season adult trapping or other methods. Where multiple pests are present or anticipated, use the highest rates suggested to control the species that are present. Optimum control of white grubs will be achieved when applications are made just prior to egg laying or to early instar larvae of target pests, followed by sufficient irrigation or rainfall to move the active ingredient into the zone where the insect pests are active. Higher rates should be used when long residual control is desired or late season curative applications are made to turf already showing grub damage

This product can be applied to lawns at 40 to 132 pounds per acre per year (1.0 to 3.0 lbs/1000 sq ft). The rate is dependent on the target pest(s), their stage of development and the desired level of control. Up to three applications can be made each growing season but do not use more than 132 pounds of product (0.4 lb of imidacloprid active ingredient, 0.2 lb bifenthrin) per acre per application on residential use sites ( i.e. around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, preschools or day care facilities). This product may be applied up to 160 pounds of product (0.5 lb of imidacloprid, 0.26 lb bifenthrin) per application on non-residential use sites ( i.e. around institutional, public, commercial or industrial buildings; parks, recreational areas or athletic fields). Note: You may also use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 pound active ingredient per acre per year for use in lawns and perimeters. Consult your local State Agricultural Experiment Station, State Extension Turf Specialists or other lawn experts for specific information concerning the timing of applications.

For control of early season insect pests and for control of white grubs, make the first application of this product at peak adult activity or when other insect pests reach a damaging population in the, spring at a rate of 40 to 100 pounds per acre (1.0 to 2.3 lbs/1000 sq ft). If a total of 100 pounds per acre has not been applied before peak egg laying period of white grub adults, make another application to bring the total up to 100 pounds per acre just prior to egg laying or to early instar larvae of white grub adults. If no applications of this product have been, made, apply 80 pounds per acre (1.8 lbs/1000 sq ft) just prior to egg laying or to early instar larvae of target pests for control of white grubs.

For control of surface insects, do not water in until at least 24 hours after the application. If the target insects are in the thatch, lightly water in after application with enough water to wet the thatch zone. For control of armyworms, cutworms and other surface feeders, apply 40 to 80 pounds per acre (1.0 to 1.8 lbs/1000 sq ft) at the first sign of insects or insect damage. Repeat the application at

6  
1.2

monthly intervals or as needed to control additional flushes of insects. Up to a total of 132 pounds per acre per year (3.0 lbs/1,000 sq ft) can be applied.

For control of existing grub populations that are causing damage to lawns, water in after application with enough water to move the insecticide into the root zone where the grubs are feeding.

Avoid mowing lawn until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

**SPECIFIC USE DIRECTIONS**  
**LAWN AND LAWN PERIMETER APPLICATIONS**

| Application              | Target Pests  | Rate                         |                 |
|--------------------------|---|------------------------------|-----------------|
|                          |   | Pounds per 1,000 square feet | Pounds per acre |
| Lawn and Lawn Perimeters | Annual Bluegrass Weevil (Adults) <sup>1</sup><br>Annual Bluegrass Weevil (Larvae) <sup>1</sup><br>Ants, Nuisance<br>Ants, Imported Fire-<br>Armyworm species<br>Banks Grass Mite<br>Billbugs <sup>2</sup><br>Black Turfgrass Ataenius <sup>3</sup><br>Black Vine Weevil<br>Centipedes<br>Chinch Bugs<br>Crickets<br>Cutworm species<br>Earwigs<br>European Crane Fly<br>Fleas<br>Frit fly<br>Grasshoppers<br>Leafhoppers<br>Mealybugs<br>Millipedes<br>Mole Crickets <sup>4</sup><br>Pillbugs<br>Sod webworms<br>Sowbugs<br>Spittle bug<br>Sugarcane grub<br>Ticks <sup>5</sup><br>White grubs<br>(Asiatic Garden Beetle, European Chafer, Green June Beetle, Japanese Beetle; Northern Masked Chafer, <i>Phyllophaga</i> spp. (May or June Beetle), Oriental Beetle, Southern Masked Chafer) | 1.8 to 3.0                   | 88 to 132       |

**1 Annual Bluegrass Weevil:** Application can be timed for adult weevils as they move from their over-wintering sites into turf. This movement usually begins when Forsythia is in full bloom and continues to full bloom of dogwoods. Larvae from adult weevils that escape the application will be controlled with the application directed at adult weevils. Applications made after adult weevil activity will control larvae. Consult your State Cooperative Extension Service for more specific information regarding timing of application. White grubs will be controlled with an application targeted towards Annual Bluegrass Weevil.

**2 Billbugs:** Application can be made when adult Billbugs are first observed, usually in April and May in temperate regions. A spring application will also control Billbug larvae. Consult your State Cooperative Extension Service for more specific information regarding timing of application. White grubs will be controlled with an application targeted towards Billbugs at a rate of 100 pounds per acre or above.

**3 Black Turfgrass Ataenius:** Application can be timed to coincide with the first generation of Black Turfgrass Ataenius that usually appears during May. White grubs will be controlled with an application targeted towards Black Turfgrass Ataenius at a rate of 100 pounds per acre or above.

**4 Mole Crickets:** Application should be made during peak adult flight and egg lay. If the soil is dry, irrigate before application to stimulate insect activity and water after application to release the insecticide from the granule carrier.

**5** Including ticks that may transmit lime disease and Rocky Mountain spotted fever.

**NOTES:**

In New York State, this product may not be applied to any grass or lawn area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Do not apply more than a total of 132 pounds of this product (0.4 lb of imidacloprid active ingredient, 0.2 lb bifenthrin) per acre per season.

If the maximum season limit of this product has been applied and pest populations require additional treatments, use another registered pesticide labeled for control of the targeted insects. You may also use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 pound active ingredient per acre per season for use in lawns and lawn perimeters.

**LANDSCAPE ORNAMENTALS**

**APPLICATION INSTRUCTIONS**

This product is a broad spectrum, contact and long lasting systemic insect control product for a wide spectrum of insects infesting landscape ornamentals. This product can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, cemeteries, arboretums and non-production greenhouses.

For control of surface insects and insects that crawl along the surface of the soil or mulch under ornamentals, water in lightly after application to move the insecticide into the areas where insects are active. Water in with enough irrigation or rainfall to move the active ingredient into the zone where the target insects are active. Soil surface applications will provide control of foliage feeding insects; however, optimum control will not be achieved until the active ingredient is taken up by the plant roots and translocated to the area where insects are feeding.

**LANDSCAPE ORNAMENTALS APPLICATION**

**SOIL SURFACE OR MULCH APPLICATIONS**

| Application   | Target Pests   | Rate                         |                 |
|---|--|------------------------------|-----------------|
|   |  | Pounds per 1,000 square feet | Pounds per acre |
| Soil Insect Pest  | Ants, Nuisance<br>Ants, Imported Fire-Beet Armyworm<br>Beetles<br>Black Vine Weevils (Adults)<br>Centipedes<br>Crickets<br>Cutworms<br>Diaprepes (Adults)<br>Earwigs<br>Fall Armyworms<br>Grasshoppers<br>Millipedes<br>Pillbugs<br>White grubs<br>(Asiatic Garden Beetle, European Chafer, Green June Beetle, Japanese Beetle, Northern Masked Chafer, <i>Phyllophaga</i> spp. (May or June Beetle), Oriental Beetle, Southern Masked Chafer) | 1.8 to 3.0                   | 88 to 132       |
| <b>APPLICATION INSTRUCTIONS:</b> Apply enough water to wet the areas where the insects are active or water in after application with enough irrigation to move the insecticide into the areas where insects are active. |  |                              |                 |

8/10

**LANDSCAPE ORNAMENTALS APPLICATION (Continued)**  
**SOIL SURFACE OR MULCH APPLICATIONS**

| Application        | Target Pests  | Rate                         |                 |
|--------------------|---|------------------------------|-----------------|
|                    |   | Pounds per 1,000 square feet | Pounds per acre |
| Foliar Insect Pest | Adlegids<br>Aphids<br>Armored scales (suppression)<br>Azalea Lace Bug<br>Citrus Thrips<br>Eucalyptus Longhorned Borer<br>Flathead Borers<br>( including Bronze Birch Borer and Alder Borer)<br>Flea Beetles<br>Florida Wax Scale<br>Fungus Gnats<br>Glassy Winged Sharpshooter<br>Japanese Beetles (Adult)<br>Lace Bugs<br>Leafbeetles<br>( including Elm and Viburnum Leaf Beetles)<br>Leafhoppers<br>Leafminers<br>Leafrollers<br>Mealybugs<br>Pine Tip Moth larvae<br>Plant Bugs (including <i>Lygus</i> spp.)<br>Psyllids<br>Root Weevils (including Black Vine Weevils)<br>Royal Palm Bugs<br>Sawfly larvae<br>Scales, Soft<br>Thrips (suppression)<br>Treehoppers<br>Whiteflies | 1.8 to 3.0                   | 88 to 132       |

**APPLICATION INSTRUCTIONS:** Apply enough water to wet the root zone of the plants. Control of leaf feeding insects will be delayed until the insecticide is taken up by the roots and moved into the areas where the insects are feeding, usually 2 to 4 weeks.

**NOTES:**

Do not apply more than a total of 132 pounds of this product (0.4 lb of imidacloprid active ingredient, 0.2 lb bifenthrin) per acre per season.

If the maximum season limit of this product has been applied and pest populations require additional treatments, use another registered pesticide labeled for control of the targeted insects. You may also use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 pound active ingredient per acre per season for use in lawns and perimeters.



10/9

**PERIMETER APPLICATIONS**

**APPLICATION INSTRUCTIONS**

When treating perimeters of structures, make the application to the ground in a bandwidth of 5 to 10 feet adjacent to the structure.

**RATE CONVERSION CHART FOR PERIMETER APPLICATIONS**

| Rate of Application<br>Pounds per<br>1,000 sq. ft. | Band<br>Width<br>(Feet) | Pounds of this product per Linear Distance (Feet) Around Structure |     |     |     |      |      |      |      |
|--|-------------------------|--|-----|-----|-----|------|------|------|------|
|  |                         | 100  | 150 | 200 | 250 | 300  | 400  | 500  | 1000 |
| 1.8  | 5                       | 0.9  | 1.4 | 1.8 | 2.3 | 2.7  | 3.6  | 4.5  | 9.0  |
| 1.8  | 10                      | 1.8  | 2.7 | 3.6 | 4.5 | 5.4  | 7.2  | 9.0  | 18.0 |
| 2.8  | 5                       | 1.4  | 2.1 | 2.8 | 3.5 | 4.2  | 5.6  | 7.0  | 14.0 |
| 2.8  | 10                      | 2.8  | 4.2 | 5.6 | 7.0 | 8.4  | 11.2 | 14.0 | 28.0 |
| 3.6  | 5                       | 1.8  | 2.7 | 3.6 | 4.5 | 5.4  | 7.2  | 9.0  | 18.0 |
| 3.6  | 10                      | 3.6  | 5.4 | 7.2 | 9.0 | 10.8 | 14.4 | 18.0 | 36.0 |

**NOTE:** This table provides amounts which include the total area to be treated. This area includes the area of a circle whose radius is the bandwidth (this accounts for the area treated at the four corners of the structure) plus the linear distance treated times the bandwidth.

**OPTIONAL  
SPREADER SETTINGS**

*Note: Following will be a list of application spreaders and approximate settings.  
This list will vary dependent on market of spreaders. (This note will not appear on final printed label.)*

**Applications Instructions**

See Spreader Settings chart below. Adjust spreader to the appropriate setting. Make sure that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace making parallel passes. Do not use a spreader that applies granules in narrow, concentrated or uneven bands.

**Important:** It is important to follow the directions as indicated on this quality product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning.

**Note:** These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

**EXAMPLES**

| Spreader | 1.0 lbs./1,000 sq.ft. | 1.8 lbs./1,000 sq.ft. | 3.0 lbs./1,000 sq.ft. | 3.6 lbs./1,000 sq.ft. |
|----------|-----------------------|-----------------------|-----------------------|-----------------------|
| aa       | XX                    | XX                    | XX                    | XX                    |
| bb       | XX                    | XX                    | XX                    | XX                    |
| cc       | XX                    | XX                    | XX                    | XX                    |

10  
10

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** This product should be stored in its original container in a cool, dry locked place out of the reach of children and out of direct sunlight. Do not use or store near heat or open flame.

**To contain a spill:** Cover the spilled product to prevent the material from spreading. If the product container is damaged, place the entire container along with the product in a secure receptacle and clearly label the contents.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. **Paper and plastic bags:** Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS; OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV110508)

Atera is a trademark of Nufarm Americas Inc.