

228-578

08/25/2008

1/9



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs

Registration Division (H7505P)

1200 Pennsylvania Avenue NW

Washington, D.C. 20460

EPA Reg.  
Number:

228-576

Date of Issuance:

AUG 25 2008

Term of Issuance: Conditional

Name of Pesticide Product:

Atera 0.36 GC Granular  
Insecticide

## NOTICE OF PESTICIDE:

☒ Registration  
☐ Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. George Meindl  
Nurfarm Americas, Inc.  
150 Harvester Drive, Suite 200  
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3)(c)(2)(B).

1. Revise the EPA Registration Number to read, EPA Reg. No. "228-576".

Signature of Approving Official:

Venus Eagle  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

Date:

AUG 25 2008

Page Two  
228-576

2. Revise the confidential statement of formula (csf) with one product name **"Atera 0.36 GC Granular Insecticide"** in box 3 and 4.

3. Remove the reference to landscape ornamental use everywhere it appears on the label. This use is not supported on the parent label.

- On page 3 under the header **"General Information"**, delete **"Ornamentals in landscapes and surrounding areas"** and **"arboretums and non-production greenhouses"**.
- Additionally, on page 3, under **"Application Instructions"** delete **"perimeters and landscaped areas"** from the subheader and paragraph. Then revise the second sentence of the paragraph as follows, "This product can be used **on turfgrass** around.....golf courses and cemeteries."

4. The following changes are necessary due to the Agency's definitive language policy. Furthermore, without these changes the user may be mislead to believe that the language is advisory instead of mandatory.

- On page 4 under the heading **"Application Instructions"** rewrite the last sentence to read **"Use higher rates when long residual control is ..."**
- Under **"Annual Bluegrass Weevil"** also on page 4 rewrite the first sentence to read **"Time applications to control adult weevils ..."**
- Under **Annual Bluegrass Weevil...** rewrite the first sentence to read: **"Make applications at the first sign of wilting..."**

5. Make the following changes on page 5.

- Under **Billbug adults** rewrite the first sentence to read: **"Make applications when adult billbugs are first observed..."**
- Under **"Import Ants"** rewrite the last sentence to read **"Treat mounds with sufficient force ..."**
- Under **"Mole Cricket adults"** rewrite the third sentence to read **"Water Applications with in up to 0.5 inches of water..."**
- Under **"Ticks"** rewrite sentence five to read **"Make applications in the late fall and/or early spring to control adult ticks..."** Rewrite the last sentence to read **"Make applications as necessary from mid-spring to early fall..."**

Page Three  
228-576

6. Under "**Restrictions**" on page 6 make the following changes:

- Change the 3<sup>rd</sup> bullet from "**should be**" to "**must be**".
- Add another bullet listing the maximum amount of bifenthrin that can be applied in a residential area which is .2 lbs. ai/acre/per application.

5. Within eighteen months of the date of this registration, submit to the Agency the required one year storage stability study (830.6317) for the proposed product under warehouse conditions. The corrosion characteristics study (830.6320) may be carried out concurrently. It is recommended that observations be made at 0, 3, 6, 9, and 12 months. Additionally, data for oxidation/reduction (830.6314) and Explodability (830.6316) must also be submitted within the 18 month timeframe.

6. Submit two copies of your final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitute acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice contact Dani Daniel at 703 305-5409.

Enclosure: Agency product chemistry review dated 6/18/08  
Agency acute toxicity review dated 6/19/08

4/9

**RESTRICTED USE PESTICIDE**  
**DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS**  
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

# Atera™

## 0.36 GC Granular

### INSECTICIDE

FOR CONTACT AND SYSTEMIC INSECT PEST CONTROL IN TURF AND LANDSCAPE ORNAMENTALS ON GOLF COURSES, LAWNS AND LANDSCAPE ORNAMENTALS AROUND RESIDENTIAL, INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, ATHLETIC FIELDS, AND SOD FARMS.

DO NOT USE THIS PRODUCT ON GOLF COURSES AND SOD FARMS IN NASSAU COUNTY OR SUFFOLK COUNTY, NEW YORK.

ACTIVE INGREDIENTS BY WEIGHT:

Imidacloprid .....	0.20%
Bifenthrin .....	0.16%
OTHER INGREDIENTS: .....	99.64%
TOTAL .....	100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-  
EPA EST. NO.

MANUFACTURED FOR  
NUFARM AMERICAS INC.  
150 HARVESTER DRIVE  
BURR RIDGE, IL 60527



NET CONTENTS

00228-00000.2008020508.EPA New  
NUP-07310

ACCEPTED  
With COMMENTS  
In EPA Letter Dated:  
**AUG 25 2008**  
Under the Federal Insecticide, Fungicide  
and Rodenticide Act, As amended, for the  
pesticide Registered under EPA Reg. No:

228-576

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b> <ul style="list-style-type: none"> <li>• Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.</li> <li>• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>	
<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
This pesticide contains a pyrethroid. If significant quantities have been swallowed, promptly eliminate the ingested quantities from the stomach and intestines. Treatment is symptomatic and supportive. Alcohol, milk, cream, and other digestible fats and oils should NOT be administered as they may increase absorption.	

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This product is toxic to bees exposed to treatment and for more than 5 days following treatment. Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Carefully read and understand the Directions for Use and restrictions before applying this product.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Apply using ground equipment only; do not apply by air.

Do not apply within a 25 foot radius of the following: lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Do not apply on the following: nurseries, commercial greenhouses, and grass grown for seed.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves (made of any waterproof material)
- Protective eyewear

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or green houses.

Do not enter or allow people (or pets) to enter the treated area until the dust has settled.

### GENERAL INFORMATION

This product is a broad spectrum, long lasting contact and systemic insecticide for control of insects infesting turfgrass, sod and ornamentals in landscapes and surrounding areas. This product can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, golf courses, cemeteries, arboretums, and non-production greenhouses.

### RESISTANCE MANAGEMENT

The active ingredients in this product are imidacloprid, a Group 4A insecticide, and bifenthrin a Group 3 insecticide. Insect biotypes with acquired resistance to Group 4A and Group 3 insecticides may eventually dominate the insect population if Group 4A or Group 3 insecticides are used alone or repeatedly in the same field or in successive years as the primary method of control for targeted species. Previous extensive use of Group 4A or Group 3 insecticides in a given area may result in resistance to selected target insects. This may result in partial or total loss of control of those species by this product.

To delay insecticide resistance in insects that are at risk of developing resistance to insecticides consider:

- Avoid the consecutive use of other Group 3 or Group 4A insecticides on the same insect species.
- Use tank-mixtures or pre-mixes of insecticides from a different target site of action group as long as both products are registered for the same use.
- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your State Agricultural Experiment Station, State Extension Specialists, or other turf experts and/or Nufarm Americas Inc. for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Nufarm Americas Inc.

### APPLICATION EQUIPMENT

Apply this product uniformly over the area being treated with normally used granular application equipment. Use either drop-type or rotary-type spreaders to apply this product.

Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately. In order to insure continuing delivery of the correct amount of this product further adjustments to the settings should be made periodically during application.

### APPLICATION INSTRUCTIONS

#### TURFGRASS, PERIMETERS, AND LANDSCAPED AREAS

This product can be used for broad spectrum insect control in turfgrass, perimeters, and landscaped areas. This product can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, golf courses, cemeteries and other similar areas.

This product has sufficient contact and residual activity to provide high levels of control. The need for an application can be based on historical monitoring of the presence of or anticipated occurrence of individual or multiple insect pest species, the site, previous records or experiences, current season adult trapping or other methods. Where multiple pests are present or anticipated, use the highest rates suggested to control the species that are present. Optimum control of white grubs will be achieved when applications are made just prior to egg laying or to early instar larvae of target pests, followed by sufficient irrigation or rainfall to move the active

ingredient into the zone where the insect pests are active. Higher rates should be used when long residual control is desired or late season curative applications are made to turf already showing grub damage.

### APPLICATION RATES

(Be sure to read "APPLICATION EQUIPMENT" Section of this label)

Consult your local State Agricultural Experiment Station, or State Extension Turf Specialists for more specific information regarding timing of application.

Application	Target Pests	Rate	
		Pounds per 1,000 square feet	Pounds per acre
Turfgrasses	<b>Larvae of subsurface feeders:</b> Asiatic garden beetle Black turfgrass ataenius Black vine weevil European chafer European crane fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	2.3 to 2.9	100 to 125
	<b>Adults and/or larvae of subsurface and surface pests:</b> Centipedes Fleas Millipedes Ticks Imported Fire ants Mole crickets Pillbugs Sowbugs	1.7 to 2.9  OR  For heavy infestations, a single annual application of 4.6 to 5.7	75 to 125  OR  For heavy infestations, a single annual application of 200 to 250
	<b>Adults and/or larvae of surface pests:</b> Annual bluegrass weevil Billbugs Chinch bugs Frit fly Leafhoppers Mealybugs Nuisance ants	1.7 to 2.9	75 to 125
	<b>Larvae of surface feeders:</b> Armyworm Cutworm Sod webworms	1.1 to 2.9	50 to 125

NOTE: For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 250 pounds of product (0.5 lb of imidacloprid active ingredient; 0.4 lb bifenthrin) per acre per year. Avoid mowing turf or lawn area until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

**Annual Bluegrass Weevil (*Listronotus maculicollis*) adults:** Applications should be times to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when Forsythia is in full bloom and concludes when flowering dogwood (*Cornus florida*) is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

**Annual Bluegrass Weevil (*Listronotus maculicollis*) larvae:** For best results, applications should be made at the first sign of wilting of bluegrass. For the first generation, this wilting often occurs after full bloom of flowering dogwood (*Cornus florida*).

**Armyworms, Cutworms and Sod Webworms:** If the grass area is being maintained at a mowing height of greater than 1 inch, then higher application rates (up to 2.9 pounds per 1,000 square feet) may be required during periods of high pest pressure.

**Billbug adults:** Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperature regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

**Chinch Bugs:** Chinch bugs infest the base of grass plants and are often found in the thatch layer. Irrigation of the grass area before treatment will optimize the penetration of the insecticide to the area where the chinch bugs are located. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates (up to 2.9 pounds per 1,000 square feet) may be required to control populations that contain both nymphs and adults during the middle of the summer.

**Imported Fire Ants:** Apply as a broadcast treatment up to 5.7 pounds per 1,000 square feet. For mound treatment, apply 1/2 cup of this product per mound and then drench the mound with 1 to 2 gallons of water. The mounds should be treated with sufficient force to break their apex and allow the insecticide to flow into the ant tunnels.

**Mole Cricket adults:** Achieving acceptable control of adult mole crickets is difficult because grass areas are subject to continuous invasion during the early spring by this large active stage. For application when both adults and/or large nymphs are present and actively tunneling, irrigate prior to application if soil is dry and apply 2.9 pounds per 1,000 square feet. Applications should be watered in with up to 0.5 inches of water immediately after treatment. Grass areas that receive pressure from adult mole crickets should be treated within one month of peak egg hatch to ensure optimum control of subsequent nymph populations (see below). For heavily infested mole cricket areas, or in areas with a history of mole cricket infestation, use a single seasonal application rate of 200 to 250 pounds per acre (4.6 to 5.7 lb per 1,000 sq ft).

**Mole Cricket nymphs:** Grass areas that received adult mole cricket pressure in the spring should be treated within one month of peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides. Control of larger, more damaging, nymphs later in the year will require higher application rates to maintain acceptable control. For maximum effectiveness, irrigate prior to application if soil is dry, apply as a broadcast treatment at 2.9 pounds per 1,000 square feet. Applications should be watered in with up to 0.5 inches of water after treatment. For heavily infested mole cricket areas, or in areas with a history of mole cricket infestation, use a single seasonal application rate of 200 to 250 pounds per acre (4.6 to 5.7 lb per 1,000 sq ft).

**Pillbugs and Sowbugs:** For heavy infestations, apply 250 pounds per acre (5.7 lb per 1,000 sq ft) of this product.

**Ticks (Including ticks that may transmit Lyme disease and Rocky Mountain spotted fever):** Treat the entire area where exposure to ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed activity. Repeat application should be limited to no more than once per seven days. Deer ticks (*Ixodes* spp.) have a complicated life cycle that ranges over a two year period and involves four life stages. Applications should be made in the late fall and/or early spring to control adult ticks that are usually located on brush or grass above the soil surface and in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter. American dog ticks (*Dermacentor variabilis*) may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

#### OPTIONAL SPREADER SETTINGS

*Note: Following will be a list of application spreaders and approximate settings.  
This list will vary dependent on market of spreaders. (This note will not appear on final printed label.)*

#### Applications Instructions

See Spreader Settings chart below. Adjust spreader to the appropriate setting. Make sure that the spreader works properly and spreads granules uniformly. Apply to the border and then fill in the center to ensure the entire area is covered completely. To make sure that the product is evenly distributed, walk at a steady pace making parallel passes. Do not use a spreader that applies granules in narrow, concentrated or uneven bands.

**Important:** It is important to follow the directions as indicated on this quality product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning.

**Note:** These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

#### EXAMPLES

Spreader	50 lbs./Acre	75 lbs./Acre	100 lbs./Acre	125 lbs./Acre
aa	XX	XX	XX	XX
bb	XX	XX	XX	XX
cc	XX	XX	XX	XX



9/9

#### RESTRICTIONS

- Avoid runoff or puddling of irrigation water following application. Avoid application of this product to areas which are waterlogged and saturated, which will not allow penetration into the root zone of the plant. Do not apply more than 250 pounds (0.5 lb of imidacloprid and 0.4 lb of bifenthrin) per acre per year.
- Treated areas may be replanted with any crop specified on an imidacloprid and bifenthrin label, or with any crop for which a tolerance exists for the two active ingredients.
- For crops not listed on any imidacloprid and any bifenthrin label, or for crops which no tolerances for the active ingredients have been established, a 12-month plant-back interval should be observed.
- Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds.
- Do not apply by air.
- Not for use in nurseries or commercial greenhouses.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** This product should be stored in its original container in a cool, dry locked place out of the reach of children and out of direct sunlight. Do not use or store near heat or open flame.

**To contain a spill:** Cover the spilled product to prevent the material from spreading. If the product container is damaged, place the entire container along with the product in a secure receptacle and clearly label the contents.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. **Paper and plastic bags:** Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

#### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(NEW)

Atera is a trademark of Nufarm Americas Inc.