

228-595

11/3/2008

10 of 11



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

228-595

NOV 3 2008

Term of Issuance: conditional

NOTICE OF PESTICIDE:

x Registration
Reregistration
(under FIFRA, as amended)

Name of Pesticide Product:

Quincept Herbicide Premium
Lawn Weed Killer
Concentrate

Name and Address of Registrant (include ZIP Code):

Nufarm America Inc.
150 Harvester Drive
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
2. Submit the data listed below:
a. Within one year of the date on this registration notice, submit Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies.
3. Make the following label changes:
a. Change Hazards to Humans and Domestic Animals to read "Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing. Wear protective eyewear. Wear: Long-sleeved shirt and long pants, socks, shoes, and gloves."
b. Change the order of the First Aid, to have "If Swallowed" first.
c. Revise the EPA Registration Number to read, "EPA Reg. 228-595."

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Handwritten signature of Joanne I. Miller

Date:

NOV 3 2008

2011

page 2
EPA Reg. No. 228-595

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

30f11

ACCEPTED
with COMMENTS
In EPA Letter Dated:
NOV 3 2008
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

QUINCEPT® HERBICIDE

228-595

PREMIUM LAWN WEED KILLER Concentrate

FOR SELECTIVE BROADLEAF AND CERTAIN GRASSY WEED CONTROL IN RESIDENTIAL AND ORNAMENTAL TURFGRASSES. TO CONTROL CRABGRASS, DANDELION, CLOVER, HENBIT, PLANTAINS, WILD ONION, AND MANY OTHER LISTED LAWN WEEDS. ALSO FOR HIGHWAYS, RIGHT-OF-WAYS AND OTHER SIMILAR NONCROP AREAS.

CONTAINS 2,4-D, QUINCLORAC AND DICAMBA
1 QUART TREATS (COVERS) UP TO 5,000 SQ. FT.

ACTIVE INGREDIENTS:

2,4-D DMA*	6.557%
Quinclorac**	4.085%
Dicamba Acid***	0.681%
OTHER INGREDIENTS:	88.670%
TOTAL:	100.000%

Isomer Specific Method, Equivalent to:

* 2,4-D	5.446%, 1.0000 lbs./gal.
** Quinclorac	4.085%, 0.7500 lbs./gal.
*** Dicamba Acid	0.574%, 0.0574 lbs./gal.


**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-xxx
EPA EST. NO. 228-IL-1

Manufactured By
NUFARM AMERICAS INC.
150 Harvester Drive
Burr Ridge, IL 60527



NET CONTENTS: ___ GALS. ___ QT. ___ PT.

000228.00000.20080630.EPA.New
NUP-07309

40911

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or on clothing. Wear protective eyewear. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks, and
- Chemical-resistant gloves

See engineering controls for additional requirements and exceptions.

Engineering Controls Statements: If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 1 70.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (tolerant species only) and similar turf areas; also for use on Sod Farms.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Centipede, St. Augustinegrass, Dichondra, Bahiagrass, Zoysiagrass, Bermudagrass and Bentgrass nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperature over 90°F, turf injury may occur.

Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed the maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

Do not apply this product by air or through any type of irrigation equipment.

Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

Make no more than 2 applications per year. For ornamental lawns and turf, apply no more than 1.5 lbs. ae 2,4-D per acre per application and no more than 3.0 lbs. ae 2,4-D per acre per year, excluding spot treatments. For turf grown for seed or sod, apply no more than 2.0 lbs. ae 2,4-D per acre per application, with a minimum of 21 days between applications.

60911

This product combines three proven active ingredients [Quinclorac, 2,4-D, and Dicamba] to provide broad spectrum control of grassy and broadleaf weeds. This herbicide is absorbed by foliage and roots of weeds and translocated throughout the plant. USE AS DIRECTED TO CONTROL OR SUPPRESS THE FOLLOWING WEEDS:

WEEDS:

GRASSY WEEDS				
Crabgrass, Large	Carolina geranium	Hedge bindweed	Pokeweed	Tarweed
Crabgrass, Smooth	Carpetweed	Hedge mustard	Poorjoe	Thistle
Barnyardgrass	Catchweed bedstraw	Hemp	Povertyweed	Tick trefoil
Foxtail, Giant	Catsear	Henbit	Prostrate spurge	Toadflax
Foxtail, Green	Catnip	Hoary cress	Prostrate vervain	Trailing crown vetch
Foxtail, Yellow	Chickweed	Horsetail	Puncture vine	Tumble mustard
Signalgrass,	Chicory	Indiana mallow	Purslane	Tumble pigweed
Broadleaf	Cinquefoil	Ironweed	Ragweed	Tumbleweed
	Common chickweed	Jewelweed	Red clover	Velvet leaf
	Common mullein	Jimsonweed	Redroot pigweed	Venice mallow
	Common sowthistle	Kochia	Red sorrel	Veronica
BROADLEAF WEEDS		Knawel	Redstem filaree	Vervain
Alder	Corn chamomile	Knotweed	Rough cinquefoil	Vetch
Annual yellow sweet clover	Creeping Jenny	Lambsquarter	Rough fleabane	Violet, wild
Artichoke	Crimson clover	Lespedeza	Roundleafed marigold	Virginia buttonweed
Aster	Croton	Locoweed	Rush	Virginia creeper
Austrian fieldcress	Cudweed	Lupine	Russian pigweed	Virginia pepperweed
Bedstraw	Curly dock	Mallow	Russian thistle	Wavyleaf bullthistle
Beggartick	Curly indigo	Marshelder	St. Johnswort	Western clematis
Biden	Dandelion	Matchweed	Scarlet pimpernel	Western salsify
Bindweed	Dead Nettle	Mexicanweed	Scotch thistle	White clover
Bird vetch	Dock	Milk vetch	Sheep sorrel	White mustard
Bitterweed	Dollarweed	Milkweed	Shepherdspurse	Wild mustard
Bitter wintercress	Dogbane	bloodflower	Slender plantain	Wild aster
Black-eyed Susan	Dogfennel	Mugwort	Smallflower galinsoga	Wild buckwheat
Black medic	Elderberry	Morningglory	Smartweed	Wild carrot
Black mustard	English daisy	Mouseear chickweed	Smooth dock	Wild four-o'clock,
Blackseed plantain	Fall dandelion	Musk thistle	Smooth pigweed	Wild garlic
Blessed thistle	False dandelion	Mustard	Sneezeweed	Wild geranium
Blue lettuce	False flax	Narrowleaf plantain	Southern wild rose	Wild lettuce
Blue vervain	Fiddleneck	Narrowleaf vetch	Sowthistle	Wild marigold
Box elder	Field bindweed	Nettle	Spanishneedle	Wild onion
Bracted plantain	Field pansy	Orange hawkweed	Spatterdock	Wild parsnip
Brassbuttons	Flea bane (Daisy)	Oxalis	Speedwell	Wild radish
Bristly oxtongue	Flixweed	Oxeye daisy	Spiny amaranth	Wild rape
Broadleaf dock	Florida betony	Parsley-piert	Spiny cocklebur	Wild strawberry
Broadleaf plantain	Florida pusley	Parsnip	Spotted catsear	Wild sweet potato
Broomweed	Frenchweed	Pearlwort	Spotted knapweed	Wild vetch
Buckhorn	Galinsoga	Pennycress	Spotted spurge	Willow
Buckhorn plantain	Garlic mustard	Pennywort	Spurge	Witchweed
Bulbous buttercup	Goathead	Peppergrass	Spurweed	Woodsorrel
Bull nettle	Goatsbeard	Pepperweed	Stinging nettle	Woolly croton
Bull thistle	Goldenrod	Pigweed	Stinkweed	Woolly plantain
Burdock	Ground ivy	Pineywoods bedstraw	Stichwort	Wormseed
Burning nettle	Gumweed	Plains coeopsis	Strawberry clover	Yarrow
Bur ragweed	Hairy bittercress	Plantain	Sumac	Yellow Rocket
Burweed	Hairy fleabane	Poison hemlock	Sunflower	Yellow flower pepperweed
Buttercup	Hawkweed	Poison ivy	Sweet clover	and other broadleaf weeds.
Canada thistle	Heartleaf drymary	Poison oak	Tall nettle	

Some of these species may require repeat spot applications and/or use of the maximum recommended rate on this product label even under ideal conditions for application.

USE TIPS

GENERAL TIPS

- Shake well before using.
- Spray during the growing season when weeds are young and growing actively.
- For best results, do not mow for 2 days before or after treating.
- Avoid entire lawn treatments when air temperature exceeds 90°F.
- Do not allow spray to get on ornamental plants or vegetables since injury may result. Do not spray around roots of ornamentals and trees.
- Do not use on lawns with desirable clovers and legumes or on ornamentals.
- Avoid sprays which are too fine because they can drift and damage other plants.
- If certain tough weeds are not dead within 3 weeks, repeat with spot treatment application.

TIMING

Apply this product to young, actively growing weeds as a postemergence entire lawn or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds.

- Applications of this product made to annual grasses at 2 to 4 tiller may not provide complete control.
- **Newly seeded areas:** Application to grass seedlings is not recommended until after the third mowing.
- **Newly sodded, sprigged or plugged areas:** Application should be delayed until 3 to 4 weeks after the sodding, sprigging or plugging operations. Delay applications for 4 weeks after seeding and emergence of Kentucky bluegrass, perennial ryegrass and fine fescues.
- **Seeding:** Delay applications for 4 weeks after seeding and emergence of turf species listed in this label.
- **Mowing:** Do not mow for two days before or two days after application. Clippings for the first three mowing should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees or in vegetable gardens.

IRRIGATION AND RAINFALL:

- If soil moisture is not sufficient prior to application, irrigation may improve weed control. For best results do not water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is required.

USE SITES

This product can be applied postemergence to the following turfgrasses:

HIGHLY TOLERANT TURFGRASSES	MODERATELY TOLERANT TURFGRASSES	DO NOT APPLY TO THE FOLLOWING TURFGRASSES
COOL SEASON TURFGRASS Bluegrass, Kentucky Fescue, Tall Ryegrass, Annual Ryegrass, Perennial	COOL SEASON TURFGRASS Bentgrass, Creeping (1,3) Bluegrass, Annual (Poa Annua) (1) Bluegrass, Rough (Poa trivialis) Fescue, Chewing's Fescue, Fine (2) Fescue, Hard Fescue, Red WARM SEASON TURFGRASS Bermudagrass, Common (1) Bermudagrass, (Hybrids) (1) Zoysiagrass	Bahiagrass Bentgrass, Colonial Bentgrass, Seaside Buffalograss Carpetgrass Centipedegrass Dichondra Paspalum, Seashore St. Augustinegrass

1. Temporary yellowing and discoloration on these species may occur after application.
2. Only apply to fine fescue when it is part of a blend.
3. Application to established creeping Bentgrass must be applied in 2 to 3 split applications at 2.4 to 4.0 fl. oz. per 1,000 sq. ft.

Warm season turfgrasses such as Common Bermuda and Zoysia may see temporary discoloration. Do not apply during growth stages from dormancy to green-up and from green-up to dormancy.

80911

HOW MUCH TO USE

SPOT TREATMENT

Pressure Sprayer [Pump-up Sprayer]: Adjust spray nozzle to give a coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

AMOUNT OF CONCENTRATE (FLUID OUNCES)	AMOUNT OF WATER (GALLONS)	AREA TREATED (SQ. FEET)
6.4	4	1,000
3.2	2	500
1.6	1	250

ENTIRE LAWN TREATMENT

Dial Type Hose-end Sprayers: Spray lawn using a coarse spray. Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

- 1) Measure the total square foot area to be sprayed. To determine the total square foot area, multiply the length by the width of the lawn area to be treated. Subtract square footage of non-treatment areas including flower beds, shrub beds, driveways, and sidewalks.
- 2) The application rate of this product is 6.4 fluid ounces per 1,000 sq. ft of lawn. Add 6.4 fl. oz. of this product to the spray bottle [jar], [reservoir] for every 1,000 sq. ft. of lawn area to be treated.
- 3) Set the dial to the 2.5 fl. oz. per gallon mix rate.
- 4) Connect to hose, turn on water and spray evenly over the lawn treatment area.

AREA TO BE TREATED (SQ. FEET)	AMOUNT OF PRODUCT NEEDED (FLUID OUNCES)
1,000	6.4
2,000	12.8
3,000	19.2
4,000	25.6
5,000	32.0

Broadcast Application: Spray lawn using a coarse spray. Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

AREA TO BE TREATED (SQ. FEET)	AMOUNT OF PRODUCT NEEDED (FLUID OUNCES)
1,000	6.4
2,000	12.8
3,000	19.2
4,000	25.6
5,000	32.0

Ready-to-spray Hose-end Sprayers: [This product is packaged in a ready-to-spray packaging containing a plastic container fitted with a spray head that sprays the solution at a fixed concentration.] Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

This sprayer bottle will treat 5,000 sq. ft. when applied at 6.4 fluid ounces of this product per 1,000 sq. ft.

Instructions

1. Attach garden hose to spray head.
2. Turn on water supply.
3. Bend safety tab down and turn control valve to "water".
4. Point nozzle toward lawn and turn control valve to "on". Product will automatically mix with water.
5. Walk back and forth at a steady pace and spray evenly over the lawn treatment area.

90911

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(NEW)

QUINCEPT is a registered trademark of Nufarm Americas Inc.

[OPTIONAL BOX]

PRODUCT FACTS	
What it Does: Kills Broadleaf Weeds including Chickweed, Clover, Dandelion and Grassy Weeds including Crabgrass (See inside for complete list).	
Where to Use: (Icon weeds) Lawns: Kills the Weeds not the Lawn.	
When to Use: Avoid broadcast applications when air temperature exceeds 90°F.	
Questions and Comments: (Icon phone) (icon computer) Call xxx-xxx-xxxx or visit our website at www.xxxxxxxx.xxx	

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING CLAIMS - FRONT PANEL

- Kill Crabgrass
- [Also] Controls Crabgrass
- Kills crabgrass, dandelion, clover and 200+ weeds
- Controls Sensitive Weeds such as...
- Controls both grassy and broadleaf weeds in one easy step
- Controls Hard-to-Kill weeds such as...
- Kills Broadleaf and grassy weeds
- Kills over 200 weeds
- Kills Crabgrass and All Major Broadleaf Weeds
- Kills Dandelion, Clover and Other Broadleaf Weeds
- Kills Dandelion, Clover and Other Listed Broadleaf Weeds
- Use When Weeds Are Actively Growing
- Kills the Weeds, Not the Lawn
- Rainproof in Hours - Won't Wash Away
- Won't Harm the Home Lawn
- Illustration(s) of various weeds listed on label
- Easy to Use
- For Thick, Weed-Free Turf
- Three Way Control
- Apply to a Moist Lawn
- Contains Three Way Power
- One Quart Covers Up To 5,000 Square Feet, (One XXX covers up to XX,XXX square feet)
- Easy to Mix
- Just Add Water
- New Formulation
- Simple to use
- Proven Performance
- New!

110e 11

OPTIONAL CONTAINER LANGUAGE:

Containers: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank, and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal.

Once clean, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer.

Returnable-Refillable Containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.