

228-625

12/09/2009

1/19

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

December 9, 2009

Lizbeth Rea
Nufarm Americas, Inc.
150 Harvester Drive, Ste. 200
Burr Ridge, IL 60527

Dear Ms. Rea:

Subject: Notification of PBN change per Pesticide Registration Notice 98-10

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10, dated November 25, 2009, for the product "Nufarm Chlorpyrifos SPC 2 Insecticide," EPA Registration Number 228-625.

The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label bearing the new primary brand name submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please contact me directly at 703-308-8735 or email me at chao.julie@epa.gov.

Sincerely,

A handwritten signature in cursive script, appearing to read "Julie A. Chao".

Julie A. Chao
Insecticide-Rodenticide Branch
Registration Division (7505P)
Office of Pesticide Programs

2/19



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number. 228-625	2. EPA Product Manager John Hebert	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Nufarm Chlorpyrifos Pro 2 Insecticide	PM# 7	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. 150 Harvester Drive, Ste. 200 Burr Ridge, IL 60527 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
DEC - 9 2009

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Notification of a primary brand name change (Nufarm Chlorpyrifos SPC 2 Insecticide) per PRN 98-10. This notification is consistent with the provisions of PRN 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PRN 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2 x 2.5 gallons		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Lizbeth Rea	Title Regulatory Manager	Telephone No. (Include Area Code) 919/655-0701
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Manager	
4. Typed Name Lizbeth Rea	5. Date 11/25/2009	

Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL		
EPA Registration #	Date Submitted to EPA	Electronic file name
000228-000625	11/25/2009	000228-000625.20091125.PBNChange. Notification

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.



 Signature

11/25/2009


 Date

Lizbeth Rea

 Name (typed)

Regulatory Manager

 Title



5/19

RESTRICTED USE PESTICIDE
For retail sale and use by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

Nufarm Chlorpyrifos Pro-SPC 2 Insecticide

For control of listed pests on golf course turf and ornamental plants around industrial sites and road medians, fire ant treatment on nursery stock, and as a non-structural wood treatment.

ACTIVE INGREDIENT:

Chlorpyrifos (O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate)..... 24.66%

OTHER INGREDIENTS: 75.34%

TOTAL:.....100.00%

Contains 2 pounds of chlorpyrifos per gallon.

Contains petroleum distillates.

NOTIFICATION

DEC - 9 2009

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

See inside label booklet for additional FIRST AID and PRECAUTIONARY STATEMENTS

EPA Reg. No. 228-625


EPA Est. No.

Manufactured for:
Nufarm Americas Inc.
150 Harvester Drive
Burr Ridge, IL 60527

Nufarm Chlorpyrifos Pro-SPC 2
Insecticide contains chlorpyrifos, the
active ingredient used in Dursban®.

Net Contents:

000228-000625.2009112504.EPR.NotificationPBNChange.Notification



6/19

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING/AVISO**

Causes skin irritation and moderate eye irritation. Harmful if swallowed. Do not get on skin or on clothing. Avoid contact with eyes.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate and viton. For more information, follow the instructions in Supplement Three of PR Notice 93-7. If you want more options, follow the instructions for category G on an EPA chemical-resistance category selection chart.

All mixers, loaders, other applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves
- Chemical-resistant apron when mixing or loading or exposed to the concentrate
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposures
- A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

When handlers use closed systems or closed cab motorized ground application equipment in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations	
Users should:	
Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.	

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

7/19

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN

Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection, is an antidote. This product is an organophosphorus ester that inhibits cholinesterase. Contains petroleum distillate – vomiting may cause aspiration pneumonia.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, small mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

COMBUSTIBLE: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made out of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposures

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides [40 CFR Part 170]. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

For ground boom applications, do not apply within 25 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries and commercial fish ponds. Apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer. Use fine or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.

The applicator also must use all other measures necessary to control drift.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

GENERAL PEST CONTROL – EXTERIOR SURFACE AND PERIMETER TREATMENTS

Perimeter Treatment Applications:

This product is to be mixed with water and applied outdoors with pressurized sprayers as a general surface spray.

Precautions for Perimeter Treatments:

- Do not feed treated grass cuttings (hay) or seed screenings to livestock, or use treated hay for livestock bedding.
- Keep out of fish pools and other bodies of water.
- Do not treat vegetable gardens.
- Do not allow livestock to graze in treated areas.

Treatment Sites:

Apply this product around localized exterior surfaces and as a soil applied perimeter treatment of industrial plant sites, manufacturing plants, food processing plants and warehouses.

Note: Phytotoxicity – Environmental factors and varietal differences affects phytotoxic expression. In situations where phytotoxicity potential is of concern, spray a small group of plants and observe them for 7 to 10 days to determine phytotoxic potential before treating large numbers of those plants.

Application Rates:

Depending on the pest species and application method, product applications of 0.03% to 0.5% can be made. Refer to the “Application Methods” section of this label for additional information.

Mixing Directions:

To make a 0.03% water based spray, add 1 2/3 fl. oz. of this product per 10 gallons of spray. To make a 0.5% water based spray, add 2 2/3 fl. oz. of this product per each gallon of spray.

A stable emulsion of this product can be formed by first adding approximately one-half the water to the spray tank followed by adding the proper amount of this product. Close the tank and agitate for 5 to 10 seconds. Add the remaining water slowly to avoid excess foaming.

Tank Mixing:

Unless prohibited by a product's label, tank mix combinations can be used with pesticides labeled for similar use patterns. Always perform a small jar compatibility test using proper proportions of chemicals, and water to check for physical compatibility prior to tank mixing. Do not tank mix this product with products containing dichlorvos (DDVP).

Application Methods:

Apply this product outdoors as a general surface spray. Treat where pests are found or normally occur.

Low Volume Directed Sprays:

Low volume, high concentration sprays (0.5%) to localized exterior surfaces can quickly reduce localized heavy pest infestations on outside surfaces. Use a low pressure system with a pinpoint or variable pattern nozzle, such as a 1-gallon hand pump sprayer, and apply the spray mixture to specific areas such as cracks and crevices along walkways, patios, under eaves, windows and door frames or other areas where insects congregate or can gain entrance to the structure.

High Volume Broadcast Sprays:

High volume, low concentration perimeter sprays (0.03% - 0.12%), such as with power spraying equipment, can help prevent infestation of buildings by reducing pests in outdoor areas. Higher rates will provide longer residual control. To make a 0.03% water based spray, mix 8 fl. oz. of this product per 50 gallons. To make a 0.12% dilution, mix 32 fl. oz. of this product per 50 gallons of water. This type of treatment provides more thorough coverage over large areas than low volume directed sprays. Apply spray mixture directly to areas such as junctions of soil and structural walls and along base of fences. To help prevent infestation of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also treat the building foundation to a height of 2 to 3 feet, where pests are active and find entrance. Apply as a coarse spray at the rate of about 7 gallons spray mixture per 1,000 sq. ft. Thoroughly and uniformly wet the treated area.

Pests Controlled by Perimeter Treatments of Product:

- | | |
|------------------------------------|---------------------------|
| Ants | Elm leaf beetles (adults) |
| Bees | Fire ants ¹ |
| Beetles | Fleas |
| Boxelder bugs (or other true bugs) | Flies |
| Carpenter ants | Hornets |
| Centipedes | Millipedes |
| Clover mites | Pillbugs |
| Cockroaches | Scorpions ² |
| (American) | Sowbugs |
| (Asian) | Spiders |
| (Brownbanded) | Springtails |
| (German) | Ticks |
| (Oriental) | Wasps |
| (Smokybrown) | Yellowjackets |
| Crickets | |
| Earwigs | |

¹Control fire ant mounds by applying this product as a drench. Dilute 2 fl. oz. of this product per 4 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2-foot diameter. For best results, apply in cool weather (65°F to 80°F), or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays disturb the ants and cause migration, reducing product effectiveness.

²To control scorpions, remove accumulations of lumber, firewood and other materials serving as harborage sites. Prior to stacking firewood or lumber, apply this product as a localized spray to surfaces immediately below such materials.

GENERAL CONTROL OF WOOD-INFESTING INSECTS

Directions for General Use to Control Wood-Infesting Insects:

Mix this product with water and apply as a general surface or localized injection treatment with pressurized sprayers or other equipment suitable for applying insecticides to localized areas.

Treatment Sites:

When used according to label directions, this product can be applied to non-residential outdoor areas for control of wood-infesting insects. Apply this product to:

- Fence posts
- Utility poles
- Railroad ties Poles
- Landscape timbers Posts
- Logs

Application Rates:

Depending on the pest species and application method, product applications of up to 0.5% can be made. Increased residual control can be expected at higher rates (but not to exceed 0.5%). Refer to the "Specific Use Directions for Control of Wood-Infesting Insects" section of this label for additional information.

Mixing Directions:

To make a 0.5% water based spray, add 2 2/3 fl. oz. of this product per each gallon of spray mixture. To mix, add approximately one-half the required amount of water to the spray tank followed by the proper amount of this product. Close the tank and shake vigorously for 5 to 10 seconds. Slowly add the remaining water.

Tank Mixing:

Unless prohibited by a product's label, tank mix combinations can be used with pesticides labeled for similar use patterns. Always perform a small jar compatibility test using proper proportions of chemicals and water to check for physical compatibility prior to tank mixing. Do not tank mix this product with products containing dichlorvos (DDVP).

Application Methods:

Apply this product either as a coarse spray or by brushing onto targeted surfaces. Use equipment capable of delivering a coarse, low-pressure (approximately 20 psi) spray for treatment of large or overhead areas. Use sufficient spray to cover the area to the point of wetness but avoid applying to the point of runoff.

Wood-Infesting Insects Controlled by This Product:

- Beetles (including Anobiidae, Bostrichidae, Cerambycidae, and Lyctidae)
- Carpenter ants and other wood-infesting ants
- Carpenter bees
- Termites

Specific Use Directions for Control of Wood-Infesting Insects:

Beetles – Beetles are controlled by applying spray mixture to infested areas, or areas where infestations are likely to occur. Use the following guidelines to determine the appropriate rates of application:

- New Wood (typically less than 10 years of age) – Apply 1 gallon of dilution per 150 sq. ft.
- Old Wood (typically greater than 10 years of age) – Apply 1 gallon of dilution per 100 sq. ft.

Wood-Infesting Ants – Control wood-infesting ants by applying spray mixture around cracks or crevices, doors and windows, or other areas where ants enter, crawl, or hide on the exterior of industrial

plants, manufacturing plants, food processing plants and warehouses. Primary colonies are typically found outside through an exterior inspection. Make landscape modifications or correction of sanitation and structural deficiencies for effective control.

Termites – This product will kill workers and/or winged reproductive forms of termites present at the time of application only. Application of this product will not provide structural protection nor substitute for mechanical alteration, soil or foundation treatment with termiticide. This product is not to be used as sole protection against termites, as it is not intended for use for elimination of termite infestations nor for protection against future infestations. A professional inspection is recommended.

GENERAL PEST CONTROL

Golf Course Turf, Turf and Ornamentals in Industrial Plant Sites and Road Medians

Use this product to control injurious pests to 1) golf course turf and 2) turf and ornamentals in industrial plant sites and road medians that are not used for human or livestock consumption. This product is compatible with commonly used insecticides, miticides and fungicides except for alkaline materials such as Bordeaux mixture and lime. A small amount of spray mixture should be prepared to check for compatibility before a large volume of spray is mixed.

Precautions for Golf Course Turf, Road Medians and Industrial Plant Sites:

- Do not treat vegetable gardens.
- Do not allow livestock to graze in treated areas.
- Keep out of fish pools and other bodies of water.
- Do not feed treated grass cuttings (hay) or seed screenings to livestock or use hay for livestock bedding.
- Do not apply more than 1 lb. a.i./acre.
- Test in a small jar using proper proportions of chemicals and water be run to check for physical compatibility prior to tank mixing.

Turf in Golf Courses, Industrial Plant Sites, and Road Medians (Not for use in California):

Use this product to control the pests listed in the following table. Dilute this product in water and apply using suitable application equipment. Turf should be moist at the time of treatment.

PEST	Amount of Product per	
	1,000 sq. ft.	Acre
Ants	1 ½ fl. oz.	2 qts.
Armyworms (Beet, Fall, Yellowstriped)		
Centipedes		
Chiggers ¹		
Chinch bugs		
Crickets		
Cutworms		
Deer ticks ²		
Earwigs		
European crane fly larvae		
Fiery skipper		
Fire ants (foraging workers)		
Fire ants (mounds) ³		
Fleas		
Gnats		
Grasshoppers		
Greenbug aphids		
Green June beetle grubs		
Leafhoppers		

Lucerne moth Millipedes Mites (Bermudagrass stunt, Clover, Winter grain) Pillbugs Sod webworms (lawn moths) ⁴ Sowbugs Ticks ¹		
Billbug adults (Bluegrass, Denver, Hunting) ⁵	1 ½ fl. oz.	2 qts.
Annual bluegrass weevil (Hyperodes) ⁶ Black turfgrass ataenius adults ⁷ Mole crickets ⁸	1 ½ fl. oz.	2 qts.
Mole crickets ⁹	1 ½ fl. oz.	2 qts.
White grubs (Black turfgrass ataenius, European chafer, Japanese beetle larvae, and Northern and Southern masked chafers) ¹⁰	1 ½ fl. oz.	2 qts.

Specific Directions:

¹Use this product for area control of ticks and chiggers infesting non-cropland areas such as golf course turf and grassy areas where these pests are present and create a nuisance or a possible public health problem. Do not allow public use of treated areas during application or until spray has dried. Apply this product in water at the rate of 1 pint per acre (1/3 fl. oz. per 1,000 sq. ft.) using a hydraulic sprayer, mist applicator, backpack sprayer, or other suitable hand or power operated spray equipment.

²Apply this product at the rate of 2 quarts per acre or 1 ½ fl. oz. per 1,000 sq. ft. for control of deer ticks. Treat turf, grassy areas, low underbrush, weeds, ground surface and debris using enough spray volume to obtain thorough coverage.

³Apply this product as a drench for individual fire ant mounds. Dilute 2 fl. oz. per 4 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2-foot diameter. Apply in cool weather (65°F -80°F), or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays disturb the ants causing migration and reducing effectiveness of the product.

⁴For sod webworms, watering or mowing of the treated area should be delayed for 12 to 24 hours after treatment.

⁵Spray early in the season just prior to, or coinciding with first appearance of adult billbugs as specified by your local Agricultural Extension Service specialist.

⁶For control of annual bluegrass weevil, spray suspected problem areas in mid-April and again in mid-May, or as specified by your local Agricultural Extension Service specialist.

⁷For black turfgrass ataenius adults, spray early in the season as specified by your local Agricultural Extension Service specialist. Make a repeat application 1 to 2 weeks later, if needed.

⁸For control of mole crickets in turfgrass, apply this product through high-pressure injection or other suitable subsurface placement application equipment. Follow the manufacturer's instructions for calibration and the volume of spray per acre needed to provide control or as specified by your local Agricultural Extension Service specialist. Apply when young nymphs are active.

⁹For control of mole crickets in turfgrass, apply this product using broadcast or suitable hand-held application equipment. Make application in a minimum of 50 gallons of water per acre. Golf course turf must be irrigated within 24 hours after treatment to wash the insecticide into the area of insect activity. Apply when early stage nymphs are active. Effectiveness is enhanced by spraying late in the afternoon or early evening and irrigating the turf within 24 hours prior to and following application to move mole crickets near the soil surface and wash the insecticide into the zone of insect activity.

¹⁰For white grubs, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as specified by your local Agricultural Extension Service specialist. Soil should be moist prior to treatment. Immediately after spraying, irrigate the treated areas with ½ to 1 inch of water to wash the insecticide deep into the thatch or underlying soil.

ORNAMENTALS (For industrial plant site and road median use only): Use this product to treat evergreens, flowers, shrubs, vines, shade and flowering trees infested with the types of pests listed in the following tables. Dilute this product with water according to directions given in the tables and apply using suitable hand or power-operated spray equipment. Ensure complete and uniform coverage. Uniform coverage is critical for effective insect and mite control. Attempt to penetrate dense foliage but avoid over-spraying to the point of excessive runoff.

Consult your State Agricultural Experiment Station or Extension Service specialist for application timing and other specific use information for your area.

Note: Phytotoxicity – Environmental factors and varietal differences affect phytotoxic expression. Where phytotoxicity potential is of concern, treat a small group of plants and observe for 7 to 10 days to determine phytotoxic potential before treating large numbers of those plants.

SUCKING INSECTS AND MITES

Do not exceed 1 lb. ai/acre.

PEST	Amount of Product per	
	1 Gallon	100 Gallons
Adelgids (Cooley and Eastern spruce galls, Pine bark) Aphids (Apple, Balsam twig, Black pecan, Chrysanthemum, Cottonwood, Crape myrtle, Elm leaf, Melon, Peach, Rose, Spirea, White pine, Woolly, Woolly apple, Yellow pecan) Boxelder bugs Lace bugs (Hawthorn) Periodical cicada Plant bugs Psyllids Spittlebugs Thornbug Whiteflies	1/6-1/3 fl. oz.	1 pt.-1 qt.
Leafhoppers Mealybugs (Citrus, Taxus) Mites (Clover, Red spider, Southern Red, Spruce spider, Two spotted spider) ¹ Thrips (exposed)	1/3 fl. oz.	1 qt.
Scale insects (Cottony cushion, Cottony maple, Dearness, Euonymus, Fletcher, Florida wax, Golden oak, Hemispherical, Lecanium, Magnolia, Oak kermes, Oak lecanium, Oystershell, Pine needle, San Jose, Tea, White birch) ²	2/3 fl. oz.	2 qts.

Specific Directions:

¹For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days in the South or 7 to 10 days in the North after initial treatment to control newly hatched nymphs.

²Time application for control of scale insects when crawlers or first two stages of settled nymphs are present.

14/19

DEFOLIATORS AND LEAFMINERS

PEST	Amount of Product per	
	1 Gallon	100 Gallons
Armyworms (Fall, Yellowstriped) Bagworms ¹ Cankerworms Catalpa sphinx Elm spanworms Fall webworms ² Grasshoppers Greenstriped mapleworms Green fruitworms Hornworms Jackpine budworms Juniper webworms Katydid Leafrollers ³ Maple leafcutters ⁴ Oak skeletonizers Oleander caterpillars Orange tortrix Poplar tentmaker Puss caterpillars Rose chafers Sawflies, exposed (European pine, Pine, Pin oak, Redheaded) Spring elm caterpillars Springtails Spruce budworms (Eastern, Western) Tent caterpillars (Eastern, Forest, Western) Walnut caterpillars Yellownecked caterpillars	1/6-1/3 fl. oz.	1 pt.-1 qt.
Balsam gall midge Beet armyworms Beetles (Fuller rose) ⁵ Browntail moth Cutworms Cypress tip moth Douglas-fir tussock moth European pine shoot moth Gypsy moth ⁶ Holly bud moth Mahogany webworms Mimosa webworms Nantucket pine tip moth Oakworms (California, Orangestriped, Redhumped) Pandora moth Pitch pine tip moth Redhumped caterpillars Subtropical pine tip moth Tussock moth	1/3-2/3 fl. oz.	1-2 qts.

PEST	Amount of Product per	
	1 Gallon	100 Gallons
Beetles (Cottonwood leaf ⁷ , Elm leaf, Flea, Willow leaf)	1/3-2/3 fl. oz.	1-2 qts.
Leafminers Needleminers (Jeffrey pine, Lodgepole pine, Spruce) Pine needle midge Rhododendron gall midge	2/3 fl. oz.	2 qts.

Specific Directions:

- ¹For bagworms, treat when larvae are small and actively feeding.
- ²Direct spray into web and immediately surrounding foliage for effective control of fall webworms.
- ³For effective control of leafrollers, apply spray before leaves are tightly rolled.
- ⁴For maple leafcutter on maple trees, apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production.
- ⁵Make applications in the spring or early summer to reduce foliar feeding on twigs and branches by beetles.
- ⁶To control migrating and invading gypsy moth larvae, treat trunks and foliage.
- ⁷For cottonwood leaf beetles, spray larvae and adults infesting cottonwoods. Applications should be made when damaging beetle populations are developing or present.

BORERS, BARK BEETLES, and WEEVILS

PEST	Amount of Product per	
	1 Gallon	100 Gallons
Weevils (Black vine ¹ , Cranberry girdler ² , Pine reproduction, Yellow poplar)	1/3 fl. oz.	1 qt.
Borers: Clearwing moths (Ash, Dogwood, Lesser peachtree, Lilac, Oak, Peachtree, Rhododendron) Longhorned beetles (Cottonwood, Locust, Red oak) Metallic wood (Bronze birch, Flatheaded appletree, Twolined chestnut) ³ Pales weevil adults Zimmerman pine moth	2/3 fl. oz.	2 qts.
Northern pine weevil ⁴ Pales weevil ⁴	2 fl. oz.	6 qts.
Beetles ⁵ (Ambrosia, Anobiidae, Black turpentine, Blister, Cottonwood leaf, Elm leaf, European elm bark, Fuller rose, Japanese, June, Mountain pine, Native elm bark ⁶ , Southern pine, Spruce, Western pine, Willow leaf)	5 1/3 fl. oz.	4 gals.
Weevils (Northern pine, Pitch eating, Twig) ⁷	10 2/3 fl. oz.	8 gals.

Specific Directions:

- ¹Black vine weevils are night feeders. Late afternoon spraying will maximize control.
- ²For cranberry girdler larvae infesting Douglas fir seedlings, direct spray at lower crown and stems following egg laying during summer and irrigate immediately after application to move the insecticide into the top 1 to 2 inches of soil.
- ³For borers, spray trunks and lower limbs of trees and shrubs when adults begin to emerge. For peachtree borers, spray flowering trees and shrubs of the genus *Prunus* as a trunk spray before newly-hatched larvae enter the trees and thoroughly wet all bark areas from ground level to scaffold limbs. Pheromone traps

aid in detection of adult clearwing moths. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat for your area.

⁴Apply as a cut stump spray or drench in winter or early spring for control of northern pine and pales weevil larvae.

⁵For preventative treatment, spray the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment, spray the main trunk of infested trees or logs before adult beetles begin to emerge.

⁶To prevent native elm bark beetles from over-wintering in uninfested trees, apply using a dilution of 2 gallons per 100 gallons of water (2 2/3 fl. oz. of this product per gallon) as a spray to the bottom 9 feet of trunk. Wet the trunk thoroughly but do not spray to runoff. Apply the spray to the base of the root flare. Applications can be made from spring to early fall. To reduce twig and branch feeding on trees deemed to be of high value, apply as a spray to the tree crown using a dilution of 2 gallons per 100 gallons of water (2 2/3 fl. oz. of this product per gallon). Make applications in the spring or early summer using a sprayer that will give thorough coverage to the tree crown.

⁷For pine seedlings, treat immediately after transplanting. Treat each seedling with enough spray to thoroughly wet the foliage and stem to the point of runoff.

ANTS and OTHER LISTED PESTS

PEST	Amount of Product per	
	1 Gallon	100 Gallons
Ants Cockroaches (American, Asian, Brownbanded, German, Oriental, Smokybrown, Wood) Fire ants (foraging workers) Fire ants (mounds) ¹ Sowbugs Springtails	1/3 fl. oz.	1 qt.
Carpenter ants ²	5 1/3 fl. oz.	4 gals.

Specific Directions:

¹For individual fire ant mounds, apply this product as a drench. Dilute 2 fl. oz. of this product in 4 gallons of water. Sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2-foot diameter circle. Apply in cool weather (65°F -80°F), or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays disturb the ants and cause migration, reducing effectiveness of the product.

²If possible, locate carpenter ant nests and drench thoroughly.

Imported Fire Ant:

NOTE: USDA quarantine use only.

Treatment of potted, containerized and balled burlapped nursery stock: Use this product to control imported fire ants in soil attached to roots of these plants by completely submerging the ball of soil in a tank containing this product diluted at the rate of 8 fl. oz. per 100 gallons of water. Do not remove the burlap wrap or plastic containers with drain holes prior to submerging. Keep soil submerged until complete saturation has occurred, normally about 30 seconds. An alternate treatment to submerging potted plants is to apply a solution containing this product (8 fl. oz. per 100 gallons of water) as a substitute for plain water during routine watering activities. Apply this solution to the point of run-off, twice daily for three consecutive days. Do not remove the burlap wrap or container from plants prior to treatment. Make applications in a well-ventilated area.

ORNAMENTALS (Dormant Spray of Tree Pests):

Apply this product as a dormant or delayed dormant spray at the rates indicated to control the listed insects. Use this product without oil; however, use oil to control additional pests such as European red mites.

For high volume (dilute) sprays (200 to 600 gallons of spray mixture per acre), tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil for dormant use in 100 gallons of water. Spray the entire tree to runoff using suitable ground spray equipment.

For low volume (concentrate) sprays (less than 200 gallons of spray mixture per acre), use the same amount of this product as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use oil as specified by your State Agricultural Experiment Station or Extension Service specialist.

Precautions: Do not apply until rain or irrigation have replenished soil moisture such that bark and twigs are not desiccated since cold dry conditions causes this product plus oil to infuse trees resulting in bud damage or drop.

Restrictions: Make only one application during the dormant season. Do not allow meat or dairy animals to graze in treated areas.

PEST	Amount of Product per	
	1 Gallon	100 Gallons
Aphids (Mealy plum, Rosy apple, Woolly apple)	1/6-1/3 fl. oz.	1 pt.-1 qt.
Borers (Peach twig)		
Cutworms (Climbing)		
Leafrollers (Obliquebanded, Pandemis)		
Pear psylla adults		
Scale insects (San Jose)		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a secured dry storage area. Do not store above 122°F for extended periods of time. Preferred storage is above 32°F. Storage below 32°F may result in solidification. If warmed to above 32°F, product will return to original form. Freezing does not adversely affect this product. Dilutions of this product are sensitive to crystal formations when exposed to freezing, or near freezing temperatures (i.e., temperatures below 35°F, for extended periods of time). Dilutions allowed to stand overnight should be stored at temperatures above 40°F

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF

19/19

MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV11250409N)

Nufarm Chlorpyrifos ~~Pro~~-SPC_2 Insecticide is not manufactured or distributed by Dow AgroSciences, seller of Dursban®.

Dursban® is a registered trademark of Dow AgroSciences LLC.