

228-626

02-04-2010

1/16



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Lizbeth Rea
Regulatory Manager
Nufarm Americas Inc.
150 Harvester Drive Suite 200
Burr Ridge, IL 60527

FEB 04 2010

Dear Ms. Rea:

SUBJECT: Nufarm T-Methyl Pro 4.5 F Fungicide
EPA Registration Number 228-626
Your Amended Label Submitted September 3, 2009
OPPIN Decision Number 421023

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided you make the following changes:

Page 1 - heading in brackets at the top:

Revise: by removing the word "see," so that the sentence reads: "[Carton Front and Back Panel with reference to booklet for FIRST AID and Directions For Use; Booklet Cover]"

Page 3 - Directions for Use, and Page 9 - Turf Applications:

Remove: the sentence "Not for homeowner use." The label should only imply the specific user of the label; the Agency can not enforce otherwise.

Page 3 - Agricultural Use Requirements box:

Remove: the parenthesis contained within the last sentence starting with "PPE required for early entry to treated areas.....", and prior to the bulleted PPE.

Page 4 - Non-Agricultural Use Requirements box:

Revise: sentence to read "Do not enter or allow others to enter the treated areas without protective clothing until spray has dried."

Page 7 - Foliar Diseases table:

Revise: the third sentence in the Remarks column of the Fusicladium and Venturia Leaf Scabs row to read "Effective control requires coverage during leaf expansion."

Page 8 - Soil Drench Applications:

Revise: the table at the bottom of the page, to read: "Volume of Spray Mix to Apply to Each Container"

Pages 10 and 11– Turf disease control label:

Revise: the Rate of Product fl oz/1000 sq ft column, by reformatting the fractions "1/3" and "3/4" so that they are the same size as the "1/2" fraction.

Page 12 – Warranty Disclaimer:

Revise: the first sentence and last paragraph so that font is consistent with the remainder of this section.

A stamped label is enclosed for your records. If you have any questions, please contact Lisa Jones of my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely,



Shaja B. Joyner
Product Manager (20)
Fungicide Branch
Registration Division (7505P)

Enclosure: Stamped label

[Carton Front and Back Panel with reference to see booklet for FIRST AID and Directions For Use; Booklet Cover]

Nufarm T-Methyl SPC 4.5 F Fungicide

Liquid Flowable Systemic Turf and Ornamental Fungicide

For control of a broad spectrum of diseases of bedding, flowering, herbaceous and tropical foliage plants, shrubs, trees and flowers, containerized woody shrubs and trees and turfgrass.

ACTIVE INGREDIENT:

Thiophanate-methyl.....46.2%

OTHER INGREDIENTS:.....53.8%

TOTAL:.....100.0%

Contains 4.5 pounds thiophanate-methyl per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

See inside label booklet for additional FIRST AID and PRECAUTIONARY STATEMENTS

EPA Reg. No. 228-626

EPA Est. No.

Manufactured for:
Nufarm Americas Inc.
150 Harvester Drive
Burr Ridge, IL 60527

Nufarm T-Methyl SPC 4.5 F Fungicide contains thiophanate-methyl, the active ingredient used in Cleary 3336™.

Net Contents:

000228-000626.20091217.Label Amendment.AdditionalChanges

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

FEB 04 2010

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

228-626

[Page 1 of Booklet to end]

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION/PRECAUCION**

Causes moderate eye irritation. Harmful if swallowed or inhaled. Harmful if absorbed through skin. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils or viton ≥ 14 mils. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Handlers mixing, loading and applying the product as a dip must wear:

- Coveralls over long-sleeved shirt and long pants;
- Chemical-resistant gloves;
- Chemical-resistant footwear plus socks;
- A chemical-resistant apron

All other mixers and loaders and applicators must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Chemical-resistant gloves for all mixers and loaders and for applicators using hand-held equipment, and
- Chemical-resistant apron for mixers, loaders and other handlers exposed to the concentrate

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

When handlers use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for homeowner use. For use by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. Do not apply with fixed wing or rotary aircraft.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- Chemical-resistant footwear plus socks,
- Chemical-resistant headgear for overhead exposures

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides [40 CFR Part 170]. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas or allow others to enter without protective clothing until spray has dried.

GENERAL INFORMATION

This product provides broad spectrum disease control on containerized woody, flowering, herbaceous, and tropical foliage ornamental plants and trees, and turfgrasses.

Add the required amount of this product after shaking well to a partially filled tank agitated by mechanical or hydraulic means then add the remaining required amount of water. Maintain continuous agitation during application to keep the material in suspension. A new spray mixture must be prepared each day; do not use following overnight storage in spray tank.

Make applications of this product with ground equipment, using sufficient spray volume to provide thorough coverage. Do not tank mix this product with copper containing materials or with highly alkaline pesticides such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Under conditions of severe disease pressure or when application intervals are shorter than 14 days due to persistent rainfall, use the higher concentration or rates provided in this label. Contact your local State Extension Service specialist for application schedule instructions.

Important: If treatments of this product are ineffective, a tolerant strain of fungus is most likely present. Consult your Etigra representative or distributor, your local State Agricultural Experiment Station or State Agricultural Extension Service for advice on prompt use of some other labeled fungicide.

CHEMIGATION INSTRUCTIONS

For Ornamental Use Only

Apply this product only through pressurized drench (flood), sprinkler, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation systems.

Do not connect chemigation system to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact your State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Should the need arise, a person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments.

Pressurized Drench (Flood) System

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from back flow if water flow stops.

Sprinkler Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Drip (Trickle) Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to

prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

HORTICULTURAL APPLICATIONS

GREENHOUSE, NURSERY AND LANDSCAPE

This product provides broad spectrum control of many foliar, stem, and below ground diseases on a wide range of horticultural plants grown or maintained under a variety of conditions. Apply this product 14-21 days prior to when a particular disease usually appears and at the very latest, upon first sign of disease. Spray intervals usually range from 7 to 14 days with 14 days being for preventative treatments and the 7 day interval for times when conditions are judged acceptable for disease development. For hard-to-wet foliage, add an acceptable wetting agent to the spray tank to increase product efficacy. Use a spreader-sticker when excessive and repeated foliar wetting occurs. Use this product to control listed diseases on fruit and nut trees. Do not use fruit, nuts or sap from trees treated with this product as food or feed.

Note: This product has been determined to be safe for use on the plant types listed in these directions for use based on cumulative data derived from research product trials and historical field use. As all species and cultivars have not been tested, trial applications be performed if a user wishes to make an application to a plant type not listed on the label but found on a similar use site and for a disease that is listed on the label. To conduct a trial application, apply at least two applications to at least 25 trial plants at the highest concentration, 7 days apart. Evaluate 7 days after the last application before initiating full-scale application. Do not use this product on the following plants: Swedish Ivy (*Nephrolepis exaltata*), Boston Fern (*Plectranthus australis*), and Easter Cactus (*Hatiora gaertneri*).

Plant Type	Such as but not limited to:
Herbaceous Bedding	Ageratum, Begonia, Canna, Coleus, Dahlia, Dusty Miller, Foxglove, Fuchsia, Geranium, Impatiens, Lavender, Marigold, Pansy, Petunia, Pinks, Primrose, Salvia, Statice, Strawflower, Tickseed, Verbena
Flowering	Chrysanthemum, Hydrangea, Hollyhock, Iris, Lily, Poinsettia
Tropical Foliage	Dieffenbachia, Dracaena, English Ivy, Philodendron, Pothos
Woody Ornamentals	Azalea, Hibiscus, Holly, Ligustrum, Rhododendron, Rose, Pyracantha
Evergreen Trees	Douglas Fir, Fir, Larch, Pine, Spruce
Deciduous Trees*	Ash, London Plane, Maple, Oak, Sycamore, Walnut
Flowering Trees*	Cherry, Crabapple, Hawthorn, Mountain Ash, Pear

*Do not use fruit, nuts or sap from trees treated with this product as food or feed.

FOLIAR SPRAY APPLICATIONS

Hydraulic Application Mixing Instructions

After shaking the product container, add the required amount of this product to a partially filled spray tank agitated by mechanical or hydraulic means and then add the remaining specified volume of water. Maintain continuous agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Application Concentrations (Mechanical or Hand Held):

Use the specified amount of this product per 100 gallons of water for the prevention and control of the diseases shown below.

Note: The maximum single application rate for foliar application to ornamentals (excluding cut flowers) is 3.0 pounds thiophanate-methyl active ingredient (85.3 fl. oz. of product) per acre. The maximum application rate for foliar application to ornamentals per crop cycle is 300 pounds thiophanate-methyl active ingredient (66.7 gallons of product) per acre.

The maximum single application rate for application to cut flowers is 0.5 pounds thiophanate-methyl active ingredient (14.2 fl. oz. of product) per acre.

FOLIAR DISEASES

Disease(s) Controlled	Concentration of Product fl. oz./100 gals.	Remarks
Anthracnose <i>Colletotrichum</i>	10.75-20	Apply as buds break or at first sign of disease. Repeat application at 7-14 day intervals as needed during disease period.
Black Spot of Rose <i>Diplocarpon rosae</i>	10.75-20	Apply early summer or at first sign of disease. Repeat application every 7-14 days as needed during disease period.
Brown Rot and Blight <i>Monilinia</i> , <i>Sclerotini</i> , <i>Whetzellinia</i>	10.75-20	Apply late spring or at first sign of disease. Repeat application every 7-14 days as needed during the disease period.
Fusicladium and Venturia Leaf Scabs on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	10.75-20	Apply as buds break. Repeat application every 7-14 days during disease period. Effective control requires coverage during expansion. Rotations with chlorothalonil or propiconazole can be utilized.
Leaf Spots and Blights caused by: <i>Ascochyta</i> , <i>Blumeriella</i> , <i>Botrytis</i> , <i>Cercospora</i> , <i>Coccomyces</i> , <i>Corynespora</i> , <i>Curvularia</i> , <i>Didymellina</i> , <i>Entomosporium</i> , <i>Fabraea</i> , <i>Fusarium</i> , <i>Ramularia</i> , <i>Rhizoctonia</i> , <i>Marssoninia</i> , <i>Mycosphaerella</i> , <i>Myrothecium</i> , <i>Phoma</i> , <i>Physalasporea</i> , <i>Schizothyrium</i> , <i>Septoria</i> ,	10-14.5	Make applications when disease symptoms first appear. Repeat every 7-14 days during disease period. Rotate with chlorothalonil.

Disease(s) Controlled	Concentration of Product fl. oz./100 gals.	Remarks
<i>Sphaceloma</i>		
Ovulinia Blight	7.25-20	Apply as flowers open. Repeat every 7-14 days during disease period.
Powdery Mildews <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i>	10-20	Apply when disease first appears and repeat application every 7-14 days. Rotate with other effective products.
Rust Diseases caused by: <i>Puccinia, Gymnosporangium, Uromyces</i>	10.75-20	Apply late spring or when symptoms first appear. Repeat application every 7-14 days during disease period. Rotate with other effective products.
Tip Blight of Pine <i>Sphaeropsis sapinea, Diplodia pinea</i>	14.5-20	Begin applications in the spring when new growth starts. Make a second application just prior to needle emergence from the sheath and a third application 7 days later. Ensure thorough coverage.
Twig Blights, Cankers, and Diebacks <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	14.5-20	Apply when symptoms first appear. Repeat application every 7-14 days during disease period.

Adjuvants: Where rainfall and/or overhead irrigation is the norm, use of a compatible spreader/sticker is suggested. Where wetting of foliage is difficult, use a compatible wetting agent. Follow the phytotoxicity precautions described in the "HORTICULTURAL APPLICATIONS" section of this label.

SOIL DRENCH APPLICATIONS

Mixing Instructions: Add the required amount of this product to a partially filled tank agitated by mechanical or hydraulic means. Add the remaining required amount of water. Maintain continuous agitation throughout application to keep the material in suspension.

Application Concentrations/Rates and Timing for Disease Control: Use 7.5 to 20 fl. oz. of this product per 100 gallons of water. Apply as a drench or heavy spray at the rate of ½ to 2 pints per square foot (100 gallons per 400 to 1,600 square feet). For small pots and shallow flats up to 4 inches in size, apply at 1 pint per square foot. For containers and pots 4 inches or larger, refer to the following table for the volume to apply. Make repeat applications at 4 to 8 week intervals depending on disease presence and conditions for disease development.

Container Type	Volume to Apply/Container	
	1 pt./sq. ft. Rate	2 pt./sq. ft. Rate
4 inch	2 fl. oz.	
5 inch	2 ½ fl. oz.	
6 inch		6 ½ fl. oz.
7 inch		8 ½ fl. oz.
8 inch		11 fl. oz.
9 inch		14 fl. oz.
10 inch		17 ½ fl. oz.

For containers larger than 10 inches, apply in a drench volume of 2 ½ to 3 pints per square foot of surface area.

Plant Types: Containerized woody shrubs; trees, herbaceous/bedding, flowering, and tropical foliage plants and flowers and bedding plants in the landscape.

Note: Do not apply this product to plug trays or seedling flats at time of seeding.

Soil Diseases Controlled: Botrytis, Cyndrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia and Thielaviopsis.

Note: Pythium, Phytophthora and Cyndrocladium spathiphylli are not controlled by this product.

PLANT DIP TREATMENT

Mixing Instructions: Mix as described in the "FOLIAR" and "SOIL DRENCH APPLICATIONS" sections of this label. Maintain continuous agitation during application.

Application Concentration and Dipping Time

Plants or Cuttings: Use 14.5 to 21.5 fl. oz. of this product per 100 gallons of water. Immerse plants or cuttings for 10 to 15 minutes, remove and allow to drain and dry. Wear protective clothing as described under the "PERSONAL PROTECTIVE EQUIPMENT" section of this label.

Bulbs, Corms, Tubers and Rhizomes - Use 14.5 to 33 fl. oz. of this product per 100 gallons of water or 2 teaspoons of this product per gallon of water. Soak cleaned bulbs for 15 to 30 minutes in warm dip (80-85°F) within 48 hours of digging. Dry bulbs after treatment. If bulbs are for forcing, treat bulbs that have been heat-cured.

Plant Types: Plants, cuttings, cane sections of woody herbaceous, flowering and tropical foliage plants. Bulbs, corms, tubers, and rhizomes of plants such as but not limited to Caladium, Easter Lily, Tulip, Gladiolus, Daffodil, Iris.

Diseases Controlled: Botryis, Cyndrocladium, Fusarium, Gliocladium, Myrothecium, Penicilium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis.

TURF APPLICATIONS

Not for homeowner use. For use only by certified applicators or those under their immediate supervision. Do not apply with fixed wing or rotary aircraft. Not for use on turf being grown for sale or other commercial use as sod.

Use this product against certain foliar and soil diseases for use on all turf types such as golf course greens, tees and fairways, athletic fields, cemeteries, parks, and commercial and home lawns. Use this product both preventatively and curatively and is not phytotoxic. Do not use this product on turf being grown for sale or other commercial uses as sod.

Mixing Instructions: Add the required amount of this product to a partially filled tank agitated by mechanical or hydraulic means. Add the remaining required amount of water. Maintain continuous agitation to keep the material in suspension. For best results, use spray mixture the same day it is prepared.

Turf Types: All cool season and warm season grasses (such as but not limited to Bentgrasses, Bermudagrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustinegrasses and Zoysiagrasses) or their mixtures.

12
16

Disease(s) Controlled	Rate of Product fl. oz./1,000 sq. ft.**	Remarks
Anthracnose <i>Collectotricum graminicola</i>	1 3/4-3 1/2 (3 1/2 - 5 1/3)*	Apply when disease first appears. Make additional applications at 14 day intervals as needed. Allow spray to dry on leaves with no "watering in".
Ascochyta Leaf Blight <i>Ascochyta</i> spp. Dollar Spot <i>Sclerotinia homoeocarpa</i> , <i>Moellerodiscus</i> spp., <i>Lanzia</i> spp. Copper Spot <i>Gloeocerospora sorghi</i> Brown Patch and Zoysia Patch <i>Rhizoctonia solani</i> Fusarium Patch <i>Fusarium nivale</i> Red Thread <i>Laetisaria fusiformis</i>	1 3/4-3 1/2	Apply when disease first appears. Make additional applications at 14 day intervals as needed. Allow spray to dry on leaves with no "watering in".
Pink Snow Mold <i>Microdochium nivale</i> (Only for those areas where snow cover is not present the entire winter)	1 3/4-3 1/2	Apply this product in middle to late November before turf has stopped all growth activity. Lightly water application into the root zone for best results. For best results, use a spreader-sticker. Second spray should dry on leaf surfaces with no "watering-in". Minimum spray interval is 14 days.
Gray Leaf Spot <i>Pyricularia grisea</i>	3 1/2 - 5 1/3	Apply when conditions are favorable for disease development. Continue applications at 14 day intervals. Allow spray to dry on leaves with no "watering in".
Summer Patch <i>Magnaportha poae</i>	3 1/2 - 5 1/3	For preventative treatment, make 3 applications at 21 day intervals beginning in early May. Water product into the root zone thoroughly after application. For suppression, apply two applications at 14 day intervals beginning when disease first appears.
Cool Season Brown Patch <i>Rhizoctonia cerealis</i> Fusarium Blight <i>Fusarium</i> spp. Necrotic Ring Spot and Spring Dead Spot <i>Leptosphaeria korrae</i>	3 1/2 - 5 1/3	Make two applications at 14 day intervals beginning when disease first appears.

13
16

Disease(s) Controlled	Rate of Product fl. oz./1,000 sq. ft. **	Remarks
Stripe Smut <i>Ustilago striformis</i>	3 1/2 - 5 1/3	Make two applications at 14 day intervals when disease first appears. Water product into the root zone after application. For prevention, apply in the spring (just before grass begins growth), and in the fall.
Bermudagrass Decline <i>Gaeumannomyces graminis</i> var. <i>graminis</i> Take-All Patch <i>Gaeumannomyces graminis</i> var. <i>avenae</i>	3 1/2 - 5 1/3	Make applications in Mid-July or when disease symptoms first appear. Repeat applications at 14-day intervals. Use higher rates under severe disease pressure. Water product into root zone.
Coprinus Snow Mold <i>Coprinus psychromorbidus</i>	3 1/2 - 5 1/3	Make 2 applications at 21-day intervals starting in late fall to early winter. Make the last application just prior to first permanent snow cover.
Leaf Spot <i>Drechsler</i> spp. Leaf Crown & Root Diseases <i>Bipolaris</i> , <i>Curvalria</i> , <i>Exserohilum</i> Rusts <i>Puccinia</i> spp., <i>Uromyces</i> spp.	3 1/2 - 5 1/3	Start applications when disease first appears and at 14-day intervals as needed.
Bentgrass Dead Spot <i>Ophiosphaerella agrostis</i>	3 1/2 - 5 1/3	Based on the local Extension Service instructions, start applications in early June to prevent bentgrass dead spot and every 14 days as needed.

*Use 3 1/2 - 5 1/3 fl. oz. rate for curative response to Basal Stem Anthracnose.

**Refer to the "Use Sites and Maximum Application Rates" table to determine allowable rates for each application.

Turf Application Directions

Apply this product uniformly over the area to be treated with a properly calibrated power sprayer. Apply after mowing or avoid mowing for 12 hours after application. Apply sufficient water to obtain thorough coverage, usually 1 1/2 to 2 1/2 gallons per 1,000 sq. ft. of turf area. When treating golf greens, always treat aprons and approaches to golf greens. The minimum retreatment interval for turf applications is 14 days.

Use Sites and Maximum Application Rates

Site	Maximum Single Application Rate fl. oz./1,000 sq. ft.	Maximum Seasonal Application Rate fl. oz./1,000 sq. ft.
Golf Course Greens, Tees and Aprons	5 1/3	14 1/4
Golf Course Fairways (Except Florida)	3 1/2	3 1/2
Golf Course Fairways (Florida Only)	1 3/4	1 3/4
Residential and Public Areas (home lawns, parks, athletic fields, schools, day care centers)	1 3/4	7

Note to User: Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

Shake well before using.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, temperature controlled area.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER

15
16

MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV090309N)

Nufarm T-Methyl SPC 4.5 F Fungicide is not manufactured or distributed by Cleary Chemical Corporation, seller of Cleary 3336™.
Cleary 3336™ is a trademark of the Cleary Chemical Corporation.

16

16

NOT TO BE PART OF THE PRINTED LABEL
FOR INTERNAL REGULATORY USE ONLY

- 07/15/2009 Submission of primary brand name and minor label (replaced product name throughout label with "this product", replaced "recommend" and "may" with more definitive language, updated the Warranty statement) changes.
- 09/03/2009 Label amendment incorporating pests currently on Cleary Chemical Corp's 3336 F (1001-69) label that is not on this label. (lr)