



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

May 22, 2025

Joseph C. Wolfe  
Regulatory Manager, T&O  
NUFARM Americas, INC.  
11901 S. Austin Avenue  
Alsip IL, 60803

Subject: Label Amendment – Lower the Product Use Rate Minimums for Foliar Anthracnose and Dollar Spot in the Turf Disease Control table on pages 13 & 14  
Product Name: NUFARM T-METHYL SPC 4.5 F FUNGICIDE  
EPA Registration Number: 228-626  
Application Date: February 27, 2024  
Case Number: 501913

Dear Mr. Wolfe:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find

or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Ernest Kraka at [kraka.ernest@epa.gov](mailto:kraka.ernest@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Ernest Kraka", with a horizontal line drawn through the middle of the signature.

Ernest Kraka, Ph.D., Biologist/Risk Manager  
Fungicide and Herbicide Branch  
Registration Division 7505T  
Office of Pesticide Programs

Enclosure

[Carton Front and Back Panel with reference to booklet for FIRST AID and Directions For Use;  
Booklet Cover]

THIOPHANATE-METHYL

GROUP

1

FUNGICIDE

## Nufarm T-Methyl SPC 4.5 F Fungicide

[Alternate Brand Name: LESCO® T-STORM™ Flowable Turf & Ornamental Fungicide]

**[Liquid Flowable Systemic Turf and Ornamental Fungicide]**

[Optional – Alternate Product Information Statements]

[For the prevention and control of turf diseases and the diseases of annual and perennial flowers,  
bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs]

[For control of a broad spectrum of diseases of bedding, flowering, herbaceous and tropical foliage  
plants, shrubs, trees and flowers, containerized woody shrubs and trees and turfgrass]

### ACTIVE INGREDIENT:

Thiophanate-methyl .....46.2%

**OTHER INGREDIENTS:**.....53.8%

**TOTAL:** .....100.0%

Contains 4.5 pounds thiophanate-methyl per gallon

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION / PRECAUCIÓN**

See inside label booklet for FIRST AID and PRECAUTIONARY STATEMENTS

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only, Call (877) 325-1840**

EPA Reg. No. 228-626

EPA Est. No. \_\_\_\_\_

MANUFACTURED FOR  
Nufarm Americas Inc.  
11901 South Austin Avenue  
Alsip, IL 60803



[Nufarm: Grow a better tomorrow.]  
[Grow a better tomorrow.]

NET CONTENTS:

**ACCEPTED**

**05/22/2025**

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No.

**228-626**

<b>FIRST AID</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION/PRECAUCIÓN**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber.
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls**

When handlers use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

<b>USER SAFETY RECOMMENDATIONS</b>
Users should: <ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>

## ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

---

## DIRECTIONS FOR USE

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- Chemical-resistant footwear plus socks,
- Chemical-resistant headgear for overhead exposures

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides [40 CFR Part 170]. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**DO NOT** enter or allow others to enter the treated areas without protective clothing until spray has dried.

---

## COMMERCIAL TURF AND ORNAMENTALS

---

### PRODUCT INFORMATION

This product is formulated as a flowable liquid. It contains the active ingredient thiophanate-methyl, a broad spectrum fungicide which exhibits preventative, curative and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply with ground equipment using sufficient spray volume to provide thorough coverage.

[ALTERNATE PRODUCT INFORMATION STATEMENTS]

**[PRODUCT INFORMATION]**

*This product is formulated as a flowable liquid. It contains the active ingredient thiophanate-methyl, a broad spectrum fungicide which controls a variety of diseases on woody and herbaceous ornamental crops. This product is also effective as a preplant dip on cuttings and bulbs. Use this product to control the listed diseases on ornamental almond, apple, apricot, cherry, nectarine, peach, pecan, plum, and prune trees.*

*Apply this product with ground equipment using sufficient spray volume to provide thorough coverage. Add required amount of this product to partially filled tank agitated by mechanical or hydraulic means, and then add remaining required amount of water. Continuous agitation is required to keep the material in suspension. **DO NOT** tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules.]*

**Resistance Management:** For resistance management, please note that Nufarm T-Methyl SPC 4.5 F Fungicide contains thiophanate-methyl, a Group 1 fungicide. Any fungal population may contain individuals naturally resistant to Nufarm T-Methyl SPC 4.5 F Fungicide and other Group 1 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Nufarm T-Methyl SPC 4.5 F Fungicide or other Group 1 fungicides within a growing season sequence with different groups that control the same pathogens.
- Avoid application of more than the maximum number of applications listed in the directions for use and follow label instructions regarding sequential applications of Nufarm T-Methyl SPC 4.5 F Fungicide or other fungicides in the same groups in a season.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local Nufarm representative, retailer, or extension specialist for any additional pesticide resistance-management and/or IPM recommendations for specific plants and pathogens. You can also contact your pesticide distributor or university extension specialist to report resistance.

**Mixing Instructions:** SHAKE WELL BEFORE USING. Some settling may occur during prolonged periods of non-use. High pH environments can cause thiophanate-methyl, the active ingredient in this product, to break down and result in loss of efficacy. Buffer the tank water to pH 6-7 before adding this product for optimum product performance. Add required amount of product to partially filled tank (1/2 total volume), agitate by mechanical or hydraulic means, add tank mix product if used. **DO NOT tank mix this product with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur** (for more information, see ‘Tank Mixing Instructions’ below). Agitate again and then add remaining required amount of water. Maintain continuous agitation in the spray tank to keep the material in proper suspension. For best results, use spray mixture the same day it is prepared.

**Tank Mixing Instructions:** It is the pesticide user’s responsibility to ensure that all products included in the tank mix are registered for the intended use. Read and follow the applicable restrictions and limitations

and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

This product is compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: water soluble bags, wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. ***DO NOT tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur.*** This product may be applied in conjunction with chemically neutral liquid fertilizers. **DO NOT** apply in conjunction with highly alkaline fertilizers, such as aqueous ammonia, as they may cause degradation of the pesticide, resulting in reduced performance.

---

## APPLICATION THROUGH IRRIGATION SYSTEMS

---

Apply this product only through the following types of sprinkler systems: center pivot, lateral move, end tow, side wheel roll, traveler, solid set, hand move (or similar); or flood (basin) irrigation system. **DO NOT** apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact your State Extension Service specialists, equipment manufacturers, or other experts.

**DO NOT** connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments in case the need arises.

### Additional Application Instructions

Remove scale, pesticide residue, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

Determine the treatment rate as indicated in the directions for use for crop and pathogen and measure the intended areas of application.

Prepare a suspension of product in the mix tank or stock bucket. Fill the tank with 1/2 or 3/4 of the desired amount of water. Start agitation and add the required amount of product to the solution along with the remaining volume of water. Use sufficient water to ensure full coverage of foliage. **DO NOT** use an amount of water that could lead to excessive runoff from target plants. The amount of water will vary according to the amount of foliage requiring coverage and type of equipment but generally 25 to 100 gallons per acre is adequate.

Maintain a gentle agitation in the mix tank during application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time. An injection ration of 1:100 is required for greenhouse systems.

Pesticide application must begin as soon as irrigation pattern is established and stabilized. Irrigation must continue after application of pesticide for a sufficient amount of time for pesticide to be flushed from end of irrigation system.

### **Specific Information for Irrigation Systems Connected to a Public Water Supply**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer, or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump or equivalent, effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

### **Specific Information for Flood (Basin), Furrow and Border Chemigation**

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.



### Specific Information for Applications through Sprinkler Irrigation Systems

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

---

## HORTICULTURAL APPLICATIONS

---

### Nursery, Greenhouse, Landscape & Interiorscape

#### Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs

#### Restriction

Except as allowed under “Backyard Fruit and Nut Trees”, **DO NOT** use fruit, nuts or sap from treated trees as food or feed.

This product is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. It is also effective as a pre-plant dip on cuttings and bulbs.

**Note:** The “Directions For Use” of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product may be phytotoxic to Swedish Ivy (*Plectranthus australis*), Boston Fern (*Nephrolepis exalta*), and Easter Cactus (*Hatiora gaertneri*).

**Application Instructions:** Apply material with properly calibrated, hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation, flood, or drip systems, or as an overhead application where applicable. See specific instructions below.

For soil drench applications, best crop protection is achieved with preventative treatments repeated every 21-28 days.

For foliar applications, begin applications when disease first appears and repeat at 7-14 day intervals or as needed during the growing season. Use the shorter interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix 8-24 fl oz of this product (0.28-0.84 lb a.i.) per 100 gallons water (0.5-1.5 teaspoons per gal) and apply as a full coverage spray to

drip for the prevention and control of the diseases listed below. Spray volume may range up to 400 gallons of finished spray per acre depending upon plant species and plant growth stage.

For applications through irrigations systems, refer to use rates indicated in the foliar application chart.

For small volume applications less than 100 gallons, divide the specified rate by 16 to get the number of teaspoons of this product per gallon.

**Adjuvants:** Where rainfall and/or overhead irrigation is the norm, use of a compatible spreader/sticker may enhance product performance. Where wetting of foliage is difficult, use a compatible wetting agent. Refer to the Tank Mixing Instructions section of this label for more information on tank mixing and compatibility.

### **Special Instructions for Proportional Injectors (e.g. Dosatron, Dosmatic, Anderson, and similar equipment)**

Determine the treatment rate for crop and pathogen from the foliar application table below. Determine the injection ratio for the individual system to be used for application. For systems using a 1:100 ratio, measure and add the specified amount of material per 100 gallons to each gallon of water in a stock bucket or tank. For systems using a 1:200 ratio, multiply the specified amount per 100 gallons by 2. For systems using a 1:50 ratio, divide the specified amount per 100 gallons by 2. For systems using 1:16 ratio, divide the specified amount per 100 gallons by 6. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time. An injection ratio of 1:100 is preferred for most greenhouse and nursery systems.

## **FOLIAR APPLICATION**

### **Restrictions**

For all ornamentals except cut flowers, **DO NOT** exceed a maximum single application rate of 85.3 fl oz product/acre (3.0 lb a.i./acre).

For cut flowers, **DO NOT** exceed a maximum single application rate of 14.2 fl oz product/acre (0.5 lb a.i./acre).

For all foliar applications to ornamentals, **DO NOT** apply more than 66.7 gallons product/acre per year (300 lb a.i./acre/year).

<b>Disease(s) Controlled</b>	<b>Application Rate fl oz Product/100 gal</b>	<b>Application Instructions</b>
<b>Anthracnose</b> <i>Colletotrichum</i>	12-20 (0.42-0.70 lb a.i.)	Apply as buds break or at first sign of disease. Repeat at 7-14 day intervals as needed during disease period.
<b>Black Spot of Rose</b> <i>Diplocarpon rosae</i>	12-20 (0.42-0.70 lb a.i.)	Apply early summer or at first sign of disease. Repeat every 7-14 days as needed during disease period.
<b>Brown Rot and Blight</b> <i>Monilinia, Sclerotinia, Whetzellinia</i>	12-20 (0.42-0.70 lb a.i.)	Apply late spring or at first sign of disease. Repeat every 7-14 days as needed during the disease period.

<b>Disease(s) Controlled</b>	<b>Application Rate fl oz Product/100 gal</b>	<b>Application Instructions</b>
<b>Fusicladium and Venturia Leaf Scabs on:</b> Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	12-20 (0.42-0.70 lb a.i.)	Apply as buds break. Repeat every 7-14 days during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with mancozeb (Protect™), chlorothalonil, or propiconazole can be utilized.  <b>Restriction: DO NOT</b> use fruit from treated crabapple or pear trees for food purposes.
<b>Leaf Spots and Blights caused by:</b> <i>Ascochyta, Blumeriella, Botrytis, Cercospora, Cocomyces, Corynespora, Curvularia, Didymellina, Entomosporium, Fabraea, Fusarium, Ramularia, Rhizoctonia, Marssoninia, Mycosphaerella, Myrothecium, Phoma, Physalasporea, Schizothyrium, Septoria, Sphaceloma</i>	12-20 (0.42-0.70 lb a.i.)	Apply when disease symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil can be utilized.
<b>Ovulinia Blight</b>	8-16 (0.28-0.56 lb a.i.)	Apply as flowers open. Repeat every 7-14 days during disease period.
<b>Powdery Mildews</b> <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i>	12-20 (0.42-0.70 lb a.i.)	Apply when disease first appears and repeat every 7-14 days during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or triadimefon can be utilized.
<b>Rust Diseases caused by:</b> <i>Puccinia, Gymnosporangium, Uromyces</i>	12-16 (0.42-0.56 lb a.i.)	Apply late spring or when symptoms first appear. Repeat every 7-14 days as needed during disease period. Rotations and/or tank mix combinations with mancozeb (Protect) or chlorothalonil are advised.
<b>Tip Blight of Pine</b> <i>Sphaeropsis sapinea, Diplodia pinea</i>	16-20 (0.56-0.70 lb a.i.)	Begin applications in the spring when new growth starts. Make a second application just prior to needle emergence from the sheath and a third application 14 days later. Thorough coverage is essential for optimal disease control.

Disease(s) Controlled	Application Rate fl oz Product/100 gal	Application Instructions
<b>Twig Blights, Cankers, and Diebacks</b> <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	16-20 (0.56-0.70 lb a.i.)	Apply when symptoms first appear. Repeat every 7-14 days as needed during disease period.

### Backyard Fruit and Nut Trees (Not for use in California)

Certified applicators:

If this product is used on a tree producing fruits or nuts which will be used for food or feed:

- **DO NOT** apply to home orchards/backyard fruit or nut trees after fruit or nut set.
- **DO NOT** apply to fruit or nut trees other than almond, apple, pear, pecan, pistachio, apricot, cherry, nectarine, peach, plum or prune.
- For bearing fruit and nut trees, use the following application rates (for a list of diseases controlled see the table above for Foliar Application):

Crop	Application Rate Fluid Ounces Product/Acre	Maximum Annual Rate Fluid Ounces Product/Acre
Almond, Apple, Pear, Pecan, Apricot, Cherry, Nectarine, Peach, Plum or Prune	19.5 (0.7 lb a.i./A)	59 (2.1 lb a.i./A)
Pistachio	19.5 (0.7 lb a.i./A)	39 (1.4 lb a.i./A)

## Vegetable Transplants (Greenhouse and Nursery Use Only):

Not for field vegetable production

Vegetable Transplants	Diseases Controlled	Rate of Product	Application Instructions
<b>Beans, dry and succulent including:</b> Lima bean, Snap bean, Kidney bean, Mung bean, Navy bean, Pinto bean, Wax bean, Broad bean, Fava bean, Asparagus bean, Blackeyed pea, Cowpea, Sweet lupine, White lupine, White sweet lupine, Grain lupine, Chick pea, Garbanzo bean	<b>Anthracnose</b> <i>Colletotrichum</i>	29-39 fl oz/Acre <sup>1</sup> (1.02-1.37 lb a.i./A)	For one application: Apply when 100% of plants have at least one open bloom or when conditions are favorable for disease development.
	<b>Gray Mold</b> <i>Botrytis</i>	or	or
	<b>White Mold</b> <i>Sclerotinia</i>	19-29 fl oz/Acre <sup>2</sup> (0.67-1.02 lb a.i./A)	For multiple applications: Make the first application when 10% to 30% of plants have at least one open bloom and follow with sequential applications on a 4 to 7 day interval. Apply prior to the development of disease for best results.
<ul style="list-style-type: none"> <li>♦ <b>DO NOT</b> apply more than 79 fl oz of product (2.8 lb a.i.)/A/year.</li> <li>♦ <b>Pre-harvest interval:</b>  <b>California only</b> - 14 days for succulent beans, 28 days for dry beans and lima beans.  <b>All other states</b> - 14 days for succulent beans and lima beans, 28 days for dry beans.</li> </ul>			
<b>CUCURBITS</b> (Cantaloupe, Casaba, Cucumbers, Melons, Pumpkins, Summer and Winter Squash, and Watermelons)	<b>Anthracnose*</b> <i>Colletotrichum</i>	10 fl oz/Acre <sup>3</sup> (0.35 lb a.i./A)	Begin applications when plants begin to run or when disease first appears, and repeat at 7 to 14 day intervals.
	<b>Gummy Stem Blight*</b> <i>Didymella</i>		For Target Spot use at 7 day intervals.
	<b>Powdery Mildew*</b> <i>Erysiphe,</i> <i>Sphaerotheca,</i> <i>Podosphaera</i>		
	<b>Target Spot*</b> <i>Corynespora</i>		
	<b>Belly Rots*</b> <i>Rhizoctonia,</i> <i>Fusarium</i>	10 fl oz/Acre <sup>3</sup> (0.35 lb a.i./A)	Apply in sufficient volume to allow runoff to the soil. Will not control <i>Pythium</i> or <i>Phytophthora</i> .
<ul style="list-style-type: none"> <li>♦ <b>DO NOT</b> apply more than 59 fl oz of product (2.1 lb a.i.)/A/year from any combination of application timings.</li> <li>♦ Follow resistance management guidelines under Directions for Use.</li> </ul>			

\* Not for this use in California.

<sup>1</sup> Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 29 fl oz per acre (1.02 lb a.i./A). If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 39 fl oz per acre (1.37 lb a.i./A). For example, if 200 gallons of water are required, use 19 fl oz (0.67 lb a.i.) per 100 gallons.

- <sup>2</sup> Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 19 fl oz per acre (0.67 lb a.i./A). If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 29 fl oz per acre (1.02 lb a.i./A). For example, if 200 gallons of water are required, use 14.5 fl oz (0.51 lb a.i.) per 100 gallons.
- <sup>3</sup> Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 10 fl oz per acre (0.35 lb a.i./A). If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 10 fl oz per acre (0.35 lb a.i./A). For example, if 200 gallons of water are required, use 5 fl oz (0.18 lb a.i.) per 100 gallons.

## SOIL DRENCH APPLICATION

### Restriction

**DO NOT** apply more than 66.7 gallons product/acre per year (300 lb a.i./acre/year).

Diseases Controlled	Rate of Product	Application Instructions
<b>Stem, Crown, and Root Rots caused by:</b> <i>Botrytis, Cythroccladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia</i> <b>Black Root Rot</b> <i>Thielaviopsis</i>	12-20 fl oz/100 gal (0.42-0.70 lb a.i.)	Apply as a drench or directed spray using hand held, mechanical, or motorized spray equipment, or as a chemigation drench or directed spray using applicable sprinkler irrigation systems, after seeding or sticking of cuttings (8 fl oz (0.28 lb a.i.)) or after transplanting (12-16 fl oz (0.42-0.56 lb a.i.)) to propagation beds, containers, pots, trays, or nursery or landscape beds at a rate to thoroughly soak the growing media through the root zone. A general guide is 0.25-3.0 pints of finished mixture per sq ft depending on the media type and depth (about 4 fl oz (0.14 lb a.i.) per 4 inch pot or 8 fl oz (0.28 lb a.i.) per 6 inch pot). Repeat every 21-28 days for adequate crop protection. <b>Note:</b> This product does not control <i>Pythium</i> or <i>Phytophthora</i> . Tank mix combinations with metalaxyl, mefenoxam, etridiazole, fosetyl-Al, or mono and dipotassium salts of phosphorous acid or propamocarb are required for the control of <i>Pythium</i> and <i>Phytophthora</i> .

## PLANT DIP APPLICATION

Diseases Controlled	Product Use Rate	Application Instructions
<b>Plant or Cutting Diseases caused by:</b> <i>Botrytis, Cylindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	16-20 fl oz/100 gal (0.56-0.70 lb a.i.)	Immerse plants or cuttings for 10-15 min. Remove and allow to drain. <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.
<b>Bulb, Corm, and Rhizome Rots caused by:</b> <i>Botrytis, Cylindrocladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis</i>	16-20 fl oz/100 gal (0.56-0.70 lb a.i.)	Soak cleaned bulbs for 15-30 min in warm solution (80-85°F). For storage disease prevention, treat bulbs preferably within 48 hours after digging. After treatment, dry well before storing. If bulbs are for forcing, treat bulbs that have been heat-cured. <b>Note:</b> Follow accepted hygiene practices to minimize the introduction and spread of water born bacterial and water mold fungal diseases.

---

## TURF APPLICATIONS

---

For use on all fine turf applications such as Bentgrasses, Bermuda grasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia or their mixtures. This product is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. Use this product for the prevention and control of the diseases listed below.

For best results use spray mixture the same day it is prepared. Spray uniformly over the area to be treated with a properly calibrated sprayer. Apply after mowing. **DO NOT** mow for twelve hours after application. Apply specified amounts in sufficient water to obtain thorough coverage (2-4 gallons per 1,000 square feet suggested). When treating golf greens, always treat aprons.

**Application Instructions:** Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation systems. Spray uniformly over the area to be treated. Apply specified amounts in sufficient water to obtain thorough coverage of treatment area (2-4 gal per 1,000 sq ft is suggested). When treating golf greens, always treat aprons. Use the higher specified rate under conditions of severe disease pressure. For best results, apply after mowing. **DO NOT** mow for twelve hours after application. For root pathogens, lightly water the treatment area to move the fungicide into the active root zone with one to two tenths inch of water. Excessive irrigation may move application below active root zone and reduce application effectiveness. Green design and drainage will influence irrigation practices. When tank mixing with contact action fungicides for foliar diseases, allow applications to dry on leaf surfaces. Normal watering may proceed after sprays have dried.

### Restrictions

#### Not for homeowner use.

For use only by certified applicators or those under their immediate supervision.

**DO NOT** use on turf being grown for sale or other commercial use as sod.

**DO NOT** graze animals on treated turf.

**DO NOT** feed clippings to livestock or poultry.

**DO NOT** apply with fixed wing or rotary aircraft.

Minimum 14-day Retreatment Interval.

**Maximum Individual Application Rates and Minimum Retreatment Intervals**

**DO NOT** exceed the amounts per acre or reduce the retreatment interval indicated below.

Use Site	Maximum Application rate of Product	Minimum Retreatment Interval	Restrictions
Residential or Public Areas	1 $\frac{3}{4}$ fl oz / 1,000 sq ft (2.7 lb a.i./A)	14 days	
Golf Course Tees, Greens, Aprons	5 $\frac{1}{3}$ fl oz / 1,000 sq ft (8.2 lb a.i./A)	14 days	
Golf Course Fairways – except Florida	3 $\frac{1}{2}$ fl oz / 1,000 sq ft (5.5 lb a.i./A)	14 days	Excludes Florida
Golf Course Fairways – Florida Only	1 $\frac{3}{4}$ fl oz / 1,000 sq ft (2.7 lb a.i./A)	14 days	Florida Only

**Maximum Annual Application Rates**

**DO NOT** exceed the following amounts of product per acre per year.

Use Site	Maximum Gallons of Product per Acre per Year	Fluid Ounces Product per 1,000 sq ft	Restrictions
Residential or Public Areas	2.4 (10.9 lb a.i./A)	7 (0.25 lb a.i./1000 ft <sup>2</sup> )	4 Applications per year.
Golf Course Tees, Greens, Aprons	4.8 (21.8 lb a.i./A)	14 $\frac{1}{4}$ (0.50 lb a.i. /1000 ft <sup>2</sup> )	8 Applications per year.
Golf Course Fairways – except Florida	1.2 (5.5 lb a.i./A)	3 $\frac{1}{2}$ (0.12 lb a.i. /1000 ft <sup>2</sup> )	2 Applications per year. Excludes Florida
Golf Course Fairways – Florida Only	0.6 (2.7 lb a.i./A)	1 $\frac{3}{4}$ (0.06 lb a.i. /1000 ft <sup>2</sup> )	1 Application per year. Florida Only

**Turf Disease Control**

Disease(s) Controlled	Product Use Rate fl oz/1,000 sq ft	Application Instructions
<b>Anthracnose, basal</b> <i>Colletotrichum cereale</i> <b>Anthracnose, foliar</b> <i>Colletotrichum cereale</i>	3 $\frac{1}{2}$ - 5 $\frac{1}{3}$ (0.12-0.19 lb a.i.)  1 $\frac{3}{4}$ - 3 $\frac{1}{2}$ (0.06-0.12 lb a.i.)	For prevention in historic areas of disease pressure, apply twice at 14 day intervals when soil temperature reaches 60°F. For curative control, apply when disease first appears and continue at 14 day intervals. Rotations and/or tank mix combinations with chlorothalonil or triadimefon can be utilized.



<b>Disease(s) Controlled</b>	<b>Product Use Rate fl oz/1,000 sq ft</b>	<b>Application Instructions</b>
<b>Bermudagrass Decline</b> <i>Gaeumannomyces graminis</i> var. <i>graminis</i>  <b>Take-All-Patch</b> <i>Gaeumannomyces graminis</i> var. <i>avenae</i>	3 ½ - 5 ⅓ (0.12-0.19 lb a.i.)	Apply in mid-July or when disease symptoms first appear and repeat at 14 day intervals for suppression. Use higher rates under most severe disease expression. Water treatment into active root zone. Follow proper agronomic directions to maintain plant vigor.
<b>Cool Season Brown Patch</b> <i>Rhizoctonia cerealis</i> <b>Necrotic Ring Spot</b> <i>Leptosphaeria korrea</i> <b>Spring Dead Spot</b> <i>Leptosphaeria korrea</i>	3 ½ - 5 ⅓ (0.12-0.19 lb a.i.)	For prevention, apply in Fall before turf has stopped all growth activity. Apply second application in early Spring when soil temperatures reach 55-60°F or when disease first appears. For curative action, apply when disease first appears in early Spring and continue at 14 day intervals. Water treatment into active root zone.
<b>Coprinus Snow Mold</b> <i>Coprinus psychromorbidus</i>	3 ½ - 5 ⅓ (0.12-0.19 lb a.i.)	Apply 2 treatments at 21 day intervals in late Fall to early Winter, with the last application made just prior to first permanent snow cover. Rotations and/or tank mix combinations with PCNB can be utilized.
<b>Dollar Spot</b> <i>Moellerodiscus</i> , <i>Lanzia</i> , <i>Sclerotinia</i> <b>Large Brown Patch</b> <i>Rhizoctonia solani</i> <b>Ascochyta Leaf Blight</b> <i>Ascochyta</i> <b>Copper Spot</b> <i>Gloeocercospora sorghi</i> <b>Fusarium Patch</b> <i>Fusarium nivale</i> <b>Red Thread</b> <i>Laetisaria fuciformis</i> <b>Zoysia Patch</b> <i>Rhizoctonia solani</i>	1 ¾ - 3 ½ (0.06-0.12 lb a.i.)	Apply when disease first appears and continue at 14 day intervals. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or mancozeb (Protect) can be utilized.
<b>Fusarium Blight</b> <i>Fusarium roseum</i> , <i>F. triticum</i>	3 ½ - 5 ⅓ (0.12-0.19 lb a.i.)	Apply when disease first appears at 14 day intervals.
<b>Gray Leaf Spot (Blast)</b> <i>Pyricularia grisea</i>	3 ½ - 5 ⅓ (0.12-0.19 lb a.i.)	Apply preventative application before expected period of disease development. Continue applications at 14 day intervals.
<b>Leaf Spot</b> <i>Drechslera</i> <b>Leaf, Crown, and Root Diseases</b> <i>Bipolaris</i> , <i>Curvularia</i> , <i>Exserohilum</i>	3 ½ - 5 ⅓ (0.12-0.19 lb a.i.)	Apply when disease first appears and make applications at 14 day intervals as needed. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or mancozeb (Protect) can be utilized under severe conditions.

Disease(s) Controlled	Product Use Rate fl oz/1,000 sq ft	Application Instructions
<b>Pink Snow Mold</b> <i>Microdochium nivale</i>	3 ½ - 5 ½ (0.12-0.19 lb a.i.)	Apply in late Fall to early Winter before turf has stopped all growth activity. A second application may be used in combination with chlorothalonil, PCNB, or thiram (Spotrete™) at specified rates before snow cover or during Spring thaw.
<b>Rusts</b> <i>Puccinia, Uromyces</i>	3 ½ - 5 ½ (0.12-0.19 lb a.i.)	Apply at 14 day intervals when disease first appears. Rotations and/or tank mix combinations with chlorothalonil or mancozeb (Protect) are specified.
<b>Stripe Smut</b> <i>Ustilago striiformis</i>	3 ½ - 5 ½ (0.12-0.19 lb a.i.)	Apply at 14 day intervals when disease first appears. For prevention, apply in spring and fall.
<b>Summer Patch</b> <i>Magnaporthe poae</i>	3 ½ - 5 ½ (0.12-0.19 lb a.i.)	For prevention, apply 3 applications starting late April or early May using 21 day intervals. Rotations and/or tank mix combinations may be used as part of the three application program. For suppression, apply at 14 day intervals when disease first appears. Water treatment into active root zone.
<b>Bentgrass Dead Spot</b> <i>Ophiosphaerella agrostis</i>	3 ½ - 5 ½ (0.12-0.19 lb a.i.)	For prevention, apply in early June or based upon local Extension Service directions. Apply at 14 day intervals. Rotations and/or tank mix combinations may be used for season long disease prevention.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container in a dry, temperature controlled area.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

#### [Nonrefillable Containers 5 Gallons or Less:]

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### [Nonrefillable containers larger than 5 gallons:]

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### **WARRANTY DISCLAIMER**

**THE DIRECTIONS FOR USE OF THIS PRODUCT MUST BE FOLLOWED CAREFULLY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.**

### **LIMITATION OF LIABILITY**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.**

**If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RVxxxxxx)

Protect and Spotrete are trademarks of Cleary Chemicals, LLC

[LESCO and T-STORM are trademarks of LESCO Technologies, LLC]

All other trademarks are the property of their respective owners.