

228-684

04/26/2011

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Nufarm Americas, Inc.
% Georgia Anastasiou
Lewis & Harrison
122 C Street N.W., Suite 740
Washington, DC 20001

APR 26 2011

Product Name: Nufarm Iprodione SPC Fungicide
EPA Reg. No.: 228-684
Subject: Your amendment dated February 14, 2011
EPA Decision Number: 445167

Dear Ms Anastasiou:

The amended label referred to above, originally submitted July 16, 2010 in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable.

One copy of the label stamped "Accepted" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure

2/15

Nufarm Iprodione SPC Fungicide

A broad-spectrum fungicide for non-residential use on turf and ornamentals


ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide* 23.3%

OTHER INGREDIENTS: 76.7%

TOTAL: 100.0%

This product contains petroleum distillate.
*Contains 2 lbs. Iprodione per gallon.



KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

See inside label booklet for additional **FIRST AID** and **PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-684
EPA EST. NO. _____

MANUFACTURED FOR
NUFARM AMERICAS INC.
150 HARVESTER DRIVE
BURR RIDGE, IL 60527
800-455-2000



Product of France or China.
Formulated in the U.S.A. with
U.S. and imported ingredients

NET CONTENTS: _____ (Gal.) (_____ liters)

Nufarm Iprodione SPC
Fungicide contains iprodione,
the active ingredient used in
Chipco® 26 GT.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION / PRECAUCIÓN**

Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes or clothing.

FIRST-AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Contains petroleum distillate – vomiting may cause aspiration pneumonia.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, others exposed to the concentrate, cleaners/repairers of equipment and applicators applying as a dip treatment must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber, neoprene rubber or viton
- Chemical-resistant apron
- Chemical-resistant footwear plus socks

Applicators using hand held equipment must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber, neoprene rubber or viton
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Dust/mist filtering respirator (NIOSH approved respirator with any R, P or HE filter)

Applicators using aircraft or mechanical ground equipment (groundboom, airblast, etc.) and flaggers for aerial applications must wear:

- Long-sleeve shirt and long pants
- Shoes plus socks

Applicators using truck-mounted equipment with a handgun at the end of a hose (i.e., for commercial turfgrass or ornamental applications) and all other handlers not specified above must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber, neoprene rubber or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical can contaminate surface water through aerial and ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

This pesticide is toxic to invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas is hazardous to aquatic invertebrates in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours for ornamental uses. The restricted entry interval for all other WPS uses is 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils
- Shoes and socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to ornamental and turf uses (golf courses, landscape and institutional areas) of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

GENERAL INFORMATION

This product is a broad-spectrum fungicide that is applied as either a foliar spray, drench or dip and controls turfgrass diseases in non-residential sites such as golf courses, sod farms, and institutional areas where fine turf is grown, as well as a wide range of ornamental flowering and foliage plants in field, landscape and conifer nursery sites. This product is effective in controlling the following diseases:

Spring, Summer and Fall Turf Diseases:

- Brown Patch
- Leaf Spots like Helminthosporium Leaf Spot caused by *Dreschlera* spp. Pathogens
- Necrotic Ring Spot
- Corticum Red Thread
- Fusarium Blight Large Patch
- Dollar Spot

Winter Turf Diseases:

- Fusarium patch (Pink snow mold)
- Gray snow mold

Ornamental and Nursery Diseases:

- Aerial web blight
- Alternaria leaf spot
- Fusarium leaf spot
- Rhizoctonia stem and root rot
- Tulip fire
- Ray blight
- Daffodil leaf scorch
- Botrytis storage rot
- Alternaria leaf blight
- Botrytis blight
- Helminthosporium leaf spot
- Ink spot
- Alternaria leaf blight
- Fusarium corm rot
- Blossom blight
- Cylindrocladium blight and wilt

USE PRECAUTIONS AND RESTRICTIONS

- For best results, be sure to follow all the precautions, limitations and instructions in this label.
- Use of this product at residential sites is prohibited.
- Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.
- For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.
- Do not apply this product when the wind direction is toward aquatic areas.

GENERAL APPLICATION GUIDELINES

- Apply the rates indicated in the following sections of the label in 0.5 - 10 gallons of water per 1000 square feet.

- Do not drench the foliage to the point of runoff.
- Product breakdown could occur if the spray mixture is allowed to stand for more than 12 hours.
- Maintain agitation during spray operations.
- Always apply using a properly calibrated sprayer.

TURF

GENERAL INSTRUCTIONS FOR TURF:

Unless otherwise noted, make applications when the disease first appears or when conditions favor disease development.

Under severe conditions, use the higher rate and/or shorter interval of applications for all diseases. When disease pressure is light to moderate, use the lower rates and longer intervals.

USE PRECAUTIONS:

- DO NOT apply more than 35 fl. oz. of this product per 1000 square feet per year (24 lbs. a.i. per acre).
- DO NOT make more than 6 applications to a single site per year.
- DO NOT mix with any sticker, extender, or wetting agent.
- DO NOT mow or irrigate treated areas until the foliage is completely dry. Wait 24 hours following treatment.
- DO NOT graze animals on treated turf, and do not feed clippings from treated turf to livestock or poultry.

TARGET PEST	RATE (fl. oz. / 1000 sq. ft.)	APPLICATION INTERVAL
Dollar spot <i>(Lanzia spp. and Moellerodiscus spp.)</i> Brown patch <i>(Rhizoctonia solanii)</i> Leaf spot <i>(Drechslera spp.)</i>	<p>3 to 4</p> <p>For dollar spot control on fairways use 2 – 4</p>	<p>Greens and Tees: Repeat at 30-day intervals as long as required.</p> <p>Fairways and Other Turf Areas: Repeat at 30-day intervals as long as required.</p>
Large patch [†] <i>(Rhizoctonia solanii)</i>	4	<p>Make first application in fall when conditions are favorable for disease development but no symptoms are visible. Repeat applications every 30 days in spring as needed.</p>
Fusarium blight <i>(Fusarium spp.)</i> Necrotic ring spot [†] <i>(Leptosphaeria korrae)</i>	8	<p>Use only preventative foliar applications when conditions first become favorable for disease development. Make additional applications at 30-day intervals as necessary.</p>
Fusarium patch <i>(Microdochium nivalis)</i> [Pacific Northwest Only – West of the Cascade Mountains]	4 to 8	<p>Repeat at 30-day intervals as long as required.</p>
Gray snow mold <i>(Typhula spp.)</i> Pink snow mold <i>(Fusarium nivale)</i>	<p>4 to 8</p> <p>(See tank mixes for additional information)</p>	<p>Make one application before first permanent snow cover and a second during a mid-winter thaw.</p>
Corticium red thread <i>(Laetisaria fuciformis)</i>	4	<p>Apply every 30 days as required for prevention.</p>
Curvularia <i>(Curvularia spp.)</i> on Bermudagrass only	4	<p>Apply every 30 days as required for prevention.</p>
Anthracnose <i>(Colletotrichum)</i> NOTE: suppression only	4 to 8	<p>Combine this product with appropriately labeled and registered trifloxystrobin or fosetyl-al products or other anthracnose control fungicides.</p>
Pythium blight	See tank mixes below	

[†] Not registered for use in California

TANK MIXTURES FOR TURF APPLICATIONS

To expand the spectrum of pests controlled, Tank mix this product with most commonly used fungicides containing flutolanil, trifloxystrobin, and azoxystrobin. When tank mixing products, be sure to follow the most restrictive instructions.

Broad Spectrum Disease Control and Resistance Management:

Tank mixing this product with an appropriately labeled and registered thiophanate-methyl product provides effective, broad spectrum turf disease control and also serves as a useful tank mixture in the resistance management program required for other resistance sensitive fungicides.

Disease Pressure	Product Rate	Thiophanate-methyl Rate
Low to Medium	3 fl. oz./1000 ft. ²	1.0 fl. oz./1000 ft. ²
High	3 fl. oz./1000 ft. ²	2.0 fl. oz./1000 ft. ²

Summer Stress Complex/Summer Decline:

Mix 2 to 4 oz. of this product with 4 to 8 oz. of an appropriately labeled and registered fosetyl-al containing product per 1000 square feet.

Pythium Blight:

Pythium blight will be controlled by the tank mixing of fosetyl-al, or propamocarb hydrochloride with this product. If using a tank mixture, follow label directions for the use of that product and apply at the rate specified for control of the target disease organism.

Gray Snow Mold:

In areas where continuous snow cover occurs, use 4to8 fl. oz. of this product per 1000 sq. ft. tank mixed with an appropriately labeled and registered chlorothalonil or pentachloronitrobenzene (PCNB) product at the labeled rate.

Make applications in the fall before snow cover occurs and use the higher rates listed if the turf remains frozen prior to snow cover. Apply with 1 to 5 gallons of spray solution per 1000 square feet. For best results, reapply if loss of snow cover occurs during a winter thaw.

8/15

ORNAMENTALS

FOR USE BY COMMERCIAL NURSERY AND LANDSCAPE PERSONNEL ONLY. NOT FOR RESIDENTIAL AREAS.

The ornamentals listed below have been tested and found to be tolerant to this product. As it is not possible to test every species or variety of ornamental plant for tolerance, the user should test for phytotoxic responses in plants not listed in this label prior to widespread application.

This product has been tested on the following ornamentals:

Ageratum	Ajuga	Almond (ornamental)	Alyssum
Andromeda	Aphelandra	Artemisia	Aster
Azalea	Boxwood	Cactus	Calendula
Carnation	Cherry (ornamental)	Chrysanthemum	Cineraria
Cistena Plum	Coleus	Columbine	Coral Bells (Heuchera)
Crape Myrtle	Crassula	Croton	Cyclamen
Daffodilis	Dahlia	Delphinium	Deutzia
Dianthus	Dieffenbachia	Dizygotheca	Dogwood
Dracena	English Ivy	Episcia	Euonymous
Ficus	Forsythia	Gazania	Geranium
Gladiolus	Gloxinia	Gypsophila	Hawthorn
Holly	Hoya	Hydrangea	Impatiens
Iris	Juniper	Kalanchoe	Lillies
Lipstick vine	Marigold	Monarda (Bee Balm)	Pachysandra
Palm	Pansy	Peach (ornamental)	Peperomia
Periwinkle	Philodendron	Phlox	Pilea
Pine	Pitosporum	Plum (ornamental)	Poinsettia
Poppy	Pothos	Primrose	Privet
Protea	Pyracantha	Rhododendron	Rose
Rose Tree of China	Salvia	Schefflera	Snapdragon
Statice	Tree Ivy	Tulip	Viburnum
Violet	Zinnia		

NOTE: DO NOT apply this product to Peace Lilly or White Anthurium (*Spathiphyllum*).

Use the following table to determine the diseases controlled and the application method to use:

Disease	Can Be Applied To	Foliar Spray	Drench	Dip
Aerial web blight	All	<input type="checkbox"/>		
Alternaria leaf blight	All	<input type="checkbox"/>		
Alternaria leaf spot	All	<input type="checkbox"/>		
Botrytis blight	All	<input type="checkbox"/>		
Fusarium leaf spot	All	<input type="checkbox"/>		
Helminthosporium leaf spot	All	<input type="checkbox"/>		
Rhizoctonia stem and root rot	All except Impatiens and Pothos		<input type="checkbox"/>	
Ink spot	Iris	<input type="checkbox"/>		
Tulip fire	Tulip	<input type="checkbox"/>		
Alternaria leaf blight	Zinnia	<input type="checkbox"/>		
Ray blight	Chrysanthemum	<input type="checkbox"/>		
Fusarium corm rot	Gladiolus			<input type="checkbox"/>
Daffodil leaf scorch	Daffodilis	<input type="checkbox"/>		
Blossom blight	Cistena Plum / Ornamental Plum	<input type="checkbox"/>		
Botrytis storage rot	Rose			<input type="checkbox"/>
Cylindrocladium blight and wilt	Azalea and Rhododendron			<input type="checkbox"/>

FOLIAR SPRAY APPLICATIONS

Apply when conditions are favorable for disease development using the following instructions:

- Application Rate: 1.0 to 2.5 quarts of product per acre
For severe pest pressure, use the highest specified rates.
For light to moderate pest pressure, use the lower rates listed.
- Application Interval: 7 to 14 days
For severe pest pressure, use the shortest application intervals.
For light to moderate pest pressure, use the longer application intervals.
- Application Instructions: Spray plants ensuring complete coverage.
- Use Precautions: DO NOT make more than 4 applications per crop per year.

DRENCH APPLICATIONS

To control Rhizoctonia, apply this product as a drench at the seeding and/or transplanting stage using the following instructions:

- Application Rate: 13 fl. oz. per 100 gallons.
- Application Interval: 14 days
- Application Instructions: Apply using 1 to 2 pints of solution per square foot.
For severe disease pressure use the higher rates.
For light to moderate disease pressure use the lower rates.
- Use Precautions: DO NOT apply more than 35 fl. oz. / 1000 sq. ft. per year (24 lbs. a.i. per acre).
DO NOT make more than 6 applications per year.
DO NOT use this product as a drench on Impatiens and Pothos.

DIP APPLICATIONS

Refer to the following table for use of this product as a dip to control Botrytis Storage Rot, Cylindrocladium Blight and Fusarium Corn Rot in the following plants:

Plant Species	Application Rate (Quarts Product / 100 Gal Water)	Dip Duration	Instructions
Rose	1.0	5 minutes	Dip bare root roses prior to cold storage.
Azalea and Rhododendron			Dip cuttings prior to planting.
Gladiolus	2.0		Dip corms prior to storage.

TANK MIXTURES

In order to broaden the spectrum of control, use this product with most commonly used fungicides. For control of diseases caused by *Pythium* and *Phytophthora* spp., use a tank mix of this product with fosetyl-al.

Read the labels of all tank mix partners for specific application rates for the target disease organism and be sure to follow the most restrictive instructions.

DIRECTIONS FOR USE THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product using a center pivot irrigation system using the following instructions:

- System Preparation: Be sure all pesticide residues, scale and other foreign materials are cleaned from the chemical tank and injector system. Flush with clean water prior to use.

Prepare a tank mix of this product by filling the tank to 1/2 to 3/4 of the final volume with water and begin agitation. Add the specified amount of this product and the remaining water until the desired volume is reached.

Application Rate: Use the specified dosage per acre per 1 to 4 gallons of water.

Application Instructions: Set the sprinkler system to deliver 0.1 to 0.3 inches of water per acre. Using a positive displacement pump, this product mixture should be injected into the main line ahead of a right angle turn to ensure adequate mixing.

Use Precautions: **Application of this product using a sprinkler system is prohibited in the state of California.**

Apply this product using only a center pivot irrigation system. Do not apply this product through any other type of irrigation system.

To prevent this product from being washed off the crop, do not irrigate the treated area for 24 hours after making the application.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop is a result from non-uniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

SPRAY DRIFT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

- 1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

2. Nozzles must always point backwards parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information** section below.

Aerial Drift Reduction Advisory Information:

(This section is advisory in nature and does not supercede the mandatory label requirements)

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated

cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

[Nonrefillable Containers 5 Gallons or Less:]

CONTAINER DISPOSAL: ~~Nonrefillable container. Do not reuse or refill this container.~~ Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[Nonrefillable containers larger than 5 Gallons:]

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration. Do not burn unless allowed by state and local ordinance. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable containers]

CONTAINER HANDLING [DISPOSAL]: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

[Refillable For return to Nufarm:]

CONTAINER HANDLING [DISPOSAL]: Refillable Container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you DO NOT agree with or DO NOT accept any of directions for use, the warranty disclaimers, or limitations on liability, DO NOT use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV071610N)

Nufarm Iprodione SPC Fungicide is not manufactured or distributed by Bayer Environmental Science, seller of Chipco® 26GT®.

Chipco® and 26 GT® are registered trademarks of Bayer.

LABEL HISTORY

File Name	Revision Mark	Comment
	RV090715N	Submission of primary brand name and minor label (replace product name with "this product", replace "recommend" and "may" with more definitive language, updated Container Disposal section and the warranty disclaimer) changes.
000228-684.20091106.PBNChange.Notification	RV110609N	Submission of a change to the primary brand name to replace "Pro" with "SPC"
000228-00684.20100716.FirstAid&S&D.Amendment	RV071610N	Amendment to add the statement "Do not give any liquid to the person" to the first aid section. Added all PRN 2007-4 Storage and Disposal language options, CHEMTREC & Medical number, and Spanish/English understanding label statement. Reformatted label to NuFarm template