



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Mr. George Meindl
Nufarm America, Inc.
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527

AUG - 6 2010

Subject: Label Amendment- Removal of all non-residential uses and associated language
Product Name-NUP-09064 Herbicide
EPA Reg. No. 228-704

Dear Mr. Meindl:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records.

Submit one (1) copy of final printed labeling before you release the product for shipment.

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague".

Kathryn V. Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

NUP-09064 HERBICIDE

PREMIUM LAWN WEED KILLER Concentrate

FOR SELECTIVE BROADLEAF AND GRASSY WEED CONTROL IN RESIDENTIAL AND ORNAMENTAL TURFGRASSES. TO CONTROL CRABGRASS, DANDELION, CLOVER, HENBIT, PLANTAINS, WILD ONION, AND MANY OTHER LISTED LAWN WEEDS.

CONTAINS 2,4-D, QUINCLORAC AND DICAMBA
1 QUART TREATS (COVERS) UP TO 5,000 SQ. FT.

ACTIVE INGREDIENTS:

2,4-D DMA*	6.557%
Quinclorac**	2.724%
Dicamba Acid***	0.681%

OTHER INGREDIENTS:	90.038%
TOTAL:	100.000%

Isomer Specific Method, Equivalent to:

* 2,4-D	5.446%, 0.473 lb/gal.
** Quinclorac	2.724%, 0.236 lb/gal.
*** Dicamba Acid	0.681%, 0.059 lb/gal.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with COMMENTS
In EPA Letter Dated:
AUG - 6 2010
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

EPA REG. NO. 228-704
EPA EST. NO.
NUP-09064

Manufactured For
NUFARM AMERICAS INC.
150 Harvester Drive
Burr Ridge, IL 60527

228-704

___ PT.

NET CONTENTS: ___ GALS. ___ QT.

000228.00707.201005131.EPA

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- Protective eyewear (goggles, face shield or safety glasses)
- Long-sleeved shirt and long pants
- Shoes plus socks, and
- Chemical-resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other spraying. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these herbicides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Only protected handlers may be in the area during application. Do not allow people (or pets) to enter the treated area until sprays have dried. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE RESTRICTIONS

Do not allow drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.

Do not pour spray solutions near desirable plants.

Do not apply sprays which are too fine because mists can drift and damage other plants.

Do not allow contact with exposed feeder roots of ornamentals and trees.

Do not apply broadcast applications when air temperature exceeds 90°F.

Do not use on lawns with desirable clovers and legumes or on ornamentals.

Do not mow for two days before or two days after application.

Do not use clippings as mulch or compost around flowers, ornamentals, trees or in vegetable gardens.

Do not apply this product by air or through any type of irrigation equipment.

USE PRECAUTIONS

Shake well before using. Apply this product to young, actively growing weeds as a postemergence entire lawn or spot spray. Follow-up applications may be required for dense infestations of broadleaf and grassy weeds. Applications of this product made to annual grasses at 2 to 4 tiller may not provide complete control. Use lawn type sprayer with coarse spray as wind drift is less likely... Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. If certain tough weeds are not dead within 3 weeks, repeat with spot treatment application. Do not apply broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperature over 90°F, turf injury may occur.

Do not use on lawns with desirable clovers and legumes or on ornamentals. In newly seeded areas, application to grass seedlings is not recommended until after the third mowing. In newly sodded, sprigged or plugged areas, application should be delayed until 3 to 4 weeks after sodding, sprigging or plugging operations. Delay applications for 4 weeks after seeding and emergence of all turf species listed in this label. Do not mow for two days before or two days after application. Clippings from the first 3 mowings should be left in the application area. Do not use clippings as mulch or compost around flowers, ornamentals, trees or in vegetable gardens.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre (or rate/1,000 sq. ft.) which does not exceed the maximum labeled rate for the turf type where the product is to be applied. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

Do not apply this product by air or through any type of irrigation equipment. If soil moisture is not sufficient prior to application, irrigation may improve weed control. Do not water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least on-half inch is required.

Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

Make no more than 2 applications per year. For ornamental lawns and turf, apply no more than 1.5 lbs. ae 2,4-D per acre (0.5 oz. ae 2,4-D per 1,000 sq. ft.) per application and no more than 3.0 lbs. ae 2,4-D per acre (1.0 oz. ae 2,4-D per 1,000 sq. ft.) per year, excluding spot treatments.

This product combines three proven active ingredients [Quinclorac, 2,4-D, and Dicamba] to provide broad spectrum control of grassy and broadleaf weeds. This herbicide is absorbed by foliage and roots of weeds and translocated throughout the plant. USE AS DIRECTED TO CONTROL OR SUPPRESS THE FOLLOWING WEEDS:

WEEDS:

GRASSY WEEDS

- Crabgrass, Large
- Crabgrass, Smooth
- Barnyardgrass
- Foxtail, Giant
- Foxtail, Green
- Foxtail, Yellow
- Signalgrass,
- Broadleaf

BROADLEAF WEEDS

- Alder
- Annual yellow sweet
Clover
- Artichoke
- Aster
- Austrian fieldcress
- Bedstraw
- Beggartick
- Biden
- Bindweed
- Bird vetch
- Bitterweed
- Bitter wintercress
- Black-eyed Susan
- Black medic
- Black mustard
- Blackseed plantain
- Blessed thistle
- Blue lettuce
- Blue vervain
- Box elder
- Bracted plantain
- Brassbuttons
- Bristly oxtongue
- Broadleaf dock
- Broadleaf plantain
- Broomweed
- Buckhorn
- Buckhorn plantain
- Bulbous buttercup
- Bull nettle
- Bull thistle
- Burdock
- Burning nettle
- Bur ragweed
- Burweed
- Buttercup
- Canada thistle

- Carolina geranium
- Carpetweed
- Catchweed bedstraw
- Catsear
- Catnip
- Chickweed
- Chicory
- Cinquefoil
- Common
chickweed
- Common mullein
- Common sowthistle
- Com chamomile
- Creeping Jenny
- Crimson clover
- Croton
- Cudweed
- Curly dock
- Curly indigo
- Dandelion
- Dead Nettle
- Dock
- Dollarweed
- Dogbane
- Dogfennel
- Elderberry
- English daisy
- Fall dandelion
- False dandelion
- False flax
- False sunflower
- Fiddleneck
- Field bindweed
- Field pansy
- Flea bane (Daisy)
- Flixweed
- Florida betony
- Florida pusley
- Frenchweed
- Galinsoga
- Garlic mustard
- Goathead
- Goatsbeard
- Goldenrod
- Ground ivy
- Gumweed
- Hairy bittercress
- Hairy fleabane
- Hawkweed
- Healall
- Heartleaf drymary
- Hedge bindweed
- Hedge mustard
- Hemp
- Henbit
- Hoary cress
- Horsetail
- Indiana mallow
- Ironweed
- Jewelweed
- Jimsonweed
- Kochia
- Knawel
- Knotweed
- Lambsquarter
- Lespedeza
- Locoweed
- Lupine
- Mallow
- Marshelder
- Matchweed
- Mexicanweed
- Milk vetch
- Milkweed
bloodf lower
- Morningglory
- Mouseear
chickweed
- Musk thistle
- Mustard
- Narrowleaf plantain
- Narrowleaf vetch
- Nettle
- Orange hawkweed
- Oxalis
- Oxeye daisy
- Parsley-piert
- Parsnip
- Pearwort
- Pennycress
- Pennywort
- Peppergrass
- Pepperweed
- Pigweed
- Pineywoods
bedstraw
- Plains coeopsis
- Plantain
- Poison hemlock
- Poison ivy
- Poison oak

- Pokeweed
- Poorjoe
- Povertyweed
- Prostrate spurge
- Prostrate vervain
- Puncture vine
- Purslane
- Ragweed
- Red clover
- Redroot pigweed
- Red sorrel
- Redstem filaree
- Rough cinquefoil
- Rough fleabane
- Roundleafed
marigold
- Rush
- Russian pigweed
- Russian thistle
- St. Johnswort
- Scarlet pimpernel
- Scotch thistle
- Sheep sorrel
- Shepherdspurge
- Slender plantain
- Smallflower
galinsoga
- Smartweed
- Smooth dock
- Smooth pigweed
- Sneezeweed
- Southern wild rose
- Sowthistle
- Spanishneedle
- Spatterdock
- Speedwell
- Spiny amaranth
- Spiny cocklebur
- Spotted catsear
- Spotted knapweed
- Spotted spurge
- Spurge
- Spurweed
- Stinging nettle
- Stinkweed
- Stitchwort
- Strawberry clover
- Sumac
- Sunflower
- Sweet clover
- Tall nettle

- Tarweed
- Thistle
- Tick trefoil
- Toadflax
- Trailing crown vetch
- Tumble mustard
- Tumble pigweed
- Tumbleweed
- Velvet leaf
- Venice mallow
- Veronica
- Vervain
- Vetch
- Violet, wild
- Virginia
buttonweed
- Virginia creeper
- Virginia pepperweed
- Wavyleaf bullthistle
- Western clematis
- Western salsify
- White clover
- White mustard
- Wild mustard
- Wild aster
- Wild buckwheat
- Wild carrot
- Wild four-o'clock,
- Wild garlic
- Wild geranium
- Wild lettuce
- Wild marigold
- Wild onion
- Wild parsnip
- Wild radish
- Wild rape
- Wild strawberry
- Wild sweet potato
- Wild vetch
- Willow
- Witchweed
- Woodsorrel
- Woolly croton
- Woolly plantain
- Wormseed
- Yarrow
- Yellow Rocket
- Yellow flower
pepperweed
and other broadleaf
weeds.

Some of these species may require repeat spot applications and/or use of the maximum specified rate on this product label even under ideal conditions for application.

USE SITES

This product can be applied postemergence to the following turfgrasses:

HIGHLY TOLERANT TURFGRASSES	MODERATELY TOLERANT TURFGRASSES	DO NOT APPLY TO THE FOLLOWING TURFGRASSES
COOL SEASON TURFGRASS Bluegrass, Kentucky Fescue, Tall Ryegrass, Annual Ryegrass, Perennial	COOL SEASON TURFGRASS Bentgrass, Creeping (1,3) Bluegrass, Annual (Poa Annua) (1) Bluegrass, Rough (Poa trivialis) Fescue, Chewing's Fescue, Fine (2) Fescue, Hard Fescue, Red WARM SEASON TURFGRASS Bermudagrass, Common (1) Bermudagrass, (Hybrids) (1) Zoysiagrass	Bahiagrass Bentgrass, Colonial Bentgrass, Seaside Buffalograss Carpetgrass Centipedegrass Dichondra Paspalum, Seashore St. Augustinegrass
1. Temporary yellowing and discoloration on these species may occur after application. 2. Only apply to fine fescue when it is part of a blend. 3. Application to established creeping Bentgrass must be applied in 2 split applications at 2.4 to 4.0 fl. oz. per 1,000 sq. ft.		

Warm season turfgrasses such as Common Bermuda and Zoysia may see temporary discoloration. Do not apply during growth stages from dormancy to green-up and from green-up to dormancy.

HOW MUCH TO USE

SPOT TREATMENT

Pressure Sprayer [Pump-up Sprayer]: Adjust spray nozzle to give a coarse spray. Aim at center of weed and spray to wet. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. Do not use a hose-end sprayer for spot treatment.

AMOUNT OF CONCENTRATE (FLUID OUNCES)	AMOUNT OF WATER (GALLONS)	AREA TREATED (SQ. FEET)
6.4	4	1,000
3.2	2	500
1.6	1	250

ENTIRE LAWN TREATMENT

Dial Type Hose-end Sprayers: Spray lawn using a coarse spray. Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

- 1) Measure the total square foot area to be sprayed. To determine the total square foot area, multiply the length by the width of the lawn area to be treated. Subtract square footage of non-treatment areas including flower beds, shrub beds, driveways, and sidewalks.
- 2) The application rate of this product is 6.4 fluid ounces per 1,000 sq. ft of lawn. Add 6.4 fl. oz. of this product to the spray bottle [jar], [reservoir] for every 1,000 sq. ft. of lawn area to be treated.
- 3) Set the dial to the 2.5 fl. oz. per gallon mix rate.
- 4) Connect to hose, turn on water and spray evenly over the lawn treatment area.

AREA TO BE TREATED (SQ. FEET)	AMOUNT OF PRODUCT NEEDED (FLUID OUNCES)
1,000	6.4
2,000	12.8
3,000	19.2
4,000	25.6
5,000	32.0

Broadcast Application: Spray lawn using a coarse spray. Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

AREA TO BE TREATED (SQ. FEET)	AMOUNT OF PRODUCT NEEDED (FLUID OUNCES)
1,000	6.4
2,000	12.8
3,000	19.2
4,000	25.6
5,000	32.0

Ready-to-spray Hose-end Sprayers: *[This product is packaged in a ready-to-spray packaging containing a plastic container fitted with a spray head that sprays the solution at a fixed concentration.]* Apply evenly over area to be treated. One application should be sufficient. Effects begin to show after a few days and weeds gradually die.

This sprayer bottle will treat 5,000 sq. ft. when applied at 6.4 fluid ounces of this product per 1,000 sq. ft.

Instructions

1. Attach garden hose to spray head.
2. Turn on water supply.
3. Bend safety tab down and turn control valve to "water".
4. Point nozzle toward lawn and turn control valve to "on". Product will automatically mix with water.
5. Walk back and forth at a steady pace and spray evenly over the lawn treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter the product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

Or

CONTAINER HANDLING:

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV050610)

QUINCEPT is a registered trademark of Nufarm Americas Inc.

[OPTIONAL BOX]

PRODUCT FACTS

What it Does:

Kills Broadleaf Weeds including Chickweed, Clover, Dandelion and Grassy Weeds including Crabgrass (See inside for complete list).

Where to Use: (Icon weeds)

Lawns: Kills the Weeds not the Lawn.

When to Use:

Do not apply broadcast applications when air temperature exceeds 90°F.

Questions and Comments: (Icon phone) (icon computer)

Call xxx-xxx-xxxx or visit our website at www.xxxxxxxx.xxx

MEASUREMENT CALCULATIONS

3 teaspoons = 1 Tablespoon	1 fluid ounce = 6 teaspoons
1/2 fluid ounce = 3 teaspoons	1 fluid ounce = 2 Tablespoons
1/2 fluid ounce = 1 Tablespoon	4 fluid ounces = 1/2 cup

OPTIONAL MARKETING CLAIMS - FRONT PANEL

Kill Crabgrass

[Also] Controls Crabgrass

Kills crabgrass, dandelion, clover and 200+ weeds

Controls Sensitive Weeds such as...

Controls both grassy and broadleaf weeds in one easy step

Controls Hard-to-Kill weeds such as...

Kills Broadleaf and grassy weeds

Kills over 200 weeds

Kills Crabgrass and All Major Broadleaf Weeds

Kills Dandelion, Clover and Other Broadleaf Weeds

Kills Dandelion, Clover and Other Listed Broadleaf Weeds

Use When Weeds Are Actively Growing

Kills the Weeds, Not the Lawn

Rainproof in Hours - Won't Wash Away

Won't Harm the Home Lawn when Used as Directed

Illustration(s) of various weeds listed on label

Easy to Use

For Thick, Weed-Free Turf

Three Way Control

Apply to a Moist Lawn

Contains Three Way Power

One Quart Covers Up To 5,000 Square Feet, (One XXX covers up to XX,XXX square feet)

Easy to Mix

Just Add Water

New Formulation

Simple to use

Proven Performance

New!