



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

228-728

Date of Issuance:

10/30/2014

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Azure Xtra Fungicide

Name and Address of Registrant (include ZIP Code):

Nufarm Americas Inc.
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Date:

10/30/2014

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 228-728.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSF:

- Basic CSF dated 10/19/2014

If you have any questions, please contact Tamue L. Gibson at (703) 305-9096 or gibson.tamue@epa.gov.

ACCEPTED

10/30/2014

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 228-728

GROUP 3 11 FUNGICIDES

AZURE™ Xtra Fungicide

Broad-spectrum fungicide for control of plant diseases

[Alternate Brand Name: NUP-14035 Fungicide]

ACTIVE INGREDIENTS

Azoxystrobin (CAS No. 131860-33-8)18.2%
Cyproconazole (CAS No. 94361-06-5).....7.3%

OTHER INGREDIENTS74.5%

TOTAL100.0%

Azure Xtra Fungicide is a 280SC formulation that contains 0.67 lb a.i. cyproconazole and 1.67 lb a.i. azoxystrobin per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Label Booklet for First Aid and Precautionary Statements

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840**

EPA Reg. No. 228-XXX
EPA Est. No. _____

**MANUFACTURED FOR
NUFARM AMERICAS INC.
11901 S. AUSTIN AVENUE
ALSIP, IL 60803**



NET CONTENTS: _____ Gallons (_____ Liters)

[Designation as “NONREFILLABLE” or “REFILLABLE” for containers > 5 GAL]

**[Grow a Better Tomorrow]
[Nufarm Grow a Better Tomorrow]**

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. • Call a poison control center or doctor for future treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (877) 325-1840 for emergency medical treatment information.</p>	

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants and shoes plus socks..

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Azoxystrobin is toxic to freshwater, estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. For terrestrial uses do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash-water or rinsate.

Ground Water Advisory

Azoxystrobin, a degradate of azoxystrobin, and cyproconazole are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Nufarm immediately if you observe any adverse environmental effects due to use of this product.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

PRODUCT USE INFORMATION

This product is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. Applications may be made as foliar sprays in alternating spray programs or in tank mixes with other crop protection products. All applications must be made according to the use directions that follow.

USE PRECAUTIONS

Do not use in nurseries, greenhouses or landscape plantings.

Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if the maximum amount of product has been applied.

The active ingredients in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your County has a Bulletin consult http://128.104.224.199/espp_front/view.jsp. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

ROTATIONAL CROPS

Soybeans, corn, or wheat may be replanted immediately, if crop is lost. Cereal grains other than wheat may be planted 180 days after the last application. Cotton may be planted 180 days after the last application. All other crops (including leafy vegetables) may be planted 270 days after the last application.

PHYTOTOXICITY

This product demonstrates some phytotoxic effects when mixed with products that are formulated as ECs. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone can contribute to phytotoxicity. Under certain environmental conditions, tank mixes of this product plus herbicides and/or fertilizers may cause crop injury.

SPRAY DRIFT MANAGEMENT

ATTENTION

This product is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray this product where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.

INTEGRATED PEST (DISEASE) MANAGEMENT

Use this product as part of an integrated disease and pest management strategy whenever the use of a fungicide is needed. Follow cultural practices known to help reduce disease development. The **CROP SPECIFIC USE DIRECTIONS** section in this label identifies specific IPM recommendations for each crop. Consult your local agricultural authorities for additional IPM strategies established for your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

GROUP	3	11	FUNGICIDES
-------	---	----	------------

This product is a mixture of Group 3 (cyproconazole) and Group 11 (azoxystrobin) fungicides. Therefore this product has two modes of action: (1) DMI (demethylation inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (QoI) [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Nufarm encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Follow crop specific recommendations that limit the total number of sprays on a crop and the required alternations with fungicides from other resistance management groups. In situations requiring multiple sprays (more than two), develop season long spray programs for Group 11 (QoI) fungicides. The program should meet the goal of no more than 1/3 of the total sprays per season, when a Group 11 fungicide is used as a solo product, or 1/2 the total sprays when a Group 11 fungicide is used in a mixture. Programs that include both solo Group 11 products and/or mixes containing Group 11 products should be no more than 1/2 the total sprays.

Avoid alternating or tank mixing this product with any fungicide to which resistance has already developed.

SPRAY EQUIPMENT

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

Use a pump with capacity to: (1) maintain at least 35-40 psi at nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute. Use a jet agitator or liquid sparge tube for agitation. Do not air sparge.

This product is a suspoemulsion concentrate. It is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state and local agricultural recommendations.

SPRAYING/MIXING

This product may be applied with all types of spray equipment commonly used for ground and aerial applications. Do not apply this product through any type of ultra low volume (ULV) spray system. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control. Higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist.

For ground applications, apply this product in a minimum of 10 gallons of water volume for adequate coverage and canopy penetration. For aerial applications, follow local recommendations.

Prepare no more spray mixture than is required for immediate use. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Azure Xtra Alone: Add 1/2 - 2/3 of the required amount of water to a spray or mixing tank. With the agitator running, add this product to the tank and continue agitation while adding the remainder of the water. Begin application of the spray solution after complete dispersion of the product into the mix water and maintain agitation until all of the mixture has been sprayed.

Azure Xtra + Tank Mixtures: Do not combine this product in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious to the crop under your conditions of use. To determine physical compatibility of this product and other products, use a

jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables (suspoemulsions), and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Add 1/2 - 2/3 of the required amount of water to a spray or mixing tank. With the agitator running, add the tank mix partner into the tank. Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and this product to the spray tank. Allow the product to completely disperse. Spray the mixture with the agitator running.

Observe the most restrictive of the labeling limitations and precautions of all products used in tank mixtures.

APPLICATION INSTRUCTIONS

Apply this product at rates and timings as described in this label. Apply by ground, by air, or via chemigation. Refer to crop section for specific directions.

Apply this product in a minimum of 10 GPA via ground application equipment and 2 GPA via aerial application equipment.

Directions for Use Through Sprinkler Chemigation Systems (apply this product only to crops for which chemigation is specified on this label):

Spray Preparation: Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.

Use Precautions for Sprinkler Irrigation Applications

Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, drip (trickle), or hand move irrigation systems. Do not apply this product through any other type of irrigation system except as specified on this label.

Apply with center pivot or continuous-move equipment distributing 1/2 acre-inch or less during treatment. In general, use the least amount of water required for proper distribution and coverage. If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, this product should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips. Do not apply when wind speed favors drift beyond the area intended for treatment. Plant injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

If you have questions about calibration you should contact State Extension Service specialist, equipment manufacturers or other experts.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system

interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

CROP SPECIFIC USE DIRECTIONS

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Cereals Wheat Triticale	Early season suppression of: Glume blotch (<i>Stagonospora nordorum</i>) Leaf blight (<i>Septoria tritici</i>) Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Tan spot (<i>Pyrenophora tritici-repentis</i>)	3.5	Apply this product in the spring at approximately Feekes Stage 5. The minimum retreatment interval is 14 days. Apply by ground, air, or chemigation. If applied by chemigation, the maximum amount of water should be 0.25 inches.
	Control of Leaf Diseases: Glume blotch (<i>Stagonospora nordorum</i>) Helminthosporium leaf blight (<i>Drechslera tritici-repentis</i>) Leaf blight (<i>Septoria tritici</i>) Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Rust (stripe, leaf, and stem) (<i>Puccinia</i> spp.) Spot blotch (<i>Bipolaris sorokiniana</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>)	5–6.8	Apply this product between Feekes Stage 8 and 10.5.1. Protecting the flag leaf is important for maximizing the potential yield. The minimum retreatment interval is 14 days. Treat up to 30 days prior to harvest. Apply by ground, air, or chemigation. Resistance Management: No more than two foliar applications of this product or other strobilurin fungicides should be made per growing season. Do not alternate or tank mix with fungicides to which resistance has developed in the pathogen population.
Specific Use Restrictions: (1) Do not apply more than 6.8 fl oz/A of this product per season. (2) Do not harvest wheat within 30 days of application. (3) Do not harvest forage or hay within 21 days of application. (4) Do not apply more than 0.036 lb a.i./A/season cyproconazole containing products. (5) Do not apply more than 0.4 lb a.i./A/year azoxystrobin containing products.			

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Corn Field Seed	Eye spot <i>(Aureobasidium zeae)</i> Gray leafspot <i>(Cercospora zeae-maydis)</i> Northern corn leaf blight <i>(Setosphaeria turcica)</i> Northern corn leaf spot <i>(Cochliobolus carbonum)</i> Rusts (common and Southern) <i>(Puccinia spp.)</i> Southern corn leaf blight <i>(Cochliobolus heterostrophus)</i> [also known as Helminthosporium leaf blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>)]	3.5–6.8	<p>Apply this product when disease first appears. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease development, apply the high rate. A second application may be made 7–14 days later.</p> <p>Application Directions: Make no more than 2 applications of this product or any other Group 11 fungicide per year. Apply by ground, air, or chemigation.</p> <p>Resistance Management: No more than two foliar applications of this product or other strobilurin fungicides should be made per growing season. Do not alternate or tank mix with fungicides to which resistance has developed in the pathogen population.</p>
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> (1) Do not apply within 30 days of harvest. (2) Do not harvest silage within 21 days of an application. (3) Do not apply more than 6.8 fl oz/A/season of this product. (4) Do not apply more than 0.036 lb a.i. cyproconazole containing products/A/year. (5) Do not apply more than 2.0 lb a.i. azoxystrobin containing products/A/year. 			

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Soybean			<p>Application Directions: Coverage and penetration are important for best results. Use sufficient water volume to provide thorough and uniform plant coverage. Apply by ground, air or chemigation. Addition of an additive with spreading and penetrating qualities will enhance coverage and efficacy.</p> <p>Resistance Management: No more than two foliar applications of this product or other strobilurin fungicides should be made per growing season. Do not alternate or tank mix with fungicides to which resistance has developed in the pathogen population.</p>
	Rust (<i>Phakopsora</i> spp.)	4.0–6.8	<p>For soybean rust: Repeat applications at 14- to 28-day intervals if conditions favorable for rust development persist. Lower use rates may require a shorter spray interval. Depending on the conditions, application timing should be at R1 (beginning flowering, approximately 50 days after planting) up to the R6 stage (seed development), but could be earlier.</p>
	Aerial blight (<i>Rhizoctonia solani</i>) Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum truncatum</i>) Brown spot (<i>Septoria glycines</i>) Cercospora blight and leaf spot (<i>Cercospora kikuchii</i>) Frogeye leafspot (<i>Cercospora sojina</i>) Pod and Stem blight (<i>Diaporthe phaseolorum</i>)	5.0–6.8	<p>For diseases other than soybean rust: Begin applications prior to disease development. Use the higher rates under conditions favorable for severe disease pressure, dense plant canopies, or when disease is present. Contact extension personnel for local economic thresholds and timings for specific diseases in your area. An adjuvant may be added at recommended rates to improve coverage.</p>
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> (1) Do not apply more than 13.6 fl oz of this product/A/season. (2) Do not apply within 30 days of harvest of soybeans (bean). (3) Do not graze forage within 14 days of an application. (4) Do not apply more than 0.072 lb a.i./A/year of cyproconazole containing products. (5) Do not apply more than 1.5 lb a.i./A/year of azoxystrobin containing products. (6) Do not use soybean forage or hay as livestock feed if making more than one application at 6.8 fl oz product/A. 			

RATE CONVERSION TABLE

Product (fl oz/A)	Cyproconazole (lb a.i./A)	Azoxystrobin (lb a.i./A)	Treated acres per gallon product
3.5	0.018	0.046	36.6
4.0	0.021	0.052	32.0
5.0	0.026	0.066	25.6
6.5	0.034	0.085	19.7
6.8	0.036	0.089	18.8
13.7	0.072	0.178	N/A

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[Note to Reviewer: The following statement will be included on all Final Printed Labels bearing multiple Container Handling statements] **“NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this product's labeling for the applicable “Nonrefillable” or “Refillable” designation. Follow the container handling instructions below that apply to your container type/size.”

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure-rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[Refillable containers larger than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product and return it unopened to the Seller, and the purchase price will be refunded.

(RV103014)

Azure is a trademark of Nufarm Americas Inc.

All other trademarks that appear on this label which are not owned by Nufarm Americas Inc. or its subsidiaries are the property of their respective owners.