



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

228-746

Date of Issuance:

1/7/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

MEPEX® 6X Plant Growth
 Regulator

Name and Address of Registrant (include ZIP Code):

Danielle Larochelle, Senior Regulatory Manager
 Nufarm Americas Inc.
 4020 Aerial Center Parkway, Suite 101
 Morrisville, NC 27560

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Signature of Approving Official:

Carmen J. Rodia Jr., Acting Product Manager 22
 Fungicide Branch, Registration Division (7505P)

Date:

1/7/22

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 228-746.”
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 7/27/20
- Alternate CSF #1 dated 7/27/20

If you have any questions, please contact Carmen J. Rodia, Jr. by phone at (703) 306-0327, or via email at Rodia.Carmen@epa.gov; or Craig Reeves by phone at (202) 566-2869, or via email at Reeves.Craig@epa.gov.

Enclosure

MEPEX® 6X

Plant Growth Regulator

ACCEPTED

01/07/2022

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 228-746

ACTIVE INGREDIENT:

Mepiquat chloride: N,N-dimethylpiperidinium chloride.....24.0%

OTHER INGREDIENTS:76.0%

TOTAL:100.0%

Contains two (2) pounds active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE INSIDE BOOKLET FOR [FIRST AID AND] [ADDITIONAL] PRECAUTIONARY STATEMENTS
[AND COMPLETE DIRECTIONS FOR USE]

For Medical Emergencies, Call (877) 325-1840
For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

FIRST AID

| | |
|-------------------------------|---|
| IF SWALLOWED | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person. |
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF IN EYES | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. |

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (877) 325-1840 for emergency medical treatment information.

EPA Reg. No. 228-TUA

EPA Est. No. _____

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Ave
Alsip, IL 60803



NET CONTENTS: ____ Gallons (____ Liters)

[Designation as “NONREFILLABLE” or “REFILLABLE” for containers > 5 GAL]

[Optional Marketing Tagline: **Grow a better tomorrow.**]

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or if absorbed through the skin. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. In case of contact immediately flush eyes or skin with plenty of water. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some of the materials that are chemical-resistant to this product are listed below. Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Non-target Organism Advisory: This product may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

Runoff Prevention Advisory Statement: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks.

SPRAY DRIFT REDUCTION

Applicators are required to use a medium to coarser droplet size, as defined by ASABE Standard S572.1.

Do not apply when wind speeds exceed 10 miles per hour at the application site.

Do not apply during temperature inversions

When using ground application equipment:

- Apply with nozzle height no more than 3 feet above the ground or crop canopy.
- Do not apply with airblast equipment.

When applying aerially:

- Do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- The spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- When applying to crops via aerial application equipment, use ½ swath displacement upwind at the edge of the field.
- Orient nozzles so the spray is directed toward the back of the aircraft.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

PRODUCT INFORMATION

Chemigation Prohibition: Do not apply this product through any type of irrigation system.

MEPEX® 6X Plant Growth Regulator is a foliar applied plant regulator. It allows the grower to manage the cotton plant for short-season production leading to reduced risk of yield and quality loss due to delayed and prolonged harvest. Benefits derived from the use of this product include:

- Increased early boll retention and/or larger bolls
- Reduced plant height which provides a more open canopy / less boll rot
- Improved defoliation
- Less trash and lower ginning costs
- Better harvest efficiency, and
- Darker leaf color

These benefits can provide for earlier maturity and often result in improved yields.

APPLICATION INSTRUCTIONS

EARLY APPLICATION

Growers may use low-rate multiple applications, or higher, less frequent dosages which provide maximum flexibility under a wide range of growing conditions. Do not apply this product to plants under stress. If stress is alleviated, evaluate plants for vegetative growth before additional applications are made. This product may be tank mixed with insecticides/miticides when application timings coincide. (See Restrictions and Limitations).

Fields must be carefully scouted. Do not apply this product if plants are under any form of stress. In the absence of stress, a maximum of five low rate applications can be made each season. The first application can be applied at the matchhead square stage. The rate and timing of subsequent applications depend on growing conditions and desired benefits. Under good growing conditions additional treatments of 1/3 to 2/3 fl oz/acre can be made at 7-14 day intervals. Apply higher rates of this product (2/3 to 2 fl oz/acre) if vegetative growth becomes excessive or a greater degree of height control is desired. Do not use more than a total of 8.4 fl oz/acre of this product in a growing season.

If significant loss of squares and/or young bolls has occurred earlier due to insect pressure or other stresses, but now these stresses have been alleviated, the need for this product is increased - excess vegetative growth is likely because of poor fruit load.

LATE SEASON APPLICATION

Late application of this product (approximately during the fourth to sixth week of blooming) can provide certain benefits to cotton. However, it will not and does not substitute for early season use, the time of the greatest benefit from the use of this product. Late season application can lead to one or more of the following:

- Reduction in late season vegetative growth or regrowth after cutout or defoliation
- More complete and manageable cutout
- Better defoliation
- Earlier maturity and reduction in trash
- Lower ginning costs

Some of these effects may favorably influence the yield potential and fiber quality. Apply a late season application of this product only if fields are not drought or nutrient stressed; that is, those fields likely to experience additional vegetative growth or regrowth. However, fields that are very rank and extremely vigorous due to a combination of poor boll load and excellent growing conditions may not respond as much as desired to late season applications at the specified rates.

Timing for Late Season Applications

- I. **On fields where cotton cuts out and then starts regrowth:** Apply when regrowth begins, as evidenced by new leaves in the terminal and stem elongation. This would often, but not always, be in the period of 5-6 weeks after the first bloom.
- II. **On fields where cotton never completely cuts out:** Apply this product when there are 4-6 nodes above the white flower (NAWF). Measure NAWF by counting the number of mainstem nodes from the first position white bloom (the one closest to the mainstem) to the terminal. Count the node with the first position white bloom as zero and the last node in the terminal, which is counted, should have a leaf at least the size of a quarter. Generally, the NAWF first reaches 4-6 during the fourth to sixth week of bloom. During this time period, the NAWF will be decreasing about one node every 5-6 days – if its rate of decrease is less, this means that the plant is not cutting out soon enough (the crop is too vigorous). If the fifth week of bloom arrives and NAWF is still above 5-6, apply this product.

Use Rate for Late Season Application

Apply this product at a rate between 1.33 to 4.2 fl oz per acre. Use the lower rates on cotton with only moderate additional growth potential, and the higher rates on fields likely to continue vigorous growth. Total seasonal use per season (early plus late applications) must not exceed 8.4 fl oz per acre.

SPRAY VOLUMES: Thorough coverage is required.

In Water:

Areas other than California

Ground application – Use a minimum of 2 gallons of water per acre.

Aerial application - Use a minimum of 2 gallons of water per acre.

California only

Ground application – Use a minimum of 5 gallons of water per acre.

Aerial application - Use a minimum of 5 gallons of water per acre.

In Oil:

Use a minimum total oil volume of 2 pints/acre for ultra low volume (ULV) aerial application. Application in oil is permitted only in AL, AR, FL, GA, KS, LA, MO, MS, NC, OK, SC, TN and TX. Use a nonphytotoxic oil concentrate which contains either a petroleum or vegetable oil base, contains only EPA-exempt ingredients, provides good mixing quality in the jar test (see "Compatibility" section), and has been used successfully in your locality. The oil diluent must contain emulsifiers which provide good mixing quality. If the oil does not contain an emulsifier, one must be added during mixing at a volume equal to 3% of the final volume of the mixing tank. Do not apply this product as ULV without using emulsifiers. If using a vegetable oil based product, use only highly refined concentrates.

Mix under constant agitation. Pour one-half of the required volume of oil into the spray tank, followed by the emulsifier (if the oil does not already contain one) at approximately 3% of the final spray tank volume, and then pour in this product while the remainder of the oil is added. Constant agitation is required during and after mixing and during application.

RAIN WASH-OFF PRECAUTION

The use of a high quality, EPA exempt surfactant will enhance the uptake of this product into the plant. Therefore, the use of a surfactant allows applications made as little as four hours prior to rainfall to be effective. Without a surfactant, the product should be used at least 8 hours prior to expected rainfall.

COMPATIBILITY

This product is water based, and is compatible with most insecticides and miticides. If compatibility is in doubt, perform a jar test to check for compatibility. This product can be used with foliar fertilizers if your experience shows the combination is compatible and will not injure cotton under your conditions. Caution must be used when applying with foliar fertilizers under conditions of extreme heat.

RESTRICTIONS AND LIMITATIONS

- Insect or mite damage to treated crops can lead to yield decreases or other undesirable effects.
- Do not apply this product to cotton that is under stress. If using the low rate multiple applications, discontinue use until your crop has overcome any stress.
- Do not apply more than 8.4 fl oz of this product per acre per season. The sum of all products and formulations containing mepiquat chloride must not exceed 0.132 pound (60 grams) of mepiquat chloride per acre per season.
- Do not apply this product within 30 days of harvest.
- Do not graze or feed cotton forage to livestock.
- This product contains a dye and effectiveness is not related to the color of the spray solution.
- Do not plant another crop within 75 days after last treatment.

TIME AND RATE OF APPLICATION

SHORT-STAPLE AND LONG STAPLE (PIMA) COTTON

Directions for use must be observed as specified below:

I. Low Rate Multiple Applications

Use these instructions when you want to maintain maximum flexibility in plant regulation treatments.

| Area | Time of Application | Rate Per Acre (fl oz) | |
|--|--|--|---|
| | | Excessive Growth NOT Expected or Lower Rates Have Worked in the Past | Excessive Growth Expected or Higher Rates Have Been Necessary in the Past |
| AL, AR, AZ, CA, FL, GA, KS, LA, MO, MS, NC, NM, OK, SC, TN, TX, VA | First Application: Apply at the match-head square* stage of growth. | 0.33 | 0.66 |
| | Second Application: 7-14 days later, or when regrowth occurs. | 0.33 | 0.66 |
| | Third Application: 7-14 days later, or when regrowth occurs. | 0.33 to 0.66** | 0.66 to 1.33** |
| | Fourth Application: 7-14 days later, or when regrowth occurs. | 0.33 to 1.33** | 0.66 to 2** |
| | Fifth Application (if needed): 7-14 days later, or when regrowth occurs. | 0.66 to 1.33** | 0.66 to 2** |
| | Late Season Application: See section titled "LATE SEASON APPLICATION". | 1.33 to 2.66** | 2 to 4.2** |

*Match-head square is when the first square of a typical cotton plant is about the size of a match head (about 1/8 - 1/4" in diameter). Make the first application when 50% of the plants have one or more match head squares.
**Use the higher rates if previous application was not made or if growing conditions favor excessive growth.

II. High Rate Single or Multiple Applications

Use these instructions when you are not able to start growth regulation treatments early, or when you want to make the fewest number of applications.

| Area | Time of Application | Rate per Acre (fl oz) |
|--|---|-----------------------|
| AL, AR, AZ, CA, FL, GA, LA, MO, MS, NC, NM, SC, TN, VA | First Application: Apply when cotton is actively growing and is between 20" and 30" tall, but not more than 7 days beyond early bloom (5-6 blooms per 25 row feet). Also apply if cotton is 24" tall and has no blooms. Use the 1.33 fl oz rate on cotton where excessive vegetative growth is not expected and use 2.66 fl oz where excessive vegetative growth has historically occurred. See Restrictions and Limitations. | 1.33 to 2.66 |
| | Second Application: Make a subsequent application in 2 to 3 weeks if additional growth is desired. | 1.33 to 2.66 |
| | Third Application for Control of Excessive Vegetative Growth: If the cotton field has a history of vigorous growth or if conditions continue to favor vigorous growth, apply a third application 1 to 2 weeks after the second application. | 1.33 to 2.66 |
| | Late Season Application: See section titled "LATE SEASON APPLICATION". | 1.33 to 4.2 |

| Area | Time of Application | Rate per Acre (fl oz) |
|---|--|-----------------------|
| AREAS WITHOUT A HISTORY OF EXCESSIVE VEGETATIVE GROWTH | | |
| KS, OK, TX (except Rio Grande Valley) | First Application: Apply when cotton is in the early bloom stage (5-6 blooms per 25 row feet) and actively growing. Also apply if no blooms are present and the cotton is 20" tall and actively growing. See Restrictions and Limitations. | 1.33 |
| | Second Application: Make a second application in 2 to 3 weeks if additional growth control is desired. | 1.33 |
| | Third Application: If conditions after the second application of this product continue to favor vigorous growth, apply a third application 1 to 2 weeks after the second application. | 1.33 |
| | Late Season Application: See section titled "LATE SEASON APPLICATION". | 1.33 to 4.2 |
| AREAS WITH A HISTORY OF EXCESSIVE VEGETATIVE GROWTH | | |
| KS, OK, TX (including Rio Grande Valley) | First Application: For best results, apply when plants are in early bloom stage (5-6 blooms per 25 row feet) and an average of 24" tall. Treatments can also be made when cotton height averages a minimum of 20" and a maximum of 30" provided cotton is not more than 7 days beyond early bloom. If cotton is 24" tall and has no blooms, apply this product. See Restrictions and Limitations. | 2.66 |
| | Second Application: For fields with a history of excessive growth, or if conditions after the first application favor excessive growth, make a second application in 2 to 3 weeks. | 1.33 to 2.66 |
| | Third Application: If conditions after the second application of this product continue to favor vigorous growth, apply a third application 1 to 2 weeks after the second application. | 1.33 to 2.66 |
| | Late Season Application: See section titled "LATE SEASON APPLICATION". | 1.33 to 4.2 |

CONVERSION CHART

| RATE OF MEPEX® 6X Plant Growth Regulator | ACRES TREATED PER | | | EQUIVALENT RATE OF MEPEX® Plant Growth Regulator (4.2% ai) |
|---|-------------------|-------|--------|--|
| | PINT | QUART | GALLON | |
| 0.33 fl oz/A | 48 | 96 | 384 | 1/8 pint/A |
| 0.66 fl oz/A | 24 | 48 | 192 | 1/4 pint/A |
| 1.33 fl oz | 12 | 24 | 96 | 1/2 pint/A |
| 2.0 fl oz/A | 8 | 16 | 64 | 3/4 pint/A |
| 2.66 fl oz/A | 6 | 12 | 48 | 1 pint/A |
| 4.2 fl oz/A | 3.8 | 7.6 | 30.4 | 1 1/2 pints/A |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store below 32°F or above 100°F. Store in a dry place away from heat or open flame.

PESTICIDE DISPOSAL: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Note to Reviewer: The following statement will be included on all Final Printed Labels bearing multiple Container Handling statements] **“NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this product’s labeling for the applicable “Nonrefillable” or “Refillable” designation. Follow the container handling instructions below that apply to your container type/size.”

[Nonrefillable Containers 5 Gallons or Less:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable containers larger than 5 gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Refillable containers larger than 5 gallons:] Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED “AS IS” BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT

BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV082621)

[Mepex is a trademark of Nufarm Americas Inc.]

[All other trademarks are the property of their respective owners.]

[All trademarks that appear on this label which are not owned by Nufarm Americas Inc. or its subsidiaries are the property of their respective owners]