

2217-846

11/26/2012

1/28



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

James Kuntsman
PBI/Gordon Corporation
1217 W 12th Street
P O Box 014090
Kansas City, Missouri 64101-0090

11-26-12

Subject EPA Reg 2217-846 / EH-1388 Herbicide Notification

Dear Mr Kuntsman

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 10-29-12 for the product EPA Reg 2217-846 / EH-1388 Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis".

Kable "Bo" Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

October 29, 2012

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U S Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn Mr Kable Bo Davis, (PM-25)

Dear Mr Davis

NOTIFICATION Labeling notification per PRN 98-10 Notifications, Non-notifications and
Minor Formulation Amendments Section II (N) Other revisions
EH-1388 Herbicide (EPA Reg No 2217-846)

I Request for advertising/marketing claims

- 1 We ask to add advertising claims that are consistent with the Directions for Use
previously approved These phrases may be presented on the container labels of EH-
1388 Herbicide (EPA Reg No 2217-846)
 - Covers up to XXX[*]
 - *When mixed XXX of product with YYY of water
 - *When mixed at the rate of XXX per YYY of water
- 2 One (1) copy of the proposed labeling is enclosed for your review Please refer to
page 24 of the draft labeling

II Certification Statement

This notification is consistent with the provisions of PR Notice 98-10 and EPA
regulations at 40 CFR 152.46, and no other changes have been made to the label or
the confidential statement of formula of this product I understand that it is a violation
of 18 U S C Sec 1001 to willfully make any false statement to the EPA. I further
understand that if this notification is not consistent with the terms of PR Notice 98-10
and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to
enforcement action and penalties under Section 12 and 14 of FIFRA

III Enclosures

11/28/18 5/28
Acceptable ER

EH-1388 HERBICIDE

Alternate Brand Names GroundWork Concentrate 25% Weed & Grass Killer

Sublabel 1

ACTIVE INGREDIENT

Isopropylamine salt of glyphosate*

25.0%

OTHER INGREDIENTS

75.0%

TOTAL 100.0%

*N (phosphonomethyl) glycine

KEEP OUT OF REACH OF CHILDREN


CAUTION

See [back/side/inside panel or booklet]
for additional Precautionary Statements and First Aid
or [Peel open for use directions and further information]

**NET CONTENTS (16 fl oz, Quart, 40 fl oz,
½ Gallon, 1 Gallon)**

954/AP102912
EPA Reg No 2217-846
EPA Est No 2217-KS-2

MANUFACTURED BY



Telephone 800 821 7925



**READ THE ENTIRE LABEL FIRST OBSERVE ALL PRECAUTIONS
AND FOLLOW DIRECTIONS CAREFULLY**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION Causes moderate eye irritation Harmful if absorbed through skin Harmful if swallowed
Avoid contact with skin eyes or clothing Wash thoroughly with soap and water

First Aid

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 - 20 minutes • Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth-to mouth if possible • Call a poison control center or doctor for treatment advice
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment	

ENVIRONMENTAL HAZARDS

Do not apply directly to water Do not contaminate water when disposing of equipment washwater and rinsate

PHYSICAL OR CHEMICAL HAZARDS

Combustible Do not use or store near heat or open flame

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Keep children and pets out of treated areas until sprays have dried

STORAGE & DISPOSAL

Do not contaminate water food or feed by storage and disposal

PESTICIDE STORAGE Store in original container in a locked storage area inaccessible to children and domestic animals Protect container from freezing

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons]

PESTICIDE DISPOSAL AND CONTAINER HANDLING Nonrefillable container Do not reuse or refill

this container **If empty** Place in trash or offer for recycling if available **Partly filled** Call your local solid waste agency or www website com for disposal instructions Never place unused product down any indoor or outdoor drain

HOW IT WORKS

Facts and Tips

- EH 1388 Herbicide is a multipurpose herbicide designed to control a broad spectrum of emerged annual and perennial weed species EH-1388 Herbicide will control unwanted species of grass sedge broadleaf weeds and certain woody species around trees shrubs fences
- This product kills plants by inhibiting the production of a substance found only in plants It enters plants through the sprayed foliage and moves to the roots
- This product is nonselective and will kill or injure all vegetation contacted by the spray Do not spray plants or grasses you like—they may die too Not recommended for spot weed control in lawns since this product kills lawn grasses
- No soil residual activity—any product not absorbed by plants breaks down into natural materials without moving in or on the soil to untreated plants
- Plants that have not emerged at the time of application will not be controlled
- Time to symptoms—best results will be obtained when the weed species are actively growing Visible effects are gradual—wilting and yellowing advancing to complete browning and root destruction Weeds usually yellow and wilt within 2 to 4 days and plant death occurs in 1 to 2 weeks

Product Description and Spray Coverage This product is a nonselective herbicide that will kill all vegetation contacted This herbicide kills unwanted plants by inhibiting the production of a substance found only in plants It enters plants through the sprayed foliage and moves to the roots Avoid direct spray or drift onto desirable plants lawn grasses shrubs and trees Protect desirable plants and vegetation with a piece of plastic or cardboard If desirable plants are accidentally sprayed wash plants with water immediately Any glyphosate not absorbed by the plant is broken down into natural materials This product has no soil residual activity and thus will not affect nearby untreated plants

WHERE TO USE

For General Weed Control and Spot Spraying

- Use to prepare sites for large plantings for planting flower beds for planting fruit trees ornamentals vegetables garden sites and to renovate lawns
- Along and on sidewalks gravel/brick walkways paths curbs paved areas (private roads and streets) driveways parking areas recreational areas
- Along ornamental lawns and turfgrass Edging along/around lawns/yards and landscape plantings
- In areas for lawn replacement or lawn renovation/replacement
- Around or near ornamental/shade trees fruit and nut trees grapevines shrubs groundcovers flower beds mulched landscape beds
- For garden plot preparation vegetable plot preparation and small fruits and berries plot preparation
- For flower garden preparation
- Along fences hedgerows under fencelines
- Around rocks signposts light posts or on fringe areas
- Around or along buildings/structures foundations patios houses mobile homes vacant lots urban areas domestic outdoor premises and other areas associated with household or home life
- Urban areas where total vegetation control is desired
- Farmsteads where total vegetation control is desired around farmstead building foundations along fences hedgerows fencerows under fencelines around shelter belts and other areas where nonselective weed control is desired
- Can be used for habitat restoration controlling exotic vegetation and for site preparation of wildlife food plots

TOUGH WEED CONTROL (ANNUALS, PERENNIALS, BRUSH)**CONTROLS WEEDS AND GRASSES INCLUDING LAWN AND TURFGRASS SPECIES**

EH-1388 Herbicide will provide full or partial control (suppression) of these lawn and turfgrass species

Bahiagrass	Canarygrass reed	Kikuyugrass
Bentgrass colonial creeping and redtop	Carpetgrass	Orchardgrass
Bermudagrass common and hybrid	Centipedegrass	Paspalum spp
Bluegrass annual rough and Kentucky	Fescues chewings hard red tall sheep	Ryegrass annual Italian perennial and hardy
Bromegrass downy and smooth	Foxtail green yellow creeping meadow giant	St Augustinegrass
Buffalograss	Johnsongrass	Zoysiagrass or Japanese lawnggrass

WOODY SPECIES

EH-1388 Herbicide will provide full or partial control (suppression) of these woody species

Alder	Hickory	Poison hemlock
Blackberry	Kudzu	Oak
Barberry	Lantana	Quaking aspen
Caenothus spp	Maple	Raspberry
Cherry	Multiflora rose	Sumac
Coyote brush	Mock orange †	Trumpet creeper
Dewberry	Manzanita	Wild blackberry
Elm	Pine	Western red cedar †
Elderberry	Poison ivy	Willow
Honeysuckle	Poison oak	†Not for use in California

ANNUAL AND PERENNIAL WEED SPECIES

EH-1388 Herbicide will provide full or partial control (suppression) of these annual and perennial weed species

Artichoke Jerusalem	Fiddleneck	Nimblewill	Tansy ragwort
Barnyardgrass	Field bindweed	Oleander †	Tansy mustard
Beggarweed	Field pennycress	Oxalis	Texas panicum
Bittercress	Field sandbur	Pampas grass	Thistle
Black medic	Filaree	Pennsylvania smartweed	Timothy
Broadleaf plantain	Florida pusley	Pigweed	Toadflax
Buckhorn plantain	Garden spurge	Plantain	Torpedograss
Bull thistle	Goosegrass	Prickly lettuce	Trailing lantana
Burclover	Green foxtail	Primrose	Tumble mustard
Canada thistle	Guineagrass	Prostrate spurge	Vaseygrass
Chickweed	Hemp sesbania	Puncturevine	Velvetleaf
Clover	Henbit	Purple cudweed	Virginia creeper
Cocklebur	Horsenettle	Purple nutsedge	White clover
Common groundsel	Horseradish	Purslane	Whitetop
Common mullien	Horseweed/marestail	Quackgrass	Wild barley
Common ragweed	Iceplant	Ragweed	Wild carrot
Coral bead	Ironweed	Red clover	Wild geranium
Crabgrass	Knapweed	Red sorrel †	Wild morningglory
Creeping beggarweed	Knawel	Redroot pigweed	Wild mustard
Creeping Charlie	Knotweed	Shepherdspurse	Wild oats
Curly dock	Lambsquarters	Smartweed	Wild sweet potato
Dallisgrass	London rocket	Smooth catsear	Witchgrass
Dandelion	Lovegrass	Smooth pigweed	Yarrow
Evening primrose	Mallow	Sowthistle	Yellow nutsedge

Fall panicum	Maresteed	Spotted spurge	Yellow oxalis
False dandelion	Milkweed	Spurge	Yellow starthistle
Fennel	Mousetear chickweed	Sunflower	and many other weeds
†Not for use in California			

WHEN TO USE [WHEN TO APPLY]

Apply to actively growing weeds when the vegetation is green. Use when the wind is calm to avoid drift. Make applications on warm sunny days when the daytime temperature is above 60°F. Weed control is accelerated by warm sunny weather. Weeds usually begin to wilt within hours with complete kill in 1 to 2 weeks. Rain or watering 2 hours after application will not wash away effectiveness. Hard-to-control weeds and thick lawn grasses (like bermudagrass) may require a second application.

HOW TO USE [HOW TO APPLY]

This product is formulated for spot treatment of individual weeds and grasses or for treatment of large areas as instructed on this label.

To apply this product use any of the following equipment:

- Plastic, aluminum or stainless steel tank sprayer
- Ortho Dial N Spray®
- Ortho Spray-Ette®
- A hose end sprayer recommended for herbicides
- Hand trigger pump sprayer
- **Do not use or store in galvanized or unlined steel sprayer, or through any irrigation system or sprinkling can**
- Apply to evenly and thoroughly wet weeds and grasses
- Use a tank type sprayer with a coarse spray nozzle for spot treatments
- Use the Ortho Dial N Spray® or other hose end sprayer to treat large areas such as for lawn renovation or to prepare sites for planting flower beds or vegetable gardens
- Treated weeds begin to wilt in about 2 to 4 days with complete kill of the weed including the root in 2 to 4 weeks
- EH-1388 Herbicide does not move or build up in the soil to affect untreated plants growing nearby
- Ornamental plantings can be made 3 days after treatment. Vegetable gardens can be seeded or planted 3 days after treatment
- For lawn replacement, lawns may be replanted 1 week after spraying
- Measure the correct amount of EH 1388 Herbicide and water to add to sprayer from the directions below




TABLE 1 SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	<ul style="list-style-type: none"> Use 2 25 fl oz of EH-1388 Herbicide in 1 gallon of water 	<ul style="list-style-type: none"> Use 4 5 fl oz of EH 1388 Herbicide in 1 gallon of water 	300 sq ft (30 ft x 10 ft)
Ortho Dial N Spray [®] Hose End Sprayer	<ul style="list-style-type: none"> Add 2 25 fl oz of EH-1388 Herbicide to sprayer jar DO NOT ADD WATER Set dial to 3 oz and apply 	<ul style="list-style-type: none"> Add 4 5 fl oz of EH-1388 Herbicide to sprayer jar DO NOT ADD WATER Set dial to 5 oz and apply 	300 sq ft
Ortho Spray Ette [®] Hose end Sprayer	<ul style="list-style-type: none"> Add 8 fl oz of EH 1388 Herbicide to sprayer jar Fill to 4 gallon mark 	<ul style="list-style-type: none"> Add 16 fl oz of EH 1388 Herbicide to sprayer jar Fill to 4 gallon mark 	1200 sq ft (30 ft x 40 ft)
Hand Trigger Sprayers	<ul style="list-style-type: none"> Use 1 tbs (0 5 fl oz) of EH 1388 Herbicide in 24 fl oz of water 	<ul style="list-style-type: none"> Use 2 tbs (1 0 fl oz) of EH-1388 Herbicide in 24 fl oz of water 	Use for spot treatments or single weeds

EQUIVALENT MEASUREMENTS
1 teaspoon (tsp)
1 tablespoon (tbs) = 3 teaspoons (tsp)
1 fluid ounce (fl oz) = 2 tablespoons (tbs)
2 fluid ounces (fl oz) = 4 tablespoons (tbs)

EH-1388 Herbicide in these container sizes will treat these approximate areas

CONTAINER SIZE	ANNUAL WEEDS (Treated Area, Sq Ft)	PERENNIAL WEEDS (Treated Area, Sq Ft)
1 pint	2 400 sq ft	1 200 sq ft
1 quart	4 800 sq ft	2 400 sq ft
2 quart (1/2 gallon)	9 600 sq ft	4 800 sq ft
1 gallon	19 200 sq ft	9 600 sq ft

CLEANING APPLICATION EQUIPMENT

To clean sprayer after use rinse spray equipment thoroughly and flush all sprayer components with water 3 times after use. Spray rinsate on bare soil or gravel surface. Do not apply rinsate to treated weeds. Only after a very thorough cleaning can the sprayer or spray equipment be used for application of other products.

WHERE TO USE

For General Weed Control Use near fences in paths patios sidewalks curbs and driveways around ornamental trees shrubs flower beds and plantings around buildings. Control weeds in brick and gravel walks. Use for trimming and edging landscapes. NOT recommended for spot weed treatment in lawns since glyphosate kills all green plants including lawn grasses.

Spot Treatments Spray directly on actively growing weeds in and around flower beds ornamental trees fruit and nut trees grapevines shrubs fences driveways and walkways. Do not spray fruit and nut trees or grapevines directly.

For Landscaping

- Prepare areas for planting of ornamentals trees shrubs desert landscapes rock gardens flower beds or similar plantings
- Delay mowing tilling or disturbing the treated vegetation for at least seven (7) days after application of this product for the best results However if the treated vegetation is not disturbed planting or transplanting of trees flowers shrubs bedding plants may be done at one (1) day after application

For Wildlife Areas Site Preparation

- Prepare areas for planting wildlife food
- Control undesirable and exotic plants to allow for recovery or planting of native species
- Make treatments to remove unwanted plants Take care to keep spray off desirable plants

For Lawn Renovation and Lawn Replacement

- Spot treatments or broadcast applications can be used to establish or renovate warm season and cool season turfgrass from seed or planting stock This product may be applied before planting seeding sprigging plugging or sodding
- Use EH 1388 Herbicide to remove the existing lawn and kill weeds and grasses Uniform and complete spray coverage of all existing vegetation is required
- Bermudagrass nutsedge torpedograss and deep rooted perennials may require a second treatment in 5 to 7 days
- For best results apply in spring or fall when daytime temperatures are at least 60° F
- Best results are obtained when one mowing is skipped before and after the application of this product
- If the lawn is dry water lawn before applying EH-1388 Herbicide Water again 2 to 3 days after application
- Hose end sprayers and tank sprayers are appropriate for localized spot treatments and broadcast (entire) applications on small turfgrass areas e g ~10 000 sq ft Refer to Table 1 for the spray preparation using hose end sprayers and tank sprayers
- Conventional sprayers fitted with spray booms and spray guns are appropriate for broadcast (entire) applications on large turfgrass areas e g ~20 000 sq ft Refer to Table 2 for the spray preparation using power spray equipment suitable for large turfgrass areas
- Seeding sprigging plugging or sodding may be done at seven (7) days after application of this product Prepare the soil for planting by raking roto-tilling vertical cutting or other tillage method
- Rake up and remove loosened thatch or debris
- Mix in starter fertilizer and soil amendments if needed Establish the lawn according to local cultural practices




TABLE 2 SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals/Acre	Amount of Product, fl oz/1000sqft	Spray Volume, Gals/1000sqft	Precautions, Restrictions & Comments
Lawn & turf seedbeds (prior to establishment)	Many annual and perennial grasses and broadleaf weeds including annual bluegrass crabgrass lambsquarters and ragweed	Annual weeds 1 8 - 3 6 Perennial weeds 5 5 9 5	≤ 25 ≤ 25	1 3 2 6 4 0 7 0	< 1 < 1	<p>SITE PREPARATION (preplant preemergence renovation) This product may be applied before planting or renovation of turf. Applications must be made prior to the emergence of the turf to avoid turf injury. For maximum control of existing vegetation, delay planting to determine if any regrowth from escape underground plant parts occurs. Where repeat treatments necessary, sufficient regrowth must be attained prior to application. For warm season grasses such as bermudagrass summer or fall applications provide best control.</p> <p>Do not disturb soil or underground plant parts before treatment.</p> <p>Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.</p> <p>This product does not provide residual control.</p>
	Bahiagrass	5 5 9 5	≤ 25	4 0 7 0	< 1	
	Bentgrass, colonial creeping	2 7 - 3 6	≤ 25	2 0 - 2 6	< 1	
	Bermudagrass common and hybrid	5 5 9 5	≤ 25	4 0 7 0	< 1	
	Bluegrass annual rough and Kentucky	1 8 3 6	≤ 25	1 3 2 6	< 1	
	Bromegrass downy and smooth	1 8 - 3 6	≤ 25	1 3 2 6	< 1	
	Orchardgrass	1 8 - 3 6	≤ 25	1 3 2 6	< 1	
	Canarygrass reed	3 6 - 5 5	≤ 25	2 6 4 0	< 1	
	Fescue tall	3 6 - 5 5	≤ 25	2 6 4 0	< 1	
	Fescues chewings hard red sheep	5 5 - 9 5	≤ 25	4 0 7 0	< 1	
	Johnsongrass and ryegrass annual	1 8 - 5 5	≤ 25	1 3 4 0	< 1	
	Kikuyugrass	3 6 - 5 5	≤ 25	2 6 4 0	< 1	
	Quackgrass and yellow nutsedge	3 6 - 5 5	≤ 25	2 6 4 0	< 1	
	Other perennial grasses listed on this label	5 5 9 5	≤ 25	4 0 7 0	< 1	

FARMSTEADS AND FENCEROWS EH 1388 Herbicide will control kochia multiflora rose thistle in areas where total vegetation control is desired Specific instructions for farmsteads are presented in Table 3

TABLE 3 SPRAY PREPARATION FOR FARMSTEADS USING POWER SPRAY EQUIPMENT

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals/Acre	Amount of Product, fl oz/1000sqft	Spray Volume, Gals/1000sqft	Precautions, Restrictions & Comments
Farmsteads where total vegetation control is desired around farmstead building foundations along fences hedgerows fencerows under fencelines around shelter belts and other area where nonselective weed control is desired	Annual and perennial weeds including crabgrass foxtail Johnsongrass kochia morningglory multiflora rose pigweed, trumpetreeper, thistle woody brush	Annual weeds 1 8 3 6 Perennial weeds 5 5 - 9 5 Woody brush & trees 5 5 - 9 5	3 40 3 40 3 - 40	1 3 2 6 4 0 7 0 4 0 7 0	≤ 1 0 ≤ 1 0 ≤ 1 0	<p>SITE renovation) PREPARATION (preplant preemergence</p> <p>BARE GROUND Repeated applications of this product may be used on emerged weeds to maintain bare ground</p> <p>NONSELECTIVE POSTEMERGENCE</p> <p>Annuals and Perennials Apply to foliage of emerged weeds Apply at early growth stages of annual weeds and when perennial weeds are approaching maturity</p> <p>Woody brush and trees Apply this product after full leaf expansion Use the higher rate for larger plants and/or dense areas of growth On vines use the higher rate for plants that have reached the woody stage of growth Best results are obtained when applications are made in late summer or fall after fruit formation</p> <p>In arid areas best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering</p> <p>Allow 7 or more days after application before tillage mowing or removal Repeat treatments may be necessary to control plants regenerating from underground parts or seed</p> <p>Reduced performance may result if fall treatments are made following a frost</p>

FOR BRUSH CONTROL

Foliar Sprays For Brush

- Cut back brush that is taller than 5 feet and then spray regrowth
- For maximum activity weeds should be growing vigorously at the time of the postemergent applications. The most favorable period for treatment occurs after full leaf stage in the spring and continues into early summer. Brush treated in the late fall may not be fully controlled.
- Spray uniformly the foliage of the vegetation to be controlled. Apply for full coverage and thorough brush/weed contact.
- Follow up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard-to control species including blackberry, poison oak, and kudzu. Another choice for effective brush control is Gordon's Cleanout® Brush and Stump Spray. Gordon's Cleanout® Brush and Stump Spray is recommended to control undesirable woody plants including buckbrush, poison ivy, multiflora rose, and sumac. Foliar applications of Gordon's Cleanout® Brush and Stump Spray are appropriate during the period of active growth of the brush. Cut stump, basal bark, and frill treatments of Gordon's Cleanout® Brush and Stump Spray may be applied any time of year.

Sprayers And Techniques For Brush Control

This product may be applied with the following application equipment:

- Hand held and high volume spray equipment: knapsack and backpack sprayers, pump-up pressure sprayers, spray guns, and other hand-held spray equipment used to direct the spray onto weed foliage.
- High volume spray equipment for spray to wet applications.

For Perennial Weeds And Brush

Apply a 3.5-9.0% vol/vol solution of this product. Use coarse sprays only.

TABLE 4 SPRAY PREPARATIONS FOR PERENNIAL WEEDS, VINES, AND BRUSH WITH HAND HELD AND HIGH VOLUME SPRAY APPLICATIONS

DESIRED VOLUME, GALLONS	AMOUNT OF PRODUCT (fl oz)	
	3.5% vol/vol	9.0% vol/vol
1 gallon	4.5 fl oz	11.5 fl oz
2 gallons	9.0 fl oz	23.0 fl oz
3 gallons	13.5 fl oz	34.5 fl oz

Cut Surface Stump Treatment

Cut stump treatments are effective on stumps with diameters larger than 3.4 inches in diameter.

- Cut trees and brush close to the soil surface. Apply this to the freshly cut surface of the stump *immediately* after cutting. Delays in application may reduce the effectiveness.
- Apply the *undiluted product* for full and thorough coverage of the cut surface. Treat the entire circumference of the stump by drenching the cambium (the outermost wood between the pith and the bark).
- Treatment will control or suppress many types of woody brush and trees such as alder, bluegum, eucalyptus, madrone, oak, giant reed, saltcedar, sweetgum, and tan oak.
- The most favorable period for cut stump treatment occurs after full leaf stage in the spring and continues into early summer. Brush and trees treated in the late fall may not be fully controlled.

- Follow up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard to-control species. Another choice for effective control with a cut-surface stump treatment is Gordon's Cleanout® Brush and Stump Spray.

For Vine Control

- If vines are growing up poles, fences, or tree trunks with mature bark, cut vines to a height of 3 to 4 feet and spray vines thoroughly.
- If vines are climbing shrubs or tree trunks with green bark, cut vines at base and treat as directed for stumps or spray regrowth. If spraying regrowth, shield shrubs and green bark from spray drift with a piece of cardboard or plastic.

Refer to Tables 4 and 5 to determine the amount of EH-1388 Herbicide to mix with water for vine and perennial weed control.

TABLE 5 SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS FOR VINES AND PERENNIAL WEED CONTROL

SPRAYER TYPE	MIXING INSTRUCTIONS	TREAT
	Tough Weed Control Perennials or Vines	
Standard Tank Sprayer	• Use 4.5 fl oz (9 tbs) of EH-1388 Herbicide in 1 gallon of water	300 sq ft (30 ft x 10 ft)
Ortho Dial N Spray® Hose End Sprayer	• Add 4.5 fl oz (9 tbs) of EH-1388 Herbicide to sprayer jar • DO NOT ADD WATER • Set dial to 5 oz and apply	300 sq ft
Ortho Spray-Ette® Hose end Sprayer	• Add 16 fl oz of EH-1388 Herbicide to sprayer jar • Fill to 4 gallon mark	1200 sq ft (30 ft x 40 ft)
Hand Trigger Sprayers	• Use 1.0 fl oz (2 tbs) of EH 1388 Herbicide in 24 fl oz of water	Use for spot treatments or single weeds

GARDENS (VEGETABLES, HERBS, SMALL FRUITS AND BERRIES)

This product may be applied before planting or transplanting of vegetables (garden crops), herbs, and small fruits and berries. This product may be used as a broad spectrum postemergent herbicide prior to planting to control annual and perennial weeds including field bindweed, Johnsongrass, or dallisgrass.

Spray the undesirable vegetation on a spray-to-wet basis. Uniform and complete spray coverage of all existing vegetation is required.

RESEEDING AND REPLANTING IN SPRAYED AREAS

ORNAMENTALS AND FLOWERS Reseed or replant one (1) day after application.

VEGETABLES Wait three (3) days before planting or transplanting the vegetables listed below.

Beans	Cucumber	Mango melon	Radish
Beet greens	Eggplant	Melons	Rhubarb
Broccoli	Endive	Muskmelon	Rutabaga
Brussels sprout	Garlic	Mustard greens	Spinach
Cabbage	Gourds	Okra	Summer squash
Cantaloupe	Honeydew melon	Onion	Sweet potato
Carrot	Honeyball melon	Rape greens	Tomatillo
Cauliflower	Horseradish	Parsley	Tomato
Casaba melon	Jerusalem artichoke	Parsnips	Turnip
Chard (Swiss)	Kale	Peas	Watercress
Chicory	Kohlrabi	Pepper	Watermelon
Collards	Leek	Persian melon	Winter squash

Corn
Crenshaw melon

Lettuce

Potato
Pumpkin

Yams

*Tomatoes planted from seed only Wait thirty (30) days for transplants For vegetables not listed or for tomato transplants wait thirty (30) days before planting

SMALL FRUITS AND BERRIES Blackberry black raspberry blueberry boysenberry currant dewberry elderberry gooseberry huckleberry loganberry olallieberry and red raspberry may be planted by seed rooted cutting or transplant 7 days after spraying EH 1388 Herbicide Wait thirty (30) days after application before planting of all other unlisted small fruits and berries

FRUIT TREES, NUT TREES, CANEBERRIES OR GRAPEVINES If this product is used to control weeds around fruit or nut trees caneberreries or grapevines wait twenty-one (21) days before harvesting and eating the fruits or nuts

Use the following table to determine the amount of EH 1388 Herbicide to mix with water for use in gardens (vegetables herbs small fruits and berries) Close cap tightly after use

TABLE 6 SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS FOR USE IN GARDENS, (VEGETABLES, HERBS, SMALL FRUITS, AND BERRIES)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	• Use 2 25 fl oz of EH 1388 Herbicide in 1 gallon of water	• Use 4 5 fl oz of EH 1388 Herbicide in 1 gallon of water	300 sq ft (30 ft x 10 ft)
Ortho Dial N Spray [®] Hose End Sprayer	• Add 2 25 fl oz of EH 1388 Herbicide to sprayer jar • DO NOT ADD WATER • Set dial to 3 oz and apply	• Add 4 5 fl oz of EH 1388 Herbicide to sprayer jar • DO NOT ADD WATER • Set dial to 5 oz and apply	300 sq ft
Ortho Spray Ette [®] Hose end Sprayer	• Add 8 fl oz of EH 1388 Herbicide to sprayer jar • Fill to 4 gallon mark	• Add 16 fl oz of EH 1388 Herbicide to sprayer jar • Fill to 4 gallon mark	1200 sq ft (30 ft x 40 ft)
Hand Trigger Sprayers	• Use 0 5 fl oz (1 tbs) of EH-1388 Herbicide in 24 fl oz of water	• Use 1 fl oz (2 tbs) of EH-1388 Herbicide in 24 fl oz of water	Use for spot treatments or single weeds

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE WHICH ARE EXPRESSLY DISCLAIMED This limited warranty does not extend to the use of the product inconsistent with label instructions warnings or cautions or to use of the product under abnormal conditions such as drought excessive rainfall tornadoes hurricanes etc These factors are beyond the control of the manufacturer or the seller Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages and shall not include indirect or consequential damages such as loss of profits or values except as otherwise provided by law

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner

[brand name] is a trademark of PBI/GORDON Corporation

Ortho Ortho Dial N Spray[®] and Ortho Spray-Ette[®] are registered trademarks of Monsanto Company

EH-1388 HERBICIDE

Alternate Brand Names GroundWork Concentrate 25% Weed & Grass Killer

Sublabel 2 This sublabel is intended for the GroundWork brand

ACTIVE INGREDIENT

Isopropylamine salt of glyphosate* 25.0%

OTHER INGREDIENTS 75.0%

TOTAL 100.0%

*N (phosphonomethyl) glycine

KEEP OUT OF REACH OF CHILDREN

CAUTION


See [back/side/inside panel or booklet]
for additional Precautionary Statements and First Aid
or [Peel open for use directions and further information]

NET CONTENTS (16 fl oz , Quart, 40 fl oz ,
1/2 Gallon, 1 Gallon)

954/AP072911
EPA Reg No 2217-846
EPA Est No 2217 KS-2
MANUFACTURED BY



Telephone 800 821 7925





**READ THE ENTIRE LABEL FIRST OBSERVE ALL PRECAUTIONS
AND FOLLOW DIRECTIONS CAREFULLY**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION Causes moderate eye irritation Harmful if absorbed through skin Harmful if swallowed
Avoid contact with skin eyes or clothing Wash thoroughly with soap and water

First Aid

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 - 20 minutes • Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible • Call a poison control center or doctor for treatment advice
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment	

ENVIRONMENTAL HAZARDS

Do not apply directly to water Do not contaminate water when disposing of equipment washwater and rinsate

PHYSICAL OR CHEMICAL HAZARDS

Combustible Do not use or store near heat or open flame

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Keep children and pets out of treated areas until sprays have dried

STORAGE & DISPOSAL

Do not contaminate water food or feed by storage and disposal

PESTICIDE STORAGE Store in original container in a locked storage area inaccessible to children and domestic animals Protect container from freezing

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons]

PESTICIDE DISPOSAL AND CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container **If empty** Place in trash or offer for recycling if available **If partly filled** Call your local

solid waste agency or www.epa.gov for disposal instructions. Never use unused product down any indoor or outdoor drain.

HOW IT WORKS

Facts and Tips

- EH 1388 Herbicide is a multipurpose herbicide designed to control a broad spectrum of emerged annual and perennial weed species. EH-1388 Herbicide will control unwanted species of grass, sedge, broadleaf weeds, and certain woody species around trees, shrubs, fences.
- This product kills plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves to the roots.
- This product is nonselective and will kill or injure all vegetation contacted by the spray. Do not spray plants or grasses you like—they may die too. Not recommended for spot weed control in lawns since this product kills lawn grasses.
- No soil residual activity—any product not absorbed by plants breaks down into natural materials without moving in or on the soil to untreated plants.
- Plants that have not emerged at the time of application will not be controlled.
- Time to symptoms—best results will be obtained when the weed species are actively growing. Visible effects are gradual—wilting and yellowing, advancing to complete browning and root destruction. Weeds usually yellow and wilt within 2 to 4 days and plant death occurs in 1 to 2 weeks.

Product Description and Spray Coverage This product is a nonselective herbicide that will kill all vegetation contacted. This herbicide kills unwanted plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves to the roots. Avoid direct spray or drift onto desirable plants, lawn grasses, shrubs, and trees. Protect desirable plants and vegetation with a piece of plastic or cardboard. If desirable plants are accidentally sprayed, wash plants with water immediately. Any glyphosate not absorbed by the plant is broken down into natural materials. This product has no soil residual activity and thus will not affect nearby untreated plants.

WHERE TO USE

For General Weed Control and Spot Spraying

- Use to prepare sites for large plantings, for planting flower beds, for planting fruit trees, ornamentals, vegetables, garden sites, and to renovate lawns.
- Along and on sidewalks, gravel/brick walkways, paths, curbs, paved areas (private roads and streets), driveways, parking areas, recreational areas.
- Along ornamental lawns and turfgrass. Edging along/around lawns/yards and landscape plantings.
- In areas for lawn replacement or lawn renovation/replacement.
- Around or near ornamental/shade trees, fruit and nut trees, grapevines, shrubs, groundcovers, flower beds, mulched landscape beds.
- For garden plot preparation, vegetable plot preparation, and small fruits and berries plot preparation.
- For flower garden preparation.
- Along fences, hedgerows, under fencelines.
- Around rocks, signposts, light posts, or on fringe areas.
- Around or along buildings/structures, foundations, patios, houses, mobile homes, vacant lots, urban areas, domestic outdoor premises, and other areas associated with household or home life.
- Urban areas where total vegetation control is desired.
- Farmsteads where total vegetation control is desired, around farmstead building foundations, along fences, hedgerows, fencerows, under fencelines, around shelter belts, and other areas where nonselective weed control is desired.

TOUGH WEED CONTROL (ANNUALS, PERENNIALS, BRUSH)

CONTROLS WEEDS AND GRASSES INCLUDING LAWN AND TURFGRASS SPECIES

EH 1388 Herbicide will provide full or partial control (suppression) of these lawn and turfgrass species

Bahiagrass	Canarygrass reed	Kikuyugrass
Bentgrass colonial creeping and redtop	Carpetgrass	Orchardgrass
Bermudagrass common and hybrid	Centipedegrass	Paspalum spp
Bluegrass annual rough and Kentucky	Fescues chewings hard red tall sheep	Ryegrass annual Italian perennial and hardy
Bromegrass downy and smooth	Foxtail green yellow creeping meadow giant	St Augustinegrass
Buffalograss	Johnsongrass	Zoysiagrass or Japanese lawnglass

WOODY SPECIES

EH-1388 Herbicide will provide full or partial control (suppression) of these woody species

Alder	Hickory	Poison hemlock
Blackberry	Kudzu	Oak
Barberry	Lantana	Quaking aspen
Caenothus spp	Maple	Raspberry
Cherry	Multiflora rose	Sumac
Coyote brush	Mock orange †	Trumpet creeper
Dewberry	Manzanita	Wild blackberry
Elm	Pine	Western red cedar †
Elderberry	Poison ivy	Willow
Honeysuckle	Poison oak	†Not for use in California

ANNUAL AND PERENNIAL WEED SPECIES

EH 1388 Herbicide will provide full or partial control (suppression) of these annual and perennial weed species

Artichoke Jerusalem	Fiddleneck	Nimblewill	Tansy ragwort
Barnyardgrass	Field bindweed	Oleander †	Tansy mustard
Beggarweed	Field pennycress	Oxalis	Texas panicum
Bittercress	Field sandbur	Pampas grass	Thistle
Black medic	Filaree	Pennsylvania smartweed	Timothy
Broadleaf plantain	Florida pusley	Pigweed	Toadflax
Buckhorn plantain	Garden spurge	Plantain	Torpedograss
Bull thistle	Goosegrass	Prickly lettuce	Trailing lantana
Burclover	Green foxtail	Primrose	Tumble mustard
Canada thistle	Guineagrass	Prostrate spurge	Vaseygrass
Chickweed	Hemp sesbania	Puncturevine	Velvetleaf
Clover	Henbit	Purple cudweed	Virginia-creeper
Cocklebur	Horsenettle	Purple nutsedge	White clover
Common groundsel	Horseradish	Purslane	Whiteweed
Common mullien	Horseweed/marestail	Quackgrass	Wild barley
Common ragweed	Iceplant	Ragweed	Wild carrot
Coral bead	Ironweed	Red clover	Wild geranium
Crabgrass	Knapweed	Red sorrel †	Wild morning glory
Creeping beggarweed	Knawel	Redroot pigweed	Wild mustard
Creeping Charlie	Knotweed	Shepherdspurse	Wild oats
Curly dock	Lambsquarters	Smartweed	Wild sweet potato
Dallisgrass	London rocket	Smooth catsear	Witchgrass
Dandelion	Lovegrass	Smooth pigweed	Yarrow
Evening primrose	Mallow	Sowthistle	Yellow nutsedge
Fall panicum	Mayweed	Spotted spurge	Yellow oxalis

2/28

False dandelion	M. eed	Spurge	Yellow starthistle
Fennel	Mouseear chickweed	Sunflower	and many other weeds
†Not for use in California			

WHEN TO USE [WHEN TO APPLY]

Apply to actively growing weeds when the vegetation is green. Use when the wind is calm to avoid drift. Make applications on warm sunny days when the daytime temperature is above 60°F. Weed control is accelerated by warm sunny weather. Weeds usually begin to wilt within hours with complete kill in 1 to 2 weeks. Rain or watering 2 hours after application will not wash away effectiveness. Hard-to-control weeds and thick lawn grasses (like bermudagrass) may require a second application.

HOW TO USE [HOW TO APPLY]

This product is formulated for spot treatment of individual weeds and grasses or for treatment of large areas as instructed on this label.

To apply this product use any of the following equipment:

- Plastic, aluminum or stainless steel tank sprayer
- Ortho Dial N Spray®
- Ortho Spray Ette®
- A hose end sprayer recommended for herbicides
- Hand-trigger pump sprayer
- **Do not use or store in galvanized or unlined steel sprayer, or through any irrigation system or sprinkling can**
- Apply to evenly and thoroughly wet weeds and grasses
- Use a tank type sprayer with a coarse spray nozzle for spot treatments
- Use the Ortho Dial N Spray® or other hose end sprayer to treat large areas such as for lawn renovation or to prepare sites for planting flower beds or vegetable gardens
- Treated weeds begin to wilt in about 2 to 4 days with complete kill of the weed including the root in 2 to 4 weeks
- EH-1388 Herbicide does not move or build up in the soil to affect untreated plants growing nearby
- Ornamental plantings can be made 3 days after treatment. Vegetable gardens can be seeded or planted 3 days after treatment
- For lawn replacement, lawns may be replanted 1 week after spraying
- Measure the correct amount of EH-1388 Herbicide and water to add to sprayer from the directions below




TABLE 1 SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	<ul style="list-style-type: none"> Use 2 25 fl oz of EH-1388 Herbicide in 1 gallon of water 	<ul style="list-style-type: none"> Use 4 5 fl oz of EH-1388 Herbicide in 1 gallon of water 	300 sq ft (30 ft x 10 ft)
Ortho Dial N Spray [®] Hose End Sprayer	<ul style="list-style-type: none"> Add 2 25 fl oz of EH-1388 Herbicide to sprayer jar DO NOT ADD WATER Set dial to 3 oz and apply 	<ul style="list-style-type: none"> Add 4 5 fl oz of EH 1388 Herbicide to sprayer jar DO NOT ADD WATER Set dial to 5 oz and apply 	300 sq ft
Ortho Spray Ette [®] Hose end Sprayer	<ul style="list-style-type: none"> Add 8 fl oz of EH-1388 Herbicide to sprayer jar Fill to 4 gallon mark 	<ul style="list-style-type: none"> Add 16 fl oz of EH 1388 Herbicide to sprayer jar Fill to 4 gallon mark 	1200 sq ft (30 ft x 40 ft)
Hand Trigger Sprayers	<ul style="list-style-type: none"> Use 1 tbs (0 5 fl oz) of EH 1388 Herbicide in 24 fl oz of water 	<ul style="list-style-type: none"> Use 2 tbs (1 0 fl oz) of EH-1388 Herbicide in 24 fl oz of water 	Use for spot treatments or single weeds

EQUIVALENT MEASUREMENTS
1 teaspoon (tsp)
1 tablespoon (tbs) = 3 teaspoons (tsp)
1 fluid ounce (fl oz) = 2 tablespoons (tbs)
2 fluid ounces (fl oz) = 4 tablespoons (tbs)

EH-1388 Herbicide in these container sizes will treat these approximate areas

CONTAINER SIZE	ANNUAL WEEDS (Treated Area, Sq Ft)	PERENNIAL WEEDS (Treated Area, Sq Ft)
1 pint	2 400 sq ft	1 200 sq ft
1 quart	4 800 sq ft	2 400 sq ft
2 quart (1/2 gallon)	9 600 sq ft	4 800 sq ft
1 gallon	19 200 sq ft	9 600 sq ft

CLEANING APPLICATION EQUIPMENT

To clean sprayer after use rinse spray equipment thoroughly and flush all sprayer components with water 3 times after use Spray rinsate on bare soil or gravel surface Do not apply rinsate to treated weeds Only after a very thorough cleaning can the sprayer or spray equipment be used for application of other products

WHERE TO USE

For General Weed Control Use near fences in paths patios sidewalks curbs and driveways around ornamental trees shrubs flower beds and plantings around buildings Control weeds in brick and gravel walks Use for trimming and edging landscapes NOT recommended for spot weed treatment in lawns since glyphosate kills all green plants including lawn grasses

Spot Treatments Spray directly on actively growing weeds in and around flower beds ornamental trees fruit and nut trees grapevines shrubs fences driveways and walkways Do not spray fruit and nut trees or grapevines directly

For Landscaping

- Prepare areas for planting of ornamentals trees shrubs desert landscapes rock gardens flower beds or similar plantings
- Delay mowing tilling or disturbing the treated vegetation for at least seven (7) days after application of this product for the best results However if the treated vegetation is not disturbed planting or transplanting of trees flowers shrubs bedding plants may be done at one (1) day after application

For Lawn Renovation and Lawn Replacement

- Spot treatments or broadcast applications can be used to establish or renovate warm season and cool season turfgrass from seed or planting stock. This product may be applied before planting, seeding, sprigging, plugging, or sodding.
- Use EH-1388 Herbicide to remove the existing lawn and kill weeds and grasses. Uniform and complete spray coverage of all existing vegetation is required.
- Bermudagrass, nutsedge, torpedograss, and deep-rooted perennials may require a second treatment in 5 to 7 days.
- For best results, apply in spring or fall when daytime temperatures are at least 60° F.
- Best results are obtained when one mowing is skipped before and after the application of this product.
- If the lawn is dry, water the lawn before applying EH-1388 Herbicide. Water again 2 to 3 days after application.
- Hose end sprayers and tank sprayers are appropriate for localized spot treatments and broadcast (entire) applications on small turfgrass areas, e.g., ~10,000 sq ft. Refer to Table 1 for the spray preparation using hose end sprayers and tank sprayers.
- Conventional sprayers fitted with spray booms and spray guns are appropriate for broadcast (entire) applications on large turfgrass areas, e.g., ~20,000 sq ft. Refer to Table 2 for the spray preparation using power spray equipment suitable for large turfgrass areas.
- Seeding, sprigging, plugging, or sodding may be done at seven (7) days after application of this product. Prepare the soil for planting by raking, roto-tilling, vertical cutting, or other tillage method.
- Rake up and remove loosened thatch or debris.
- Mix in starter fertilizer and soil amendments if needed. Establish the lawn according to local cultural practices.




TABLE 2 SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals /Acre	Amount of Product, fl oz/1000sqft	Spray Volume, Gals/1000sqft	Precautions, Restrictions & Comments
Lawn & turf seedbeds (prior to establishment)	Many annual and perennial grasses and broadleaf weeds including annual bluegrass crabgrass lambsquarters and ragweed	Annual weeds 1 8 3 6 Perennial weeds 5 5 9 5	≤ 25 ≤ 25	1 3 2 6 4 0 7 0	< 1 < 1	<p>SITE PREPARATION (preplant, preemergence, renovation) This product may be applied before planting or renovation of turf Applications must be made prior to the emergence of the turf to avoid turf injury For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs Where repeat treatments necessary, sufficient regrowth must be attained prior application For warm season grasses such as bermudagrass summer or fall applications provide best control</p> <p>Do not disturb soil or underground plant parts before treatment Tillage or renovation techniques such as vertical mowing coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts</p> <p>This product does not provide residual control</p>
	Bahia grass	5 5 - 9 5	≤ 25	4 0 7 0	< 1	
	Bentgrass colonial creeping	2 7 - 3 6	≤ 25	2 0 2 6	< 1	
	Bermudagrass common and hybrid	5 5 - 9 5	≤ 25	4 0 7 0	< 1	
	Bluegrass annual rough and Kentucky	1 8 - 3 6	≤ 25	1 3 2 6	< 1	
	Bromegrass downy and smooth	1 8 - 3 6	≤ 25	1 3 2 6	< 1	
	Orchardgrass	1 8 - 3 6	≤ 25	1 3 2 6	< 1	
	Canarygrass reed	3 6 - 5 5	≤ 25	2 6 - 4 0	< 1	
	Fescue tall	3 6 - 5 5	≤ 25	2 6 4 0	< 1	
	Fescues chewings hard red sheep	5 5 - 9 5	≤ 25	4 0 7 0	< 1	
	Johnsongrass and ryegrass annual perennial	1 8 - 5 5	≤ 25	1 3 4 0	< 1	
	Kentucky bluegrass	3 6 - 5 5	≤ 25	2 6 4 0	< 1	
	Quackgrass and yellow nutsedge	3 6 - 5 5	≤ 25	2 6 4 0	< 1	
	Other perennial grasses listed on this label	5 5 - 9 5	≤ 25	4 0 - 7 0	< 1	

FARMSTEADS AND FENCEROWS EH 1388 Herbicide will control kochia multiflora rose thistle in areas where total vegetation control is desired Specific instructions for farmsteads are presented in Table 3

TABLE 3 SPRAY PREPARATION FOR FARMSTEADS USING POWER SPRAY EQUIPMENT

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals /Acre	Amount of Product, fl oz/1000sqft	Spray Volume, Gals/1000sqft	Precautions, Restrictions & Comments
Farmsteads where total vegetation control is desired around farmstead building foundations along fences hedgerows fencerows under fencelines around shelter belts and other area where nonselective weed control is desired	Annual and perennial weeds including crabgrass foxtail Johnsongrass kochia morningglory multiflora rose pigweed trumpetcreeper thistle woody brush	Annual weeds 1 8 3 6 Perennial weeds 5 5 - 9 5 Woody brush & trees 5 5 - 9 5	3 - 40 3 - 40 3 - 40	1 3 - 2 6 4 0 7 0 4 0 7 0	≤ 1 0 ≤ 1 0 ≤ 1 0	<p>SITE PREPARATION (preplant preemergence renovation) Repeated applications of this product may be used on emerged weeds to maintain bare ground</p> <p>NONSELECTIVE POSTEMERGENCE Annuals and Perennials Apply to foliage of emerged weeds Apply at early growth stages of annual weeds and when perennial weeds are approaching maturity Woody brush and trees Apply this product after full leaf expansion Use the higher rate for larger plants and/or dense areas of growth On vines use the higher rate for plants that have reached the woody stage of growth Best results are obtained when applications are made in late summer or fall after fruit formation In and areas best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering Allow 7 or more days after application before tillage mowing or removal Repeat treatments may be necessary to control plants regenerating from underground parts or seed Reduced performance may result if fall treatments are made following a frost</p>

TABLE 4 SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS FOR USE IN GARDENS, (VEGETABLES, HERBS, SMALL FRUITS, AND BERRIES)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	<ul style="list-style-type: none"> Use 2 25 fl oz of EH-1388 Herbicide in 1 gallon of water 	<ul style="list-style-type: none"> Use 4 5 fl oz of EH-1388 Herbicide in 1 gallon of water 	300 sq ft (30 ft x 10 ft)
Ortho Dial N Spray [®] Hose End Sprayer	<ul style="list-style-type: none"> Add 2 25 fl oz of EH 1388 Herbicide to sprayer jar DO NOT ADD WATER Set dial to 3 oz and apply 	<ul style="list-style-type: none"> Add 4 5 fl oz of EH-1388 Herbicide to sprayer jar DO NOT ADD WATER Set dial to 5 oz and apply 	300 sq ft
Ortho Spray Ette [®] Hose end Sprayer	<ul style="list-style-type: none"> Add 8 fl oz of EH 1388 Herbicide to sprayer jar Fill to 4 gallon mark 	<ul style="list-style-type: none"> Add 16 fl oz of EH-1388 Herbicide to sprayer jar Fill to 4 gallon mark 	1200 sq ft (30 ft x 40 ft)
Hand Trigger Sprayers	<ul style="list-style-type: none"> Use 0 5 fl oz (1 tbs) of EH 1388 Herbicide in 24 fl oz of water 	<ul style="list-style-type: none"> Use 1 fl oz (2 tbs) of EH 1388 Herbicide in 24 fl oz of water 	Use for spot treatments or single weeds

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE WHICH ARE EXPRESSLY DISCLAIMED This limited warranty does not extend to the use of the product inconsistent with label instructions warnings or cautions or to use of the product under abnormal conditions such as drought excessive rainfall tornadoes hurricanes etc These factors are beyond the control of the manufacturer or the seller Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values except as otherwise provided by law

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner

[brand name] is a trademark of PBI/GORDON Corporation

Ortho Ortho Dial N Spray[®] and Ortho Spray-Ette[®] are registered trademarks of Monsanto Company

APPENDIX

I Advertising claims that may be presented on container labeling of supplemental registrants

- kills common broadleaf weeds and grasses
- Kills weeds and grasses
- for general multipurpose weed control
- water based formula
- no odor
- for nonselective post emergent control of unwanted vegetation
- contains glyphosate
- treats approximately # _____ sq ft /treats over # _____sq ft (dependent upon container size)
- systemic weed and grass killer for spot treatment of unwanted weeds and grasses

- ◆ eliminates undesirable vegetation in 2-4 weeks
- ◆ kills roots and all in 2-4 weeks
- ◆ kills to the root/down to the root/roots and all
- ◆ no residual activity in soil
- ◆ for use around flower beds trees and shrubs fences walks mulch beds gravel and driveways
- ◆ for use around flower beds trees shrubs fences and walks
- ◆ kills weeds near/under fences in driveways on patios in paths sidewalks cracks walkways gardens trees shrubs flower beds mulch beds graveled areas and curbs
- ◆ controls weeds and unwanted vegetation around trees shrubs and in flower beds
- ◆ use to prepare sites for planting fruit trees shrubs and ornamentals
- ◆ multipurpose grass and broadleaf weed control
- ◆ for outdoor use only
- ◆ use for lawn renovation
- ◆ can reseed lawns 7 days after treatment
- ◆ can be applied with a tank sprayer or hose end sprayer
- ◆ visible results in 3 days
- ◆ complete kill down to the root in 3 to 4 weeks
- ◆ [More Concentrated][Makes ____ gallons of spray][Value pack]
- ◆ [Rain-Proof in hours][Rain-Proof in less than ____ hours]
- ◆ Won't wash off
- ◆ No root No weed No problem
- ◆ Gets to the root of the problem
- ◆ Use for planting site preparation for large plantings such as flowerbeds fruit trees ornamentals vegetables garden sites and to renovate lawns
- ◆ Can be used around small fruits and vegetables
- ◆ Kills weeds you don't want in the garden
- ◆ Gets rid of weeds in flowerbeds and along fences around trees shrubs driveways walkways in and around fences patios paths sidewalk cracks walkways and trails
- ◆ Can be used for habitat restoration controlling exotic vegetation and for site preparation of wildlife food plots
- ◆ [For product information and a complete list of weeds controlled] [For more information] [For questions or comments] [call toll free 1-8xx-yyy-zzzz or visit (website address)]
- ◆ Covers up to XXX[*]
- ◆ Makes up to [30] gallons[*]
- ◆ * When mixed XXX of product with YYY of water
- ◆ * When mixed at the rate of XXX per YYY of water
- ◆ Lawn reseeding in 3 to 7 days
- ◆ Lawn renovation
- ◆ Garden renovation
- ◆ Fenceline cleanup and more
- ◆ Rain fast in hours
- ◆ Even kills the roots
- ◆ 25% More Free!
- ◆ Contains Glyphosate and Surfactant
- ◆ Lawn Renovation—Driveways—Fenceline and More!
- ◆ For outdoor use
- ◆ Great for garden preparation
- ◆ Allows grass reseeding in one week
- ◆ Glyphosate 25%

