

Mushroom Zineb Dust

(15% Active)

for

Control of Mycogone (Bubble), Verticillium (Dry Bubble, Brown Spot) and Dactylium (Mildew)

DIRECTIONS FOR USE

MSCo No. 15X Zineb Dust is used at the rate of \(\frac{1}{2}\) to \(\frac{3}{4}\) lb. per 4000 sq. ft. of bed area and is applied with a good duster. Begin applications soon after casing soil is applied and continue at 3 to 7 day intervals until small buttons begin to form: Also between breaks as required.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Keep container closed and in a dry place. This material may cause irritation of nose and throat; protect the operator's eyes, nose and throat by use of respirator and goggles. Avoid inhalation as material is harmful if taken internally. Do not breathe the dust. Avoid contact with skin or clothing. Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms of the label and subject thereto, the buyer assumes any risk to persons or property arising out of use of handling and accepts the product on these conditions.

ACTIVE INGREDIENTS:

Zineb (Zinc Ethylene Bisdithiocarbamate) (Metallic Zinc content 3.55) 15.0°

85.0%

100.0℃

INERT INGREDIENTS

Manufactured by

MUSHROOM SUPPLY CO

TOUGHKENAMON, PA.

USDA Reg. No 237-9

ACCEPTED

JUL 27 1967

O ATTACHED CONNEITS