

ACCEPTED
FEB 19, 69

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 239-16. SUBJECT
TO ATTACHED COMMENTS.



VOLCK oil spray

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

Shake well. Use a forceful spray, wetting pests and all parts of plants thoroughly. Do not store diluted spray. Make new dilution for each use.

FOR DORMANT USE—To Control Scale Insects, Red Spiders, Mites, and Mealybugs on Dormant Shrubs, Evergreens (note caution), Woody Plants, Fruit Trees and Shade Trees: ½ pint (5½ ozs.) per gal. water.

GROWING SEASON—Scale Insects, Red Spiders, Mites, Mealybugs and White Fly Larvae on Citrus, Fruit Trees, Shade Trees, Evergreens, Camellias, Azaleas, Hardy Shrubs: 5 Tablespoonfuls (2½ ozs.) per gal. water.

COMBINATION WINTER SPRAY

FRUIT TREES, ROSES AND ORNAMENTALS (Dormant, deciduous only)—for Control of San Jose Scale, Rose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale and Overwintering Insect Eggs: **FOR BEST RESULTS:** Use 5 Tablespoonfuls VOLCK Oil Spray and 5 Tablespoonfuls ORTHORIX Spray in 1 gallon water. Keep agitated during spraying. Note cautions on ORTHORIX Spray label before use.

COMBINATION WITH ORTHO Malathion 50 Insect Spray on Citrus, Avocados, Mangoes, Gardenias and Ixora for Control of Scale Insects, Red Spider Mites, Mealybugs, White Flies, Aphids, Thrips and for the removal of Sooty Mold Fungus through control of White Fly Larvae—Use 1 Tablespoonful VOLCK Oil Spray and 1 Tablespoonful ORTHO Malathion 50 Insect Spray for each gallon of diluted spray. May be applied on Avocados up to within 7 days of harvest; to Grapefruit, Lemons, Limes and Oranges up to within 7 days of harvest but not during full bloom; and to Mangoes up to within 2 days of harvest.

AS A STICKER-SPREADER—For ISOTOX Insect Spray, Lead Arsenate and Certain Other Insecticides and Fungicides used at regular strength: Use 4 teaspoonfuls (¾ oz.) per gal. water to increase efficiency of a spray treatment, provide better contact with insects or plant surfaces and to retard weathering off of the residual materials applied.

GREENHOUSES: To control Scale Insects, Red Spiders, Mites, White Fly, Mealybugs on Hardy Shrubs, Gardenias, Camellias and Azaleas—2 Tablespoonfuls (1 oz.) per gal. water. Allow at least 60 days between oil sprays and limiting to 3 applications per year.

PALM TREES: Scale Insects, Mealybugs, Red Spider Mites—3 Tablespoonfuls (1½ ozs.) per gal. water. Note Caution. **AZALEAS, CAMELLIAS:** Scale Insects, Red Spider Mites, Mealybugs, White Fly Larvae, Lace Bug Larvae—5 Tablespoonfuls (2½ ozs.) per gal. water. Apply in spring after blooming and in fall prior to blooming.

IF SPRAY-ETTE Garden Hose Sprayer is used—Follow directions on the bottle.


CAUTION: Harmful if swallowed. Wash hands after use. Do not use VOLCK Oil Spray on foliage within 30 days before or after using Sulfur. Avoid spraying when temperature is excessively high or when plants are suffering from drought. Oil sprays should not be applied to broad-leaved evergreens or palms when freezing or near-freezing temperatures may be expected. Do not use on Maiden-hair Ferns, Cocos Palms, Maples, etc., or non-woody plants unless specified. VOLCK Oil Spray may cause discoloration to Blue Spruce. Do not use in combination with or closely following treatment with ORTHOCIDE Garden Fungicide.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



CHEVRON CHEMICAL COMPANY
ORTHODIVISION—San Francisco, Calif 94120

RICHMOND, CALIF. Atlanta, Georgia Des Plaines, Illinois
Form 571-1 USDA Reg. No. 235-16 Product No. 116 Made in U.S.A.



VOLCK oil spray

DORMANT & GROWING SEASON INSECT SPRAY



CONTROLS SCALE INSECTS
MEALYBUGS - WHITE FLIES
RED SPIDER MITES and
Certain Other Garden Pests

239-16

02/19/1969

111