

239-16

2-11-71


1 of 2

Best Available copy


ORTHO

ORTHO

\$34

VOLCK®
Oil
Spray
VOLCK®
Oil Spray
EXTRACTED REFINED
OIL SPRAY
INSECTICIDE
For Use Against
Scale Insects
Red Spiders
Mealybugs
White Flies
and Certain Other Pests
ALSO
MAKES AN EXCELLENT
LEAF POLISH
PERMANENT & PERSISTENT SEASON
INSECT SPRAY
Controls Scale Insects
Mealybugs White Flies
Red Spiders Mites and
Certain Other Garden Pests

ACCEPTED
FEB 11 1971
UNITED STATES FEDERAL INSECTICIDE
ACT AND FIFTEEN-SECOND ACT
REGISTERED PRODUCT LISTED
239-16

Active Ingredient	By Wt.
Petroleum Oil	97%
Inert Ingredients	3%
(Pet. Oil-Min. U.R. 91% - Class. Lt.-Med.)	

CAUTION: Keep out of reach of children.
 See back panel for additional cautions.

NET CONTENTS 1 GALLON LIQUID

Best Available COPY

2 of 2



ORTHO



ORTHO VOLCK® Oil Spray

VOLCK®
Oil
Spray

RECOMMENDED
USE SPRAY
INSECTICIDE

For Use Against
Scale Insects
Red Spiders
Mealybugs
White Flies
and Certain Other Pests

ALSO
MAKES AN EXCELLENT
LEAF POLISH

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

Shake well before using. Use a forceful spray, wetting pests and all parts of plant thoroughly. Do not store diluted spray. Make new dilution for each use.

FOR DORMANT USE—To Control Scale Insects, Red Spiders, Mites, and Mealybugs on Dormant Shrubs, Evergreens (note caution), Woody Plants, Fruit Trees and Shade Trees: Use 1 part VOLCK Oil Spray in 25 parts of water (½ pint (5½ ozs.) per gal. water).

GROWING SEASON—To Control Scale Insects, Red Spiders, Mites, Mealybugs and White Fly Larvae on Citrus, Fruit Trees, Shade Trees, Evergreens, Camellias, Azaleas, Hardy Shrubs: Use 1 part VOLCK Oil Spray in 50 parts water (5 Tablespoonfuls (2½ ozs.) per gal. water). This application is also effective in the removal of sooty mold fungus through the control of white fly larvae.

AS A STICKER-SPREADER—For ISOTOX Insect Spray, Lead Arsenate and Certain Other Insecticides and Fungicides used at regular strength: Use 1 part VOLCK Oil Spray in 200 parts water (4 teaspoonfuls (½ oz.) per gal.) to increase efficiency of spray treatment, provide better contact with insects or plant surfaces and to retard weathering off of the residual materials applied.

COMBINATION WITH ORTHO Malathion 50 Insect Spray on Citrus, Avocados, Mangoes, Gardenias and Ixora for Control of Scale Insects, Red Spider Mites, Mealybugs, White Flies, Aphids, Thrips and for the removal of Sooty Mold Fungus through control of White Fly Larvae—Use 1 Tablespoonful VOLCK Oil Spray and 1 Tablespoonful ORTHO Malathion 50 Insect Spray for each gallon of diluted spray. May be applied on Avocados up to within 7 days of harvest; to Grapefruit, Lemons, Limes and Oranges up to within 7 days of harvest but not during full bloom; and to Mangoes up to within 2 days of harvest.

COMBINATION WINTER SPRAY

FRUIT TREES, ROSES AND ORNAMENTALS (Dormant, deciduous only)—for Control of San Jose Scale, Rose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale and Overwintering Insect Eggs: FOR BEST RESULTS: Use 5 Tablespoonfuls VOLCK Oil Spray and 5 Tablespoonfuls ORTHORIX Spray in 1 gallon water. Keep agitated during spraying. Note cautions on ORTHORIX Spray label before use.

GREENHOUSES: To control Scale Insects, Red Spiders, Mites, White Fly, Mealybugs on Hardy Shrubs, Gardenias, Camellias and Azaleas—2 Tablespoonfuls (1 oz.) per gal. water. Allow at least 60 days between oil sprays and limit to 3 applications per year.

PALM TREES: Scale Insects, Mealybugs, Red Spider Mites—Use 1 part VOLCK Oil Spray in 100 parts water (3 Tablespoonfuls (1½ ozs.) per gal.). Note Caution.

AZALEAS, CAMELLIAS: Scale Insects, Red Spider Mites, Mealybugs, White Fly Larvae, Lace Bug Larvae—Use 1 part in 50 parts water (5 Tablespoonfuls (2½ ozs.) per gal.). Apply in spring after blooming and in fall prior to blooming.

DILUTION TABLE

	tsp. = teaspoonful		Tbs = Tablespoonful	
Add Water to make	1 to 25 (4%)	1 to 50 (2%)	1 to 100 (1%)	1 to 200 (½%)
One Gal.	½ Pt.	5 Tbs.	3 Tbs.	4 tsp.
Three Gals.	1 Pt.	½ Pt.	¼ Pt.	4 Tbs.

IF SPRAY-ETTE Garden Hose Sprayer is used—Follow directions on the bottle.

CAUTION: Harmful if swallowed. Wash hands after use. Do not use VOLCK Oil Spray on foliage within 30 days before or after using sulfur. Avoid spraying when temperature is excessively high or when plants are suffering from drought. Oil sprays should not be applied to broad-leaved evergreens or palms when freezing or near-freezing temperatures may be expected. Do not use on Maidenhair Ferns, Cocos Palms, Maples, etc. or non-woody plants unless specified. VOLCK Oil Spray may cause discoloration to Blue Spruce. Do not use in combination with or closely following treatment with ORTHOCIDE Garden Fungicide.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Chevron Chemical Company

Ortho Division, San Francisco, California 94120

Lafayette, Calif., Des Plaines, Ill.

Perth Amboy, N.J., Atlanta, Ga.

Form 4327-M Product 710

USDA Reg. No. 239-16 Made in U.S.A.