

ZINC COPOSIL® 35 Dust

ACCEPTED
DEC 12 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT GIDE ACT
FOR ECONOMIC POISON, REG. STER-
ED UNDER NO. 239-129 SUBJECT
TO ATTACHED COMMENTS.

ORTHO

(FUNGICIDE)

Active Ingredient	By Wt.
Copper (in basic cupric zinc sulfate complex) expressed as elemental	6.3%
Inert Ingredients	93.7%
Contains 6.3% zinc as elemental derived from basic cupric zinc sulfate complex.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE

This product is formulated for application by ground or air equipment.

CAULIFLOWER, CABBAGE BRUSSELS SPROUTS, BROCCOLI: Downy Mildew—25 to 50 lbs. per acre. Apply at first sign of disease and repeat at 7 to 10-day intervals to maintain control.

CELERY: Early and Late Blights—25 to 50 lbs. per acre. Apply at first sign of disease and repeat at 10 to 14-day intervals to maintain control.

CUCUMBERS, SUMMER & WINTER SQUASH, PUMPKINS, MELONS: Powdery and Downy Mildew, Leaf Spot—30 to 35 lbs. per acre. Apply when diseases appear. Repeat at 7 to 10-day intervals.

LETTUCE: Downy Mildew—36 lbs. per acre. Apply prior to or at first sign of disease. Repeat at regular

5 to 10-day intervals to maintain control.

ONIONS: Downy Mildew—35 to 50 lbs. per acre. Apply when blight appears. Repeat at 7 to 10-day intervals.

PEARS: Fire Blight (Pear Blight)—15 to 30 lbs. per acre. Begin dusting at 10% bloom and repeat at 5 to 7-day intervals during bloom. Especially important just prior to rainy periods and when rat-tail blossoms are present. Scab—30 lbs. per acre. Apply in cluster-bud, bloom and petal-fall or at other times prior to rain.

POTATOES: Late Blight—Use 25 to 50 lbs. per acre depending on plant size. Dust thoroughly and repeat the applications at 7-day intervals, as necessary. The first application should be made before the disease is expected to appear.

SPINACH: Downy Mildew—40 to 50 lbs. per acre. Apply prior to or at first sign of disease. Repeat at

regular 5 to 10-day intervals to maintain control.

STRAWBERRIES: Leaf Spot—35 to 50 lbs. per acre, apply when disease first appears and repeat as necessary.

SUGAR BEETS: Cercospora Leaf Spot—40 to 50 lbs. per acre. Apply at first sign of disease and repeat as necessary.

TOMATOES: Late Blight—Use 25 to 35 lbs. per acre depending on plant size. Dust thoroughly before disease appears, usually starting before early fall rains are expected. Repeat dust applications according to local practice.

WALNUTS: Walnut Blight—50 lbs. per acre. Apply first in late pre-bloom and second treatment in post-bloom.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

NET WEIGHT 50 LBS.

Chevron Chemical Company
Ortho Division/San Francisco, California 94119
Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
Form 861-J2 Product 918 Made in U.S.A.
EPA Reg. No. 239-129-AA