


BHC 10 Wettable



239-224



ORTHO

(INSECTICIDE)

Active Ingredients	By Wt.
Gamma Isomer of Benzene Hexachloride	10%
Other Isomers of Benzene Hexachloride	10%
Inert Ingredients	80%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

ORTHO BHC 10 Wettable mixes readily with water for spray application. It is also suitable for blending with the proper carrier for manufacture of dusts. This product can be mixed with Summer Oil Emulsions, ORTHO Spreaders, FLOTOX Wettable Sulfur, Insoluble Copper fungicides, Bordeaux Mixture, Lime Sprays, Lead Arsenate, Calcium Arsenate Sprays, Nicotine, Derris, Rotenone, VAPOTONE, Toxaphene, and ORTHOCIDE. ORTHO BHC 10 Wettable is not compatible with liquid Lime Sulfur, Lime Sulfur.

DIRECTIONS

Repeat application as necessary.

USE INDICATED AMOUNT IN 100 GALLONS WATER UNLESS OTHERWISE SPECIFIED.

APPLES, PEARS: Aphids (including Woolly Apple Aphid), Plum Curculio, Tent Caterpillars, Pear Psyllid, Tarnished Plant Bug—2 to 3 pounds. Do not apply within 60 days of harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 2643 B3 Product 819 Made in U.S.A.

EPA Reg. No. 239-224-AA

PEACHES, PLUMS, PRUNES: First brood only of Plum Curculio, Stink Bugs, Aphids, Thrips, Tarnished Plant Bug—2 to 3 pounds. Do not apply within 60 days of harvest.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, KALE, COLLARDS: Aphids, Thrips—2 to 3 pounds per acre in water to cover. Do not apply to Cabbage after thinning. Do not apply to Cauliflower, Broccoli, Brussels Sprouts, Kale and Collards after the seeding stage.


CELERY (seedling only): Aphids, Thrips, Certain Caterpillars—2 to 3 pounds per acre in water to cover.

GLADIOLUS: Thrips—2 pounds.

GRAPES: Bud Beetles (Grape Hoplia)—4 lbs. per acre in 200 to 250 gals. water. Do not apply after edible parts start to form.

GRASSHOPPERS, CRICKETS and EARWIGS: Fence Rows, Barrier Zones, Wastelands, Uncultivated Areas—As a dust: Apply 4 to 8 pounds per acre directly to infested area. Where grasshoppers are migrating into field put a 10-foot barrier strip of dust around field or on side from which they are approaching. As a spray: Apply 2 pounds per 100 gallons water. As a bait: Mix 4 pounds with each 100 pounds regular formula that in place of Arsenic or Sodium Fluosilicate.

PEPPERS: Aphids, Leaf Miners—3 pounds per acre in water to cover. Do not apply after the edible parts start to form.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION:

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Do not store in a cool, dark place near feed or food. Do not use on foliage or fruit of Apples, Potatoes, Beans, Lima Beans, Peas, or Cucumbers. Do not use on ornamentals. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Toxic to fish and wildlife. Do not apply where any body of water will be contaminated by drift, application or runoff. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

SOIL INSECTS: Wireworms, Cutworms, White Grubs, Diabrotica larvae and certain other Soil Insects on Celery, Cucumbers, Kale, Lettuce, Melons, Pumpkins, Spinach—Pre-Plant Treatment—Apply 5 lbs. in sufficient water to cover 1 acre. Apply as spray or dust directly on soil, then disc as soon as practicable to give maximum penetration in the upper 4 to 6 inches of soil. ORTHO BHC 10 Wettable can be mixed with acid or neutral fertilizers and applied with fertilizer drill. Do not use on root crops or in soil in which root crops are to be planted.

STRAWBERRIES: Spittle Bugs, Strawberry Weevils—1 pound per acre in water to cover. Do not apply after the edible parts start to form.

TOMATOES: Colorado Potato Beetles, Flea Beetles, Thrips—2 lbs. per acre in water to cover. Aphids, Leaf Miners—2 to 3 lbs. per acre in water to cover. Do not apply after the edible parts start to form.

ORNAMENTAL TREES: PINE BARK BEETLES, BORERS, WOOD BORING BEETLES—15 pounds per 45 gals. water and wet trunks to runoff.

TICKS AND CHIGGERS ON VEGETATION: As a spray: 1 pound. As a dust: Dust collected at 16 to 20 pounds per acre. Do not use on areas to be grazed by dairy or meat animals.

BULB TREATMENT: WIREWORMS, THIRPS: Dust dry bulbs with 5 to 8 ounces to each 100 pounds of bulbs at time of planting. Do not dust and let stand before planting. ALTERNATE TREATMENT: Dust open trench before planting bulbs at rate of 5 lbs. per acre.

BULB TREATMENT: SYMPHYLIDS—Use 20 pounds per acre worked into the top 4 inches of soil.

SARCOPTIC MANGE, FLEAS, LICE, TICKS, GNATS, MOSQUITOES on Dogs: Use 1 lb. per 3 gallons water as a spray or 1 lb. per 100 lbs. of material per 100 lbs. of material. Repeat after 10 days if necessary. Do not use on cats and other mammals. Do not use indoors.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that the material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning the material.

2. Certain and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from handling, storage, and use of this material.