


# ISOTOX<sup>®</sup> Lindane Wettable 25



# ORTHO



## (INSECTICIDE)

Active Ingredient	By Wt.
Lindane—	
Gamma Isomer of Benzene Hexachloride	25%
Inert Ingredients	75%

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE OR FOR DRUMS, WASH AND DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

**APPLES:** Aphids—1 lb. in 100 gals. water (300 to 400 gals. spray per acre). Concentrate Sprayer—3 to 4 lbs. per acre in water to cover. Treat under leaves also. Do not apply within 60 days of harvest.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KALE, COLLARDS:** Aphids, Flea Beetle, Caterpillars, Thrips, Loopers—1/2 lb. in 100 gals. water (200 gals. spray per acre). Do not apply after the edible parts start to form. Cabbage Maggots—4 lbs. per acre in water to cover. Apply to soil around base of plants.

**CELERY:** Aphids, Thrips, Tarnished Plant Bugs, Certain types of Lepidopterous Larvae, Climbing Cutworms—1/2 lb. in 100 gals. water (200 gals. diluted spray per acre). Do not apply after the celery is half-grown or after it begins to bunch.

**CUCUMBERS, MELONS, SQUASH, SUMMER SQUASH, PUMPKINS:** Cucumber Beetles, Thrips—1/2 lb. in 100 gals. water (200 gals. spray per acre). Treat under leaves also. Squash Bugs, Leaf Miners, Grasshoppers, Lygus Bugs, Aphids, Melon Worms, Pickle Worms—1 lb. in 100 gals. water (200 gals. spray per acre). Late treatment of melons and squash may result in reduction of quality at harvest. Do not apply to Cucumbers, Squash and Summer Squash within 1 day of harvest. Remove excess residues.

**GLADIOLUS AND DAFFODIL BULBS:** Wireworms—Dust dry bulbs at time of planting with 2 to 3 ounces to each 100 lbs. bulbs for temporary protection while rooting.

## NET WEIGHT 20 POUNDS

**Chevron Chemical Company**  
 Ortho Division/San Francisco CA 94119 Richmond CA 94804  
 Product 958 Made in U.S.A.  
 Form Gen 9012-B  
 EPA Reg. No. 239-229 ZA

**GRASSHOPPERS:** Use on listed crops (Except Strawberries) at recommended dosages.

**ONIONS:** Thrips—1 lb. per 100 gals. water (200 gals. spray per acre). First application when thrips average 5 to 6 per plant. Repeat in 5 to 6 days. A third application may be necessary. Do not use on green or spring onions. Onion Maggot—4 lbs. per acre in water to cover. Apply to soil around base of plants when plants are 2 to 3 inches tall. Repeat at 7 to 10-day intervals.

**PEPPERS:** Aphids, Leaf Miners—1/2 lb. in 100 gals. water (200 gals. spray per acre). Armyworms—2.4 lbs. per acre in water to cover. Do not apply after the edible parts start to form.



**DO NOT TAKE INTERNALLY  
 DO NOT INHALE  
 DO NOT GET ON SKIN**

**WARNING:** May Be Fatal If Swallowed! Do not breathe dust or spray mist. May Be Absorbed Through Skin. Avoid contact with skin and eyes. Wash thoroughly after handling.  
**Note to Physicians:** Emergency Information — call (415) 233-3737. Do not store or transport near feed or food. Do not apply to edible crops within 30 days of harvest unless otherwise noted. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply to any area not specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**SOIL INSECTS:** Wireworms, Cutworms, White Grubs, Diabrotica Larvae and certain other soil insects on Celery, Cucumbers, Cabbage, Broccoli, Brussel Sprouts, Cauliflower, Strawberries, Kale, Lettuce, Melons, Pumpkins, Spinach, Tomatoes, Mint (Post harvest in fall prior to winter period)—Pre-Plant Treatment—1 to 2 lbs. per acre. Apply as spray or dust directly on soil. Disc as soon as practicable to give thorough distribution in upper 4 to 6 inches of soil. Can be mixed with acid or neutral fertilizers and applied with fertilizer drill. Do not apply to root crops (except potatoes) within 1 year. Do not use in soil to be planted to potatoes within 2 years.

**SPINACH:** Aphids, Leaf Miners—3/4 lb. in 100 gals. water (200 gals. spray per acre). Use only in seedling stage.

**STORED PRODUCTS INSECTS:** Granary Weevil, Sawtoothed Grain Beetles, Confused Flour Beetle, Corn or Rice Weevil, Bean Weevil, Angoumois Grain Moth, Indian Meal Moth (Grain Bins, Storage Bins, Warehouses)—16 1/2 lbs. per 100 gals. water. Apply as a spot treatment. Spray into breeding places, areas where insects

congregate or onto insects directly. Repeat as necessary. Controls only accessible stages of these pests. Apply only so that foodstuffs or grain will not be contaminated.

**STRAWBERRIES:** Strawberry Weevil, Spittlebug—1/3 lb. in 100 gals. water (200 gals. spray per acre). Concentrate Sprayer—2/3 lb. per acre in water to cover. Apply in spring before blossoms open when insects appear. Thoroughly treat flower buds. Do not apply after the fruit starts to form.

**TOMATOES (for fresh fruit market only):** Aphids, Thrips—1/2 lb. per 100 gals. water (200 gals. spray per acre). Concentrate Sprayer—1 lb. per acre in water to cover. Leaf Miners—3/4 lb. in 100 gals. water (200 gals. spray per acre). Concentrate Sprayer—1 1/2 lbs. per acre in water to cover. Do not apply after fruit starts to form.

### USES ON ANIMALS

Do not treat animals under 3 months of age. Do not spray beef cattle, sheep, goats and hogs within 30 days of slaughter. Do not dip them within 60 days of slaughter. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Do not use on dairy cattle or dairy goats. Avoid treating animals in cold, stormy weather. Do not dip or spray thirsty, overheated or sick animals. Using dosages given below, spray entire animal or infested areas as necessary in following amounts. **Adult Animals (Horses, Cattle):** 2 qts. diluted spray per head, using proportionately lesser amounts for smaller animals. Use at from 300 to 400 lbs. pressure. **Young Animals (those over 3 months of age):** 1 qt. diluted spray per animal. Wash hands thoroughly with soap and warm water after handling the concentrate or applying the diluted product to animals.

**BEEF CATTLE, SHEEP, GOATS, HOGS, HORSES:** Spray or Dip—Psoroptic and Sarcoptic Mange (Scab or Scabies) — 1 1/2 lbs. per 100 gals. water. For most effective treatment, apply 2 sprays one week apart. Do not use on sick animals. **Lice, Ticks, Horn Flies, Stable Flies**—1 lb. per 100 gals. water. **Sheep Ticks (Keds) on Sheep** — 1 1/2 lbs. in 100 gals. water. Spray along back and around ears, spraying against wool. Spray in fall. In some areas a repeat treatment after shearing is also advisable. About 350 lbs. pressure is desirable.

Also for use in accordance with the recommendations and instructions issued by the United States Department of Agriculture for quarantine programs and to be used only by or under the direction of federal/state personnel for quarantine treatments.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from handling, storage and use of this material.