

1/2 H 221-271
1/11/85
1.1/2


ISOTOX Lindane Wettable 25
EPA Reg. No. 239-229
Submitted: September 13, 1985

COPY A

If swallowed, give water or milk to drink and telephone for medical advice. Consult medical personnel before inducing vomiting. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment center or hospital. In case of eye contact, flush eyes with fresh water for at least 15 minutes. See a doctor if irritation persists. If on skin, wash with plenty of soap and water. If inhaled, move person to fresh air.

CONFIDENTIAL

PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA



ACCEPTED

JAN 22 1986

Un...
 For...
 as...
 registered under
 EPA Reg. No. 239-229

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING
 May be fatal if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Do not breathe dust or spray mist. Wash thoroughly after handling. Laundry can turn soiled clothing.

STATEMENT OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS
 This product is toxic to fish and wildlife. Do not apply directly to water.
 This pesticide is highly toxic to bees exposed to direct treatment or residues remaining on the treated areas. Do not apply when the bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Ortho

ISOTOX® Lindane
Wetttable 25
(INSECTICIDE)

Active Ingredient
 Lindane—25%
 Gamma Isomer of Benzene Hexachloride—25%
 Inert Ingredients—50%

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

BEETERY STATEMENT
 Keep all unprotected persons out of operating areas or vicinity where there may be drift.
 Do not enter treated areas without protective clothing until sprays have dried.
 Certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information. Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

Actual application of Lindane is prohibited.
 Applicators must wear the following protective clothing during the application process: a light weight protective suit or coveralls, water resistant hat, unlined, waterproof gloves, and unlined, light weight socks. Masks and bandanas must also wear goggles or a face shield, waterproof gloves and a waterproof apron.
 Do not apply to edible food crops within 30 days of harvest unless otherwise noted.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KALE, COLEAROS. Aphids, Flea Beetles, Caterpillars, Thrips, Leaf Miners—1 lb. in 100 gals. water (200 gals. spray per acre). Do not apply until the aphids have started to form. Cabbage Maggot—4 lbs. per acre in water to cover. Apply to soil around base of plants.

CITRUS. Aphids, Thrips, Leafminers, Flea Beetles, Caterpillars, Thrips, Leaf Miners—1 lb. in 100 gals. water (200 gals. spray per acre). Do not apply after the citrus is half grown or when the fruit begins to bunch.

KEEP OUT OF REACH OF CHILDREN
WARNING-AVISO
FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.
PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

NET WEIGHT

Chevron Chemical Company © 1984
 Ortho Agricultural Chemicals Division
 San Francisco CA 94120-7141 Richmond CA 94604-0036
 Product 958 Made in U.S.A. EPA Est. 239-CA-1
 Tel. (415) 741-1111
 EPA Reg. No. 239-22

CUCUMBERS, MELONS, SQUASH, SUMMER SQUASH, PUMPKINS: Cucumber Beetles, Thrips—1 lb. in 100 gals. water (200 gals. spray per acre). Treat with 1 lb. of Squash Bugs, Leaf Miners, Grasshoppers, Leafhoppers, Aphids, Flea Beetles, Pickle Weevils—1 lb. in 100 gals. water (200 gals. spray per acre). Late in season of melons and squash may result in reduction of quality of harvest. Do not apply to Cucumbers, Squash and Summer Squash within 1 day of harvest. Pumpkin seeds residues.

GRASSHOPPERS: Use on listed crops (except Strawberries) at recommended dosages.
ONIONS: Thrips—1 lb. per 100 gals. water (200 gals. spray per acre). First application when plants average 3 to 6 per plant. Repeat in 5 to 6 days. A third application may be necessary. Do not use on green or spring onions. Onion Maggot—4 lbs. per acre in water to cover. Do not apply to plants when plants are 2 to 3 inches tall except at 1 lb. per 100 gals. water.

PEPPERS: Aphids, Leaf Miners—1 lb. in 100 gals. water (200 gals. spray per acre). Armyworms—2 1/2 lbs. per acre in water to cover. Do not apply after the seedlings start to form.

SOIL INSECTS: Wireworms, Cutworms, White Grubs, Diabrotica larvae and certain other soil insects on Celery, Cucumbers, Cabbages, Broccoli, Brussels Sprouts, Cauliflower, Strawberries, Kale, Lettuce, Melons, Pumpkins, Spinach, Tomatoes, Mint (Post harvest in fall prior to winter storage)—Per Flea Treatment—1 to 2 lbs. per acre. Apply as spray or dust directly on soil. Disc as soon as practicable to give thorough distribution in upper 4 to 6 inches of soil. Can be mixed with acid or neutral fertilizers and applied with fertilizer drill. Do not apply to root crops (except potatoes) within 1 year. Do not use in soil to be planted to potatoes within 2 years.

SPINACH: Aphids, Leaf Miners—1 lb. in 100 gals. water (200 gals. spray per acre). Use only in seedling stage.

STORED PRODUCTS INSECTS: Granary Weevil, Sawtoothed Grain Beetles, Confused Flour Beetle, Corn or Rice Weevil, Bean Weevil, Angoumois Grain Moth, Indian Meal Moth (Grain Bins, Storage Bins, Warehouses)—16 1/2 lbs. per 100 gals. water. Apply as a spot treatment. Spray into feeding places, areas where insects congregate or collect directly. Repeat as necessary. Control or if excessive stages of these pests. Apply only so that foodstuffs or grain will not be contaminated.

STRAWBERRIES: Strawberry Weevil, Spittlebug—1 lb. in 100 gals. water (200 gals. spray per acre). Concentrate Sprayer—1/2 lb. per acre in water to cover. Apply in spring before blossoms open when insects appear. Thoroughly treat flower buds. Do not apply after the fruit starts to form.

TOMATOES: (For fresh fruit market only): Aphids, Thrips—1 lb. per 100 gals. water (200 gals. spray per acre). Concentrate Sprayer—1 lb. per acre in water to cover. Leaf Miners—1 lb. in 100 gals. water (200 gals. spray per acre). Concentrate Sprayer—1 1/2 lbs. per acre in water to cover. Do not apply after the fruit starts to form.

DIP USES ON ANIMALS
 Do not treat animals under 3 months of age. Do not dip beef cattle, sheep, goats and hogs within 60 days of slaughter. Do not treat swine within 2 weeks of farrowing nor for at least 3 weeks thereafter. Do not use on dairy cattle or dairy goats. Avoid treating animals in cold, stormy weather. Do not dip in dirty, overcasted or sick animals. Wash hands thoroughly with soap and warm water after handling the concentrate or applying the diluted product to animals.

DEER, CATTLE, SHEEP, GOATS, HOGS, HORSES: Dip Permethrin and Sarcopile Mixture (Scale or Scabiosis)—1 1/2 lbs. per 100 gals. water. Do not use on sick animals. Lice, Ticks, Horn Flies, Stable Flies—1 lb. per 100 gals. water. Sheep Ticks (Ixodes)—on Sheep—1 1/2 lbs. in 100 gals.

Also for use in accordance with instructions issued by the United States for quarantine programs and to direction of federal/state personnel.

Note: Used dip solutions must be collected in the Resource Conservation and Recycling Act. Do not use more than 1000 lb. used or other hazardous waste, the material must be subject to subpart C of treatment, storage or disposal of hazardous waste as a hazardous waste for.

Do not apply under conditions in large or other plantings that may be rendered unusable for sale, use.

STORAGE AND PROHIBITIONS
 Do not contaminate water, food or clothing with equipment.
 Open dumping is prohibited.

STORAGE
 Keep pesticide in original container. Dilute into food or drink container near feed or food.
 For help with any spill, leak, fire, material, call day or night (415) 741-1111.

PESTICIDE DISPOSAL
 This product is toxic. Improper spray mixture, or misuse is a threat. Do not dispose of this material in any way. For disposal instructions, contact your State Pesticide Agency or the Hazardous Waste nearest EPA Regional Office for **CONTAINER DISPOSAL**. Completely empty bag into appropriate container. Dispose of empty container, or if allowed by burning. If burned, stay out of.

CONDITIONS OF SALE: 1. Chevron warrants that this material conforms to the label and is reasonably free from adulteration. Chevron neither makes, nor authorizes, any other warranty of fitness, guarantee or representation, express or implied.
 2. Critical and unforeseeable to prevent it from eliminating all of chemicals. Such risks include, but are not limited to, the use of plants and crops to which the material is applied, and damage caused by such risks even though the user stated hereon and followed Bayer and user action liability (except those assumed by Bayer from handling, storage, or use).

RQ LINDANE, MIXTURE
ORM-A