


Parathion 2 Dust



ORTHO



(INSECTICIDE)

Active Ingredient	By Wt.
Parathion*	2%
Inert Ingredients	98%

*O,O-diethyl O-p-nitrophenyl thiophosphate.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

COMPLETELY EMPTY BAG AND DISPOSE OF WASTE PESTICIDES AND BAG BY BURYING. DO NOT BURN. NEVER RE-USE.

This product is formulated for application by ground or air equipment. This product is not to be used around homes or home gardens.

BEANS: Mexican Bean Beetle, Red Spider, Two-Spotted Mites, Aphids, Leaf Miners, Leaf Hoppers, Lygus Bugs, Looper Caterpillars, Climbing Cutworms, Armyworms—25 to 40 lbs. per acre. Do not apply within 15 days of harvest.

BLUEBERRIES: Blossom Weevils—15 to 25 lbs. per acre. Make full-coverage applications when leaf buds show green and 7 to 9 days later. Do not apply within 14 days of harvest.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS: Aphids, Cucumber Beetle, Darkling Ground Beetle, Spider Mites, Leafhoppers—25 pounds per acre. Do not apply to Cabbage within 10 days of harvest and to Cauliflower, Broccoli and Brussels Sprouts within 7 days of harvest.

CARROTS: Aphids—20 lbs. per acre. Do not use treated carrot tops for food or feed. Do not apply within 15 days of harvest.

CELERY, CUCUMBERS, MELONS: Aphids, Leafminer, Thrips, Darkling Ground Beetle, Spider Mites—20 lbs. per acre. Do not apply to Cucumbers before vining or within 15 days of harvest; to Celery within 21 days; and to Melons within 7 days of harvest.

CHERRIES: Cherry Slug, Tortrix, Leaf Roller, Case Bearers, Black Aphid, Tussock Moth Caterpillar—30 to 40 pounds per acre. Do not apply within 14 days of harvest.

CORN: Aphids—25 lbs. per acre. Do not apply within 10 days of picking, cutting or forage.

COTTON: Cotton Aphids, Red Spiders, Leafhoppers, Spittlebugs—20 pounds per acre. Do not apply more than 2 or 3 times during the season. Louisiana—10 pounds per acre. Workers entering fields within 24 hours after application should wear protective clothing. Do not hand pick or harvest within 7 days.

FILBERTS: Filbert Aphid, Apple Mealybug, Eye-spotted Bud Moth—20 lbs. per acre. Consult State Agricultural Experiment Stations for proper timing. Do not apply after hulls begin to open.



POISON



ANTIDOTE: If Swallowed, Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY!** Note to Physicians: Atropine is Antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. **If On Skin:** In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

DANGER:

KEEP OUT OF REACH OF CHILDREN

DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

DANGER:

Poisonous If Swallowed, Inhaled or Absorbed Through Skin! Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for parathion protection. **Rapidly Absorbed Through Skin!** Do not get in eyes, on skin or in clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of treating areas or vicinity where there may be danger of drift. Avoided areas should not be reentered until drying insecticides are visible on treated area. Do not store or transport near food or feed. Do not apply under conditions involving possible drift to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Do not use in or near water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff may occur. Do not apply to areas not listed on this label. This product is highly toxic to bees exposed to direct treatment or drifting on nearby forage. Protective information may be obtained from your cooperative Agricultural Extension Service.

GRAPES: Mealybug, Leaf Folder, Red Spider Mites—30 to 35 lbs. per acre. Do not apply within 14 days of harvest.

LETTUCE: Aphids, Leaf Miner—25 lbs. per acre. Reduction of Mosaic may be accomplished by aphids control. Do not apply to Head Lettuce within 7 days of harvest and to Leaf Lettuce within 21 days of harvest.

ONIONS: Thrips—Use 20 pounds per acre. Do not apply within 15 days of harvest.

PEAS: Pea Aphid, Pea Weevil—Use 20 pounds per acre. Do not apply within 10 days of harvest.

PEPPERS: Aphids, Serpentine Leaf Miner, Loopers, Climbing Cutworms, Spider Mites, Armyworms—20 to 40 lbs. per acre. Do not apply within 15 days of harvest.

PLUMS, PRUNES: Thrips, Red Spider Mites—50 to 60 lbs. per acre at Jacket Stage. Do not apply within 14 days of harvest.

POTATOES: Aphids, Colorado Potato Beetle, Flea Beetle, Leafhopper, Serpentine Leafminer, Grasshoppers, Whiteflies—40 lbs. per acre. Do not apply within 5 days of harvest.

SPINACH: Leaf Miner—25 pounds per acre. Do not apply within 14 days of harvest.

STRAWBERRIES: Red Spider, Aphids, White Fly, Cyclamen Mite, Thrips, Lygus Bugs, Potato Leafhoppers—20 pounds per acre. Repeat against Red Spider in 7 to 10 days. Do not apply within 14 days of harvest.

SUGAR BEETS: Webworm—17 pounds per acre. Do not apply within 15 days of harvest.

SQUASH, SUMMER SQUASH: Aphids, Leafminer, Thrips, SUMMER SQUASH (SEED): Squash Bug—25 lbs. per acre. Do not apply within 15 days of harvest.

TOBACCO (SEEDBED): Aphids—1/2 to 3/4 lb. per 100 sq. yds. (25 to 35 lbs. per acre). Not more than 2 applications. Handlers of treated plants should wear gloves. Do not apply to plant beds with shaker cans or cloth bags.

TOMATOES: Aphids, Spider Mites, Hornworm—20 to 30 lbs. per acre. Do not apply within 10 days of harvest.

WALNUTS: Walnut Aphids—Use 30 to 40 pounds per acre. Do not apply on walnuts after the outer husks open.

WHEAT, OATS, BARLEY: Green Bug (Aphid)—20 pounds per acre. Do not apply within 15 days before harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that the material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Chevron neither makes nor authorizes any agent or representative to make any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemical. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above).

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 2113-73 Product 1153 Made in U.S.A.

EPA Reg. No. 229-305-AA