



# ORTHO

## Chlordane 40 Wettable

**(INSECTICIDE)**

Active Ingredient  
Technical Chlordane\*

Inert Ingredients ..... 60%

\*Equivalent to 24% Octachloro-4, 7-Methanotetrahydroindane and 16% related compounds.

ACCEPTED  
FEB 2, 1973  
UNDER FEDERAL INSECTICIDE  
FUNGICIDE AND ROENTGENIC ACT  
FOR ECONOMIC TOXIC SUBSTANCE  
REGISTRATION  
ED UNDER 40 CFR 155.303



### READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Harmful if swallowed. Avoid breathing dust or spray mist. Contact with skin causes toxic symptoms. In case of contact with skin, wash with soap and water. Do not store or transport near feed or food. Keep children and pets off treated areas until material is watered into the soil and grass is dry. Forage crops treated with this material should not be fed to dairy animals or to animals being finished for slaughter. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product will kill fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

**DIRECTIONS ON BACK**

## NET WEIGHT 4 POUNDS

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

**Chevron Chemical Company**

Ortho Division/San Francisco, Calif. 94119  
Richmond, California Fresno, California Honolulu, Hawaii  
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida  
Form 3496-J Product 1176 Made in U.S.A.

EPA Reg. No. 239-323-AA

## ORTHO® Chlordane 40 Wettable

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

### DIRECTIONS

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

Repeat treatment as necessary. Do not apply to food crops within 30 days of harvest unless otherwise specified. Remove residues remaining at harvest by washing, wiping or other effective means.

**ANTS:** In lawns and golf courses, apply 1½ ounces in sufficient water to cover 1,000 sq. ft. It is essential that the turf be thoroughly soaked with water and rolled after treatment. Birds feeding on treated areas may be killed.

**GLADIOLI:** Thrips—Use 2½ pounds per 100 gallons diluted spray.

**MOLE CRICKETS** (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Collards, Tomatoes, Ornamentals): In Seed Beds—9 ozs. per 100 gals. water on 1,000 sq. ft. of bed. In the Field: 2½ lbs. per 100 to 125 gals. water per acre. ORTHO Chlordane 40 Wettable may be mixed in non-alkaline mineral fertilizer which is spread broadcast and disked into topsoil. The amount combined will depend on rate of fertilizer application. Use sufficient ORTHO Chlordane 40 Wettable so that 3 to 4 pounds will be distributed per acre.

**ONIONS** (Make pre-planting soil applications at time of planting or transplanting only): Onion Maggots—Use 5 pounds per acre. Do not use on green or spring onions.

**SOIL INSECTS:** Ants, the larvae of White-fringed and Japanese Beetles on Beans, Corn, Cucumbers, Onions, Sweet Potatoes and Tomatoes—Use 12½ to 15 pounds per acre. FOR BEST RESULTS: Apply either as a dust or spray. Work into the top few inches of soil before planting.

**VEGETABLES** (BEETS: Climbing Cutworms, Surface Cutworms, Flea Beetles, Armyworms. BEANS: Flea Beetles, Green Stink Bug; CARROTS: Carrot Rust Fly. BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KALE, COLLARDS: Imported Cabbageworm, Diamondback Moth Larvae, Cross-striped Cabbageworm, Cabbage Looper, Armyworm, Green Stink Bug, Harlequin Bug. CUCUMBERS: Squash Bug, Squash Vine Borer, Cucumber Beetles. ONIONS: Thrips. TOMATOES: Tomato Hornworm, Armyworm, Serpentine Leaf Miner, Flea Beetles, Climbing Cutworms)—2½ lbs. per 100 gals. water at 10 to 14 day intervals. Beets, Beans, Carrots, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Collards, Onions and Tomatoes—Apply up to 200 gals. per acre. Cucumbers—Apply up to 150 gals. per acre. Do not apply to Beets after the seedling stage if tops are to be used for food. Do not apply to green or snap beans after pods start to form. Do not apply to Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Collards, Cucumbers or Tomatoes after the edible parts start to form. Do not use on green or spring onions.

FORM 3496-J