


ISOTOX Seed Treater (75)
EPA Reg. No. 239-353
Submitted: June 14, 1985

COPY A

If swallowed, DO NOT make person vomit. If person is conscious, give a large quantity of water or milk to drink and call a doctor or emergency hospital. In case of eye contact, flush eyes with fresh water for at least 15 minutes. See a doctor if irritation persists. If on skin, wash with plenty of soap and water. If inhaled, move person to fresh air.



CONFIDENTIAL

PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS WARNING

May be fatal if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Do not breathe dust or spray mist. Wash thoroughly after handling. Launder contaminated clothing. This product should be applied in a well ventilated area. Applicators who apply this product manually or without the use of a closed-system treatment procedure must wear the following protective clothing during the application process: long-sleeved shirt, long pants, gloves, and a disposable, paper dust mask which covers at least one-third of the face.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, give a large amount of water to drink, make person vomit, and call a doctor. In case of eye contact, wash eyes immediately with fresh water for at least 15 minutes. See a doctor if irritation persists. For skin contact, wash skin thoroughly with soap and water.

Note to Physicians: Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on exposed seed may be killed. Do not apply directly to water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL, USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

ISOTOX Seed Treater (75) is a special formulation for use in the treatment of seeds against wireworms and seed corn maggots. Also repellent to pheasants. It is suitable for use in combination with ORTHOCIDE 75 Seed Protectant, and other generally used fungicides, and may be applied either dry or through slurry treaters, generally used for wet application of fungicides on seeds.

GRAIN AND OTHER STORAGE BINS FOR GRAIN AND STORED PRODUCTS INSECTS (exposed stages): Saw Toothed Grain Beetle, Granary Weevil, Corn-Infused Flour Beetle, Corn or Rice Weevil, Bean Weevil, Angoumois Grain Moth, Indian Meal Moth—5½ lbs. in 100 gals. water. Spray empty storage bins prior to filling. Apply 2 gals. diluted spray per 1000 sq. ft. of bin area.

(Cwt. = 100 lbs.)	SLURRY METHOD		DRY METHOD	
	Amt. Seed Treater/ Gal. Slurry	Cup Size in Cubic Centimeters	Lbs. Seed Treated/ Gal. Slurry	Amt. Seed Treater/ Measure Seed
SEED				
CORN	2 lb. 11 oz.	23	1650	2½ oz./cwt.
BEANS—(Use with a protective fungicide) Baby bush limas, Bush snap and all others	11 oz.	23	1650	¾ oz./cwt.
Standard and Fordhook large limas	6 oz.	23	1650	½ oz./cwt.
SUGAR BEETS	2 lb. 7 oz.	46	720	5½ oz./cwt.
TOMATOES	1 lb. 3 oz.	46	720	2½ oz./cwt.
PEAS	1 lb. 6 oz.	23	1630	1½ oz./cwt.
CUCUMBERS, CANTALOUPE	1 lb. 5 oz.	23	1550	1½ oz. wt.
CARROTS	8 oz.	46	620	1½ oz./cwt.
COTTON				
Acid Delinted or Reginned	1 lb. 6 oz.	46	820	2½ oz./cwt.
Fuzzy—not delinted	1 lb. 9 oz.	46	820	3 oz./cwt.
MILO, SORGHUM	13 oz.	46	780	1½ oz./cwt.
ALFALFA, CLOVER	6 oz.	46	740	¾ oz./cwt.
BARLEY	1 lb. 6 oz.	23	1650	1½ oz./cwt.
SPINACH	10 oz.	46	720	1½ oz./cwt.
OATS	9 oz.	23	1400	¾ oz./cwt.
SAFFLOWER	11 oz.	23	1560	¾ oz./cwt.
RYE	11 oz.	23	1560	¾ oz./cwt.
SUDAN GRASS	5 oz.	46	660	¾ oz./cwt.
SOYBEAN	11 oz.	23	1700	¾ oz./cwt.
WHEAT	11 oz.	23	1730	¾ oz./cwt.
WATERMELON	1 lb. 5 oz.	23	1500	1½ oz./cwt.
LETTUCE	9 oz.	46	630	1½ oz./cwt.
OKRA	1 lb. 6 oz.	23	1650	1½ oz./cwt.
SUNFLOWER	1 lb. 5 oz.	23	1550	1½ oz./cwt.



ORTHO

ISOTOX® Seed Treater (75)

(INSECTICIDE)

ALSO REPELLENT TO PHEASANTS

Active Ingredient

Lindane—

Gamma Isomer of Benzene Hexachloride 75%

Inert Ingredients 25%

By Wt.

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

Chevron Chemical Company © 1981

Ortho Agricultural Chemicals Division

San Francisco CA 94120-7144

Richmond CA 94804-0036

Product 1972

Made in U.S.A.

R Form 9251-G

EPA Reg. No. 239-353-ZB

NET WEIGHT 20 POUNDS

Directions for Use (continued)

Treated seed must not be used for processing for oil. For on the spot for immediate use, minimizing. Do not store excess treated seed. Do not store excess treated seed with this applicability provisions of the commercially treated seed must be mixed with food or animal seed may be hazardous to by burial away from bodies.

If soil temperature is extremely hot and mid-summer planting is storage of treated seed to in a slurry treater do not seed has had a chance to dry. There may be some crinkling moisture entering seed causing quality or germination via organism damage, combining.

STORAGE

PROHIBITIONS

Do not contaminate water or equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original food or drink containers. For help with any spill, le day or night (415) 233-3737.

PESTICIDE DISPOSAL

This product is acutely hazardous. Do not spray, mix, or rinse. cannot be disposed of by State Pesticide or Environmental representative at the nearest.

CONTAINER DISPOSAL

Completely empty liner. Do not loosen clinging particles. Do not reuse container. Disposal if allowed by State and cannot be reused, disposed.

CONDITIONS OF SALE: 1. that this material conforms reasonably fit for use as directed by any agent or representative of MERCHANTABILITY, guaranteeing this material.

2. Critical and unforeseeable eliminating all risks in conclusion, but are not limited material is applied, lack of other plants or crops, 5 inch fit for the uses stated here. Buyer and user acknowledge assumed by Chevron under of this material.