


ISOTOX[®] Transplanter Solution

(INSECTICIDE)

Active Ingredient	By Wt.
Lindane—	
Gamma Isomer of Benzene Hexachloride	7.5%
Inert Ingredients	92.5%
Lindane per gallon—8.5 ounces	



Comments

CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

ISOTOX Transplanter Solution is a special lindane formulation for use in transplanting water against wireworms and cabbage maggots. It mixes readily with water and stays in suspension without agitation during the transplanting operation. The use of ISOTOX Transplanter Solution in transplanting water establishes a zone of protection around the roots of newly set plants. This allows the roots to develop and the plants to become established.

DIRECTIONS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KALE, COLLARDS: Wireworms, Cutworms, Cabbage A maggot. Add 8.5 fluid ounces (1/4 pint) in 50 gallons of transplanting water, or about 1/2 pint per plant or 200 to 250 gallons per acre. Apply ISOTOX Transplanter Solution directly to the tank with water.

NET CONTENTS
1 GALLON LIQUID

Chevron Chemical Company

San Francisco, Calif. 94119
 Berkeley, California Fresno, California Honolulu, Hawaii
 Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
 EPA Reg. No. 239-531 Product 1469 Made in U.S.A.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid spray mist. May be absorbed through skin. Avoid skin contact. In case of contact, wash immediately with soap and water. Do not use on household pets or humans. Avoid contamination of feed and foodstuffs. **WARNING—FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.** Do not apply to root crops. Do not apply in soil to be planted to root crops within 1 year, except for potato transplants. Do not use on foliage of white potatoes. When container is empty, immediately wash and destroy. Never re-use.

This product will kill fish and wildlife. Keep out of any water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from area. Do not apply where runoff is likely to occur. Do not discharge into water by cleaning of equipment, or disposal of wastes.