




# ORTHO Tomato Vegetable Dust



READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS, WARNINGS AND DIRECTIONS

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS

## DIRECTIONS

ORTHO Tomato Vegetable Dust may be used as a dust or spray. For best results, dust or spray in early morning or late evening when temperatures are low and humidity is high.

**As a Spray:** Mix 3 to 4 Tablespoons in a little water to make a thin paste; add to 1 gallon water and agitate during application. Do not store diluted spray. Make new dilution for each use.

**For Insect Control:** Apply when insects first appear and repeat at 5 to 10 day intervals or as necessary.

**For Disease Control:** Start program of dusting as soon as foliage appears. Repeat at 1 to 2 week intervals for best protection.

Can be used on Beans up to within 3 days of harvest; on Raspberries within 14 days of harvest; on Potatoes within 1 day of harvest; and on other listed food crops up to within 7 days of harvest. Remove residues from food crops by washing before use.

| Plants                              | Insects                                                                                                                                                                                | Diseases                                               |
|-------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------|
| Tomatoes                            | Aphids, Thrips, Leafhoppers, Flea Beetles, Blister Beetles, Tomato Fruitworm, Japanese Beetle, Fleahoppers                                                                             | Early and Late Blight, Septoria Leaf Spot, Anthracnose |
| Beans                               | Aphids, Thrips, Leafhoppers, Mexican Bean Beetle, Japanese Beetle, Fleahoppers                                                                                                         |                                                        |
| Cantaloupes, Watermelons, Cucumbers | Aphids, Thrips, Squash Bugs, Immature Leafhoppers, Blister Beetles, Melon Worms, Pickle Worms, Predation of Damages, Cucumber Beetles, Squash Vine Borer, Japanese Beetle, Fleahoppers |                                                        |
| Eggplant                            | Aphids, Flea Beetles, Blister Beetles, Japanese Beetles, Fleahoppers                                                                                                                   | Phomopsis, Alternaria, Anthracnose                     |
| Peppers                             | Thrips, Flea Beetles, Fleahoppers, Japanese Beetles                                                                                                                                    | Anthracnose, Cercospora                                |
| Potatoes                            | Aphids, Leafhoppers, Flea Beetles, Blister Beetles, Colorado Potato Beetles, Japanese Beetles, Fleahoppers                                                                             | Early and Late Blight                                  |
| Strawberries                        | Aphids, Leafhoppers, Omnivorous Leaf Tiers, Flea Beetles                                                                                                                               | Botrytis Rot                                           |
| Raspberries                         | Raspberry Fruitworm, Aphids                                                                                                                                                            | Spot Blight, Anthracnose, Fruit Rot                    |

**CAUTION:** Harmful if swallowed. Avoid inhalation of dust or spray mist. Avoid contact with skin. Wash hands after use. Avoid contamination of feed and foodstuffs. Dispose of container promptly when empty.

Highly toxic to fish. Do not contaminate any body of water.

This product is toxic to bees and should not be applied when bees are active visiting the area.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

## Chevron Chemical Company

Ortho Division, P.O. Box 134117

Richmond, Va. 22181, Lafayette, La. 70501, Perth Amboy, N.J. 08861, Atlanta, Ga. 30301

EPA Reg. No. 239-555-AA

# Tomato Vegetable Dust

**CONTROLS**  
Many Insects  
and Diseases  
on Tomatoes,  
Potatoes, Melons  
and Certain Other  
Vegetables


## ACCEPTED

JAN 25, 1972

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISONS REGISTERED  
UNDER NO. 239-555-AA SUBJECT  
TO ATTACHED COMMENTS.



**Tomato  
Vegetable Dust**



**ORTHO**

\$1.79



**ORTHO**

**Tomato  
Vegetable Dust**

**Tomato  
Vegetable  
Dust**

A Multi Purpose Garden Dust That  
Controls Many Insects and Diseases  
on Tomatoes, Melons, Potatoes and  
Certain Other Vegetables.



**CONTROLS**  
Many Insects  
and Diseases  
on Tomatoes,  
Potatoes, Melons  
and Certain Other  
Vegetables

**CAUTION:** Keep out of reach of children  
See side panel for additional cautions

**NET WEIGHT 1 POUND**

**ACCEPTED**

**JAN 25, 1972**

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 37-545 SUBJECT  
TO ATTACHED COMMENTS.



**Tomato  
Vegetable Dust**