

PLEASE NOTE

This image contains more than one label approved for this product on this date.



HOME ORCHARD spray

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

INSECTS	Apple	Peach	Plum	Cherry	Apricot	Almond	Pistachio
Aphids	X	X	X	X	X	X	X
Apple Maggot	X						
Cankeryorm		X	X	X	X		X
Codling Moth		X	X		X		
Curculio	X	X	X	X			
Flea Beetle		X	X	X	X	X	X
Grape Berry Moth							X
Japanese Beetle		X	X	X	X	X	X
Leafhopper	X	X	X	X	X	X	X
Leafroller	X	X	X	X	X	X	X
Oriental Fruit Moth		X	X	X	X		
Peach Tree Borer			X	X			
Peach Twig Borer			X	X			
Pear Psylla					X		
Pear Slug	X				X		
Plant Bugs				X			
Red Spider Mites	X	X	X	X	X	X	X
Rose Chafer							X
Spittlebug							X
Strawberry Leaf Beetle							X
Strawberry Weevil							X
Sucking Plant Bugs		X	X		X		
Tent Caterpillar	X	X	X	X	X		X
Thrips	X	X	X	X	X	X	X
DISEASES							
Bitter Rot		X					
Black Rot (fruit & leaf spots)	X	X					X
Botrytis Rot							X
Botryosphaeria		X					
Black Pox		X					
Brooks Fruit Spot		X					
Brown Rot	X		X	X			
Dead Arm							X
Fly Speck		X					
Scab		X	X		X		

FOR USE AS A DUST: Apply at the rate of 1/2 pound to an average size fruit tree. Dust thoroughly so as to contact all plant parts. Repeat as necessary.

FOR USE AS A SPRAY: Mix at the rate of 5 Tablespoonfuls per gallon of water or 2 cups to 6 gallons water (14 ounces in 12 gallons water). Spray thoroughly, contacting all upper and lower leaf surfaces and fruits until entire tree is dripping wet. For medium size trees (6 to 8 ft. high), apply 2 to 3 gallons of diluted spray per tree. Large trees may require 6 gallons or more. Do not store diluted spray. Make new dilution for each use. **Mixing Directions—**Mix designated amount of ORTHO Home Orchard Spray in a little water to make a thin paste, add to required amount of water and agitate during application. Where disease control is of primary importance, use a suitable fungicide alone closer to harvest.



ORTHO

Form 5-1974


AMERICAN ROYAL CANINE COMPANY

NEW YORK, N.Y.


NEW YORK, N.Y.

NEW YORK, N.Y.

APPROX. 100 YEARS AGO...
 1900...
 1910...
 1920...
 1930...
 1940...
 1950...
 1960...
 1970...
 1980...
 1990...
 2000...



**ORTHO[®]
HOME ORCHARD
spray**



395

ORTHO® HOME ORCHARD spray



HOME ORCHARD spray



A MULTI-PURPOSE SPRAY TO CONTROL MANY FRUIT INSECTS AND DISEASES

Active Ingredients:	31.1%
Captaf	15.1%
Malathion	15.0%
Methoxychlor (as methyl ester)	1.0%
Inert Ingredients:	68.9%
Net Weight (16.25 oz. 453.6 g)	
Net Content (1.25 gal. 4.73 L)	
Equivalent to 16.25% (2.25 L) formulation	
For use on citrus fruits and other trees	
Insects and other animals	

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 16.25 POUNDS 453.6 G

ORTHO HOME ORCHARD spray

4 1/2 5

U.S. Pat. Nos. Deleted



HOME ORCHARD spray

... ..

... ..

... ..

... ..

ACCEPTED

FEB 10 1970

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CONTROL ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 239-568

NEXT

LABEL

OR DILUTE INTO FOOD OR DRINK CONTAINERS.

INSECTS	FRUITS						
	Grapes	Apples	Peaches	Plums	Pears	Strawberries	Grains
Aphids	X	X	X	X	X	X	X
Apple Maggot		X					
Cankerworm		X	X	X	X		X
Codling Moth		X	X		X		
Curculio	X	X	X	X			
Flea Beetle		X	X	X	X	X	X
Grape-Berry Moth							X
Japanese Beetle		X	X	X	X	X	X
Leafhopper	X	X	X	X	X		X
Leafroller	X	X	X	X	X	X	X
Oriental Fruit Moth		X	X	X	X		
Peach Tree Borer			X	X			
Peach Twig Borer			X	X			
Pear Psylla					X		
Pear Slug	X				X		
Plant Bugs				X			
Red Spider Mites	X	X	X	X	X	X	X
Rose Chafer							X
Spittlebug						X	
Strawberry Leaf Beetle						X	
Strawberry Weevil						X	
Sucking Plant Bugs		X	X		X		
Tent Caterpillar	X	X	X	X	X		X
Thrips	X	X	X	X	X	X	
DISEASES	FRUITS						
Bitter Rot		X					
Black Rot (fruit & leaf spots)	X	X					X
Botrytis Rot						X	
Botryosphaeria		X					
Black Pox		X					
Brooks Fruit Spot		X					
Brown Rot	X		X	X			
Dead Arm							X
Fly Speck		X					
Scab		X	X		X		

FOR USE AS A DUST: Apply at the rate of 1/2 pound to an average size fruit tree. Dust thoroughly so as to contact all plant parts. Repeat as necessary.

FOR USE AS A SPRAY: Mix at the rate of 5 Tablespoonfuls per gallon of water or 2 cups to 6 gallons water (14 ounces in 12 gallons water). Spray thoroughly, contacting all upper and lower leaf surfaces and fruits until entire tree is dripping wet. For medium size trees (6 to 8 ft. high), apply 2 to 3 gallons of diluted spray per tree. Large trees may require 6 gallons or more. Do not store diluted spray. Make new dilution for each use.

Mixing Directions—Mix designated amount of ORTHO Home Orchard Spray in a little water to make a thin paste, add to required amount of water and agitate during application. Where disease control is of primary importance, use a suitable fungicide alone closer to harvest.

APPLES AND PEACHES: Spray immediately after blossoms appear and repeat at 7 to 10 day intervals. Do not apply to apples and peaches until 15 days before harvest.

APPLES: Make first application at the first appearance of apple maggot. Repeat every 7 to 10 days until harvest. Do not apply to apples until 15 days before harvest.

PEACHES: Make one application after the blossoms have fallen and repeat at 7 to 10 day intervals. Do not apply to peaches until 15 days before harvest. When disease control is of primary importance, use a suitable fungicide alone closer to harvest.

PLUMS AND PEAR: Make one application after the blossoms have fallen and repeat at 7 to 10 day intervals. Do not apply to plums and pears until 15 days before harvest.


STRAWBERRIES, RASPBERRIES: Make one application at the first sign of new leaf growth and repeat at 7 to 10 day intervals. Do not apply to strawberries and raspberries until 15 days before harvest and raspberries until 30 days before harvest.

BERRIES, RASPBERRIES: Apply at the first sign of new leaf growth and repeat at 7 to 10 day intervals. Do not apply to raspberries until 30 days before harvest.




RICHMOND BRAND
 (Reg. U.S. Pat. & Tm. Off.)

ORTHOMERCK COMPANY, INC., NEW YORK, N. Y.
 Sole U.S. Distributor
 Atlanta, Georgia
 Boston, Massachusetts
 Chicago, Illinois
 Cincinnati, Ohio
 Dallas, Texas
 Denver, Colorado
 Detroit, Michigan
 Houston, Texas
 Kansas City, Missouri
 Los Angeles, California
 Louisville, Kentucky
 Memphis, Tennessee
 Miami, Florida
 Milwaukee, Wisconsin
 Minneapolis, Minnesota
 New Orleans, Louisiana
 New York, New York
 Philadelphia, Pennsylvania
 Portland, Oregon
 St. Louis, Missouri
 St. Paul, Minnesota
 San Francisco, California
 Seattle, Washington
 Springfield, Illinois
 Tampa, Florida
 Washington, D. C.
 Wichita, Kansas



ORTHO®
HOME ORCHARD
spray



5 2 5

ACCEPTED
FEB 10 1970
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 239-568

237
1400