



ORTHO
**Home Orchard
Spray**
ORTHO
\$1.49

Chevron


ORTHO
**Home Orchard
Spray**
**Home
Orchard
Spray**
**A Multi-Purpose Spray to Control
Many Fruit Insects and Diseases**


Controls many fruit insects and diseases. Kills on contact. Residual action. Use as directed. For more information, see the back of the container.

CAUTION: Avoid breathing of dust or spray mist. Avoid contact with skin; wash thoroughly after using.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

CAUTION: Avoid breathing of dust or spray mist. Avoid contact with skin; wash thoroughly after using.

Avoid contamination of feed and foodstuffs. Dispose of container promptly when empty.

This product is toxic to fish. Keep out of any body of water. Fish may be killed.


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

283



ORTHO Home Orchard Spray



READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

FOR USE AS A BUST: Apply at the rate of 1/4 pound to an average size fruit tree. Bust thoroughly so as to contact all plant parts. Repeat as necessary.

FOR USE AS A SPRAY: Mix at the rate of 5 Tablespoons per gallon of water or 2 cups to 6 gallons water (14 ounces in 12 gallons water). Spray thoroughly, contacting all upper and lower leaf surfaces and fruits until entire tree is dripping wet. For medium size trees (6 to 8 ft. high), apply 2 to 3 gallons of diluted spray per tree. Large trees may require 6 gallons or more. Do not store diluted spray. Make new dilution for each use. **Warning:** *Bruceana* - Mix designated amount of ORTHO Home Orchard Spray in a little water to make a thin paste, add to required amount of water and agitate during application. Where disease control is of primary importance, use a suitable fungicide alone closer to harvest.

APPLES AND PEARS: Spray when new spring growth appears and repeat at 7-day intervals up to bloom. After petal fall, repeat at 10-day intervals. Do not apply to apples and pears within 7 days of harvest.

PEACHES: Make first application at pink bud stage. Apply from 5 to 7 sprays at 10-day intervals. Peach Tree Borer - 1/4 lb. in 1 gal. water, spraying or painting trunks of trees from crotch to ground about the first of July, August and September. Do not apply to peaches within 21 days of harvest. Where disease control is of primary importance, use a suitable fungicide alone closer to harvest.

PLUMS AND PRUNES: Make one application just before blossoms open. After petal fall, repeat at 10-day intervals. Apply from 5 to 7 sprays. Do not apply to plums and prunes within 7 days of harvest.

STRAWBERRIES, GRAPES: Make first application at first sign of new leaf growth, then at 10 to 14-day intervals. Do not apply to strawberries within 3 days of harvest and remove residues at harvest by washing. Do not apply to grapes within 14 days of harvest.

CHERRIES, APRICOTS: Apply at full bloom, petal fall, 2 weeks after petal fall and again 2 weeks before harvest. Do not apply to Cherries within 7 days of harvest and to Apricots within 21 days.

Plants	Insects	Diseases
Cherries	Aphids, Curculio, Leafhopper, Leafroller, Pear Slug, Red Spider Mites, Tent Caterpillar, Thrips	Black Rot (fruit & leaf spots)
Apricots		Brown Rot
Apples	Aphids, Apple Maggot, Cankerworm, Codling Moth, Curculio, Flea Beetle, Japanese Beetle, Leafhopper, Leafroller, Oriental Fruit Moth, Red Spider Mites, Sucking Plant Bugs, Tent Caterpillar, Thrips	Bitter Rot, Black Rot (fruit & leaf spots) Botrytis, Black Fox, Brooks Fruit Spot, Fly Spots, Scab
Peaches	Aphids, Cankerworm, Codling Moth, Curculio, Flea Beetle, Japanese Beetle, Leafhopper, Leafroller, Oriental Fruit Moth, Peach Tree Borer, Peach Twig Borer, Red Spider Mites, Sucking Plant Bugs, Tent Caterpillar, Thrips	Brown Rot, Scab
Plums, Prunes	Aphids, Cankerworm, Curculio, Flea Beetle, Japanese Beetle, Leafhopper, Leafroller, Oriental Fruit Moth, Peach Tree Borer, Peach Twig Borer, Plant Bugs, Red Spider Mites, Tent Caterpillar, Thrips	Brown Rot
Pears	Aphids, Cankerworm, Codling Moth, Flea Beetle, Japanese Beetle, Leafhopper, Leafroller, Oriental Fruit Moth, Pear Psylla, Pear Slug, Red Spider Mites, Sucking Plant Bugs, Tent Caterpillar, Thrips	Scab
Strawberries	Aphids, Flea Beetle, Japanese Beetle, Leafroller, Red Spider Mites, Spittlebug, Strawberry Leaf Beetle, Strawberry Weevil, Thrips	Botrytis Rot
Grapes	Aphids, Cankerworm, Flea Beetle, Grape-Berry Moth, Japanese Beetle, Leafhopper, Leafroller, Red Spider Mites, Rose Chafer, Tent Caterpillar	Black Rot (fruit & leaf spots), Dead Arm

Home Orchard Spray

A Fine
Multi-Purpose
Orchard Spray
CONTROLS
Insects and
Diseases on
Fruits - Berries

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, Calif. Lafayette, Calif. Des Plaines, Ill. Perth Amboy, N.J. Atlanta, Ga.

Form 3025-82 EPA Reg. No. 239-568 Product 1587 Made in U.S.A.

343

