




ORTHO

ORTHO CID Seed Protection

(FUNGICIDE)
A Slurry Product — CONTAINS BINDERS
Prevents Seed Rot and Damp-off Disease



Active Ingredients: Captan 70.9%, Related Derivatives 4.1%, Inert Ingredients 25%.
N-[1-trichloromethylthio]-4-cyclohexene-1,2-dicarboximide

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panel for additional cautions.

NET WEIGHT

THE FOLLOWING TABLE gives amount of ORTHOCID 75 Seed Protectant to use on various seeds. Do not use on baby lima beans.

| SEED | Dosage Rate | Lbs. Seeds Treated % of Product |
|--|------------------|------------------------------------|
| CORN - Field | 1 1/4 oz 100 lbs | 1280 |
| | 2 3/4 oz bu | |
| CORN - Sweet | 2 3/4 oz 100 lbs | 600 |
| | 1 1/2 oz bu | |
| WHEAT | 1 3/4 oz 100 lbs | 1280 |
| RICE } For Seed and Seedling Rots Not for Smut | 2 1/4 oz 100 lbs | 710 |
| | 2 oz 100 lbs | |
| OATS } | 2 3/4 oz bu | 800 |
| | 1 1/2 oz 100 lbs | |
| BARLEY } | 1 1/2 oz 100 lbs | 1070 |
| | 2 3/4 oz bu | |
| FLAX | 2 1/2 oz 100 lbs | 640 |
| | 1 1/2 oz bu | |
| BEANS (Except Baby Limas) | 1 3/4 oz 100 lbs | 910 |
| PEAS | 1 oz bu | |
| LENTILS (Washington & Idaho Only) | 1 1/4 oz 100 lbs | 910 |
| CARROT | 6 oz 100 lbs | 265 |
| CUCUMBERS CANTALOUPE | 2 oz 100 lbs | 800 |
| SQUASH, PUMPKIN WATERMELON | 1 1/2 oz 100 lbs | 1200 |
| BEETS, SWISS CHARD | 8 oz 100 lbs | 200 |
| RUTABAGA | 3 oz 100 lbs | 530 |
| SUGAR BEETS | 8 oz 100 lbs | 200 |
| CLOVER | 5 1/2 oz 100 lbs | 290 |
| ALFALFA | 3 1/2 oz bu | |
| SOYBEANS | 1 3/4 oz 100 lbs | 960 |
| | 1 oz bu | |
| TURNIP | 2 oz 100 lbs | 800 |
| COTTON - Acid Delinted | 2 oz 100 lbs | 800 |
| COTTON - Machine Delinted Fuzzy (not delinted) | 3 oz 100 lbs | 800 |
| BLUE GRASS | 5 1/2 oz 100 lbs | 290 |
| CABBAGE, CAULIFLOWER RADISH BROCCOLI, BRUSSELS SPROUTS | 1 oz 100 lbs | 1600 |
| SPINACH, EGGPLANT | 4 oz 100 lbs | 400 |
| SORGHUM hulled (Kernel Smut) | 3 oz 100 lbs | 800 |
| PEPPERS | 2 oz 100 lbs | 800 |
| COWPEAS | 2 oz 100 lbs | 800 |
| | 1 1/2 oz bu | |
| RYE | 1 1/4 oz 100 lbs | 910 |
| SESAME | 1 oz 100 lbs | 1600 |

Chevron Chemical Company

Ortho Agricultural Chemicals Division
San Francisco CA 94119 Richmond LA 94804
Product 2006 Made in U.S.A. EPA Est. 239-MO-1
P Form 9286 B EPA Reg No 239 579 AA


Chevron

ORTHO

ORTHO[®] 75 Seed Protectant

(FUNGICIDE)

A Slurry Product — CONTAINS BINDER
Prevents Seed Rot and Damp-off Diseases



| | |
|--|--------|
| Active Ingredients | By Wt. |
| Captan | 70.9% |
| Related Derivatives | 4.1% |
| Inert Ingredients | 25% |
| N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dithione | |

CAUTION: KEEP OUT OF REACH OF CHILDREN.
See side panel for additional cautions.

NET WEIGHT

TYPE

Ortho 75 Seed Protectant is a fungicide for use on a wide range of crops. It is available in a slurry formulation.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

ORTHO[®] 75 Seed Protectant is a fungicide for use on a wide range of crops. It is available in a slurry formulation. It is used to prevent seed rot and damp-off diseases. It is used on crops such as corn, beans, and other legumes. It is used on crops such as wheat, barley, and other grains. It is used on crops such as soybeans, cotton, and other broadleaf crops. It is used on crops such as alfalfa, clover, and other forage crops. It is used on crops such as sorghum, millet, and other small grains. It is used on crops such as rice, sugarcane, and other tropical crops. It is used on crops such as tobacco, cotton, and other specialty crops. It is used on crops such as citrus, grapes, and other fruit crops. It is used on crops such as ornamentals, shrubs, and other landscape plants. It is used on crops such as lawns, golf courses, and other turf areas. It is used on crops such as trees, shrubs, and other woody plants. It is used on crops such as flowers, herbs, and other garden plants. It is used on crops such as lawns, golf courses, and other turf areas. It is used on crops such as trees, shrubs, and other woody plants. It is used on crops such as flowers, herbs, and other garden plants.

USE PRECAUTION

When using formulations that do not contain dye, to comply with 21 CFR Chapter 1, Section 2.5, all seed treated with a pesticide must be colored to distinguish and prevent subsequent or inadvertent use as a food for man or feed for animals.

DIRECTIONS

SLURRY TREATMENT — For use on a wide range of crops. It is used to prevent seed rot and damp-off diseases. It is used on crops such as corn, beans, and other legumes. It is used on crops such as wheat, barley, and other grains. It is used on crops such as soybeans, cotton, and other broadleaf crops. It is used on crops such as alfalfa, clover, and other forage crops. It is used on crops such as sorghum, millet, and other small grains. It is used on crops such as rice, sugarcane, and other tropical crops. It is used on crops such as tobacco, cotton, and other specialty crops. It is used on crops such as citrus, grapes, and other fruit crops. It is used on crops such as ornamentals, shrubs, and other landscape plants. It is used on crops such as lawns, golf courses, and other turf areas. It is used on crops such as trees, shrubs, and other woody plants. It is used on crops such as flowers, herbs, and other garden plants.

USE ISOTRA Seed Treater (75) in combination with ORTHO[®] 75 Seed Protectant when treating seed. It is used to prevent seed rot and damp-off diseases. It is used on crops such as corn, beans, and other legumes. It is used on crops such as wheat, barley, and other grains. It is used on crops such as soybeans, cotton, and other broadleaf crops. It is used on crops such as alfalfa, clover, and other forage crops. It is used on crops such as sorghum, millet, and other small grains. It is used on crops such as rice, sugarcane, and other tropical crops. It is used on crops such as tobacco, cotton, and other specialty crops. It is used on crops such as citrus, grapes, and other fruit crops. It is used on crops such as ornamentals, shrubs, and other landscape plants. It is used on crops such as lawns, golf courses, and other turf areas. It is used on crops such as trees, shrubs, and other woody plants. It is used on crops such as flowers, herbs, and other garden plants.

GLADIOLUS CORMS For Prevention of Bulb Rot — Use 1.6 oz. of ORTHO[®] 75 Seed Protectant per 100 lbs. of corms. It is used to prevent bulb rot. It is used on crops such as gladiolus, iris, and other bulb crops. It is used on crops such as gladiolus, iris, and other bulb crops. It is used on crops such as gladiolus, iris, and other bulb crops.

GRASS SEED (Range Grasses such as Wheatgrass and Bromegrass) To control *Podosporella verticillata* — Use 1.6 oz. of ORTHO[®] 75 Seed Protectant per 100 lbs. of seed. It is used to control *Podosporella verticillata*. It is used on crops such as wheatgrass, bromegrass, and other range grasses. It is used on crops such as wheatgrass, bromegrass, and other range grasses. It is used on crops such as wheatgrass, bromegrass, and other range grasses.

ONION SEED PELLETTING — Use 1.6 oz. of ORTHO[®] 75 Seed Protectant per 100 lbs. of seed. It is used to prevent seed rot and damp-off diseases. It is used on crops such as onion, garlic, and other bulb crops. It is used on crops such as onion, garlic, and other bulb crops. It is used on crops such as onion, garlic, and other bulb crops.

POTATO SEED PIECES

Do not use formulations that contain dye. For protection against damping-off use 1.6 oz. of ORTHO[®] 75 Seed Protectant per 100 lbs. of seed pieces. Suspend product in the least amount of water required to achieve uniform coverage. Potatoes should be treated immediately after cutting. If planting is delayed 1 or more days, the treated seed should be dried before storing. ORTHO[®] 75 Seed Protectant is a fungicide and will not control bacterial decay.

CAUTION:

May be hazardous if inhaled or absorbed. Avoid contact with eyes, nose, mouth, and skin. Avoid contact with clothing. Wash thoroughly after use. Do not use in areas where children or animals are present. Do not use in areas where water is used for drinking or irrigation. Do not use in areas where water is used for drinking or irrigation. Do not use in areas where water is used for drinking or irrigation.

Note to Physicians: — This product is a fungicide. It is used to prevent seed rot and damp-off diseases. It is used on crops such as corn, beans, and other legumes. It is used on crops such as wheat, barley, and other grains. It is used on crops such as soybeans, cotton, and other broadleaf crops. It is used on crops such as alfalfa, clover, and other forage crops. It is used on crops such as sorghum, millet, and other small grains. It is used on crops such as rice, sugarcane, and other tropical crops. It is used on crops such as tobacco, cotton, and other specialty crops. It is used on crops such as citrus, grapes, and other fruit crops. It is used on crops such as ornamentals, shrubs, and other landscape plants. It is used on crops such as lawns, golf courses, and other turf areas. It is used on crops such as trees, shrubs, and other woody plants. It is used on crops such as flowers, herbs, and other garden plants.

TREATED SEED DO NOT USE FOR FOOD, FEED OR OIL

For more information, contact your local agricultural extension office or the manufacturer. For more information, contact your local agricultural extension office or the manufacturer. For more information, contact your local agricultural extension office or the manufacturer.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Protect from excessive heat.

PESTICIDE DISPOSAL

Pesticide spray mixture or rinseate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL

Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

CONDITIONS OF SALE

This product is sold as a fungicide. It is used to prevent seed rot and damp-off diseases. It is used on crops such as corn, beans, and other legumes. It is used on crops such as wheat, barley, and other grains. It is used on crops such as soybeans, cotton, and other broadleaf crops. It is used on crops such as alfalfa, clover, and other forage crops. It is used on crops such as sorghum, millet, and other small grains. It is used on crops such as rice, sugarcane, and other tropical crops. It is used on crops such as tobacco, cotton, and other specialty crops. It is used on crops such as citrus, grapes, and other fruit crops. It is used on crops such as ornamentals, shrubs, and other landscape plants. It is used on crops such as lawns, golf courses, and other turf areas. It is used on crops such as trees, shrubs, and other woody plants. It is used on crops such as flowers, herbs, and other garden plants.